

US 20180177866A1

(19) United States (12) Patent Application Publication (10) Pub. No.: US 2018/0177866 A1

Pauza

(10) Pub. No.: US 2018/0177866 A1 (43) Pub. Date: Jun. 28, 2018

(54) HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY

- (71) Applicant: American Gene Technologies International, Inc., Rockville, MD (US)
- (72) Inventor: Charles David Pauza, Rockville, MD (US)
- (21) Appl. No.: 15/736,284
- (22) PCT Filed: Jul. 8, 2016
- (86) PCT No.: PCT/US2016/041456
 § 371 (c)(1),
 (2) Date: Dec. 13, 2017

Related U.S. Application Data

(60) Provisional application No. 62/190,139, filed on Jul. 8, 2015.

Publication Classification

(51)	Int. Cl.	
Č, ź	A61K 39/21	(2006.01)
	A61K 35/17	(2006.01)
	C12N 15/86	(2006.01)
	C12N 15/113	(2006.01)
	C12Q_1/06	(2006.01)
	A61P̃ 31/18	(2006.01)
(52)	U.S. Cl.	· · · ·
	CPC	461K 39/21 (2013.01); A61K 35/17
	(2013	.01); C12N 15/86 (2013.01); C12N
	1	5/1138 (2013 01) 461K 2030/5158

<i>15/1138</i> (2013.01	1); À61K	2039/5158
(2013.01); A61P 31/	Í8 (2018.	.01); C12N
2740/15043 (2013	.01); C12	N 2310/14
(2013.01); <i>C</i>	'12Q 1/06	5 (2013.01)

(57) **ABSTRACT**

The present invention relates generally to immunization and immunotherapy for the treatment or prevention of HIV. In particular, the methods relate to in vivo and ex vivo enrichment of HIV-specific CD4 T cells. In certain embodiments, the disclosed compositions and methods can incorporate therapy in order to further enhance the HIV-specific CD4 T cells.

Enrich (vaccine) and protect (AGT-LV) the HIV-specific CD4 T cells.



Enrich (vaccine) and protect (AGT-LV) the HIV-specific CD4 T cells.



Figure 1



Figure 2



Figure 3

miR21 Vif end miR185 Tat start miR185 Tat end CCGCGTCTTCGTCGCGGCCGCTCGAGCATGCAT

Figure 4

TTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGGGCGCCGTCCAGGCACCTCGATTAGTTCTCGAG GACTGAAGTTAGGCCAGCTTGGGCACTTGATGTAATTCTCCCTTGGAATTTGCCCTTTTTGAGTTTGGATCTTGGTTC ATTCTCAAGCCTCAGACAGTGGTTCAAAGTTTTTTTCTTCCATTTCAGGTGTCGTGAGGAATTGGCGAAGCTAATTC miR30 CCR5 start TGCAGATTCGACTGTACAAGGTATATTGCTGTTGACAGTGAGCGACTGTAAACTGAGCTTGCT CTACTGTGAAGCCACAGATGGGTAGAGCAAGCACAGTTTACCGCTGCCTACTGCCTCGG miR30 CCR5 end miR21 Vif start <u>ACTTCAAGGGGCTT</u>ccccGG<u>CATCTCCATGGCTGTACCACCTTGTCGGGGGGATGTGTACTT</u> CTGAACTTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGGTA

ACCGGTGCCTAGAGAAGGTGGCGCGGGGGGAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCTTTTTCCCGAG GGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTCGCAACGGGTTTGCCGCCAGAACA CAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTGCGTGCCTTGAATTACTT TGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAAATTTTTGATGACCTGCTGCGAC SCGGGCGGCGACGGGGCCCGTGCGTCCCAGCGCACATGTTCGGCGAGGCGGGGCCTGCGAGCGCGGCCACCGA CCT6GGCGGCAAGGCTGGCCCGGTCG6CACCAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCA

Elongation Factor-1 alpha (EF1-alpha) promoter (ITALICS)





Figure 5



Figure 6



Figure 7



B



Figure 8





1 atgcagacee teatetttt egacatggag gecaetgget tgeeettete eeageeeaag
 61 gteaeggage tgtgeetget ggetgteeae agatgtgeee tggagageee eeeeettet
 121 eagggeeae eteeeaegt teeteeaea eegegtgtgg tagacaaget eteeetgtgt
 181 gtggeteegg ggaaggeetg eageeetgea gecagegaga teaeaggtet gageeeaget
 241 gtgetggeg egeetggegg teaatgtttt gatgacaace tggegaee egeetagge
 361 teeeeetge gecageeag geetggegtg etggegeee eeaggeggg egeetgggggee
 361 teeeeetge teeaageag getggetag geetggeeta eeagtet gatggigee
 361 teeeeetge teeaageag getggetag geetggegg gageageag eeeteaga
 481 eaeggeeeaa ggaaggeta eegeetagge ageetggeg gageageag eeeteaga
 481 eaeggeeeaa ggaaggeta eageetage ageatetae etggegete

TREX1 - human; open reading frame; three prime repair exonuclease 1

541 octocagaci ogcacaoggo tgagggtgat giootggood tgotoagoat ofgicagtgg 601 agaccacagg occtgotgog gigggtggat gotcacgooa ggoolllogg caocatcagg 661 occatgtatg gggtcacago ototgotagg accaagocaa gaccatotgo igtcacaaco 721 actgoacaco iggocacaao caggaacaci agioccagoo itogagagag caggggiaco 781 aaggalotto olocagigaa ggacoolgga gooclatoca gggaggggol goiggococa 841 olggiolgo iggocatool gacollgoa giagocacao iglaiggaol alcoolggo 901 acacolggog allaa

Figure 9

TREX2 - human; open reading frame; three prime repair exonuclease 2 -1 cacettygic acaetecaga gologettea tececaogag eggeacetga telecagtga 61 oggggagaot gaggootggc aaagaggaoo tgttgcottt gtgtaactgg coccagcatg 121 gaggaggeat typecchage atgggggagg cattyaccee aggeetgeea ggeetgeag 181 ecacageota getaggigga gattactuce ticegeacea etetagette eteceegige 241 ctgtgaagag ologgggoot gettootaat ttgtaaacae ggggegtgtg teteagigge 301 tytyagetag ogagggggtg gegagegage eggetgegea ggteetgagg ecceaggeet 361 cattgttggc caacaggcag ctgggggggg gotgoggcog otgattaaag geogootaga 421 geagestigty tiggegalage tigeocagalage cocaggalage eggteagtige ecgeceelagt 481 octoagggtt tgtgcototo gotoggacag tttgaggaot tgotatocco gtgggaacat 541 caccatetee gaggeaceee gggoegagae ottigtotte etggaeeigg aageeaeigg 601 getecceagt gtggageeeg agattgeega getgteeete tttgetgtee acogeteete 661 octgyagaac coggagcacy acgagtotgg tgeoctagta ttgeocoggg tootggacaa 721 getcaegetg tgeatgtgee eggagegeee etteactgee aaggeeageg agateaeegg 781 cetgageagt gagggeetgg egegatgeeg gaaggetgge tttgatggeg eegtggtgeg 841 gaogetgeag geotteetga geogeeagge agggeeeate tgeettgtgg eccaeaatgg 901 ottlgattat gaittseece tgetgigs egagetgegg egeetgegig eesgeelgee 961 cogggaract gtotgcotgg acaegetger ggcotgegg ggestggaee gegeeeasag 1021 ccacggcace egggceeggg geegeeaggg ttacageete ggcageetet tecacegeta 1081 ottoogggoa gagoosagog cagoosacto agoogagggo gaogtgoaca cootgotoot 1141 gatetteetg cacegegeeg cagagetget egeetgggee gatgageagg coegtgggtg

Figure 9 Con't

1201 ggcccacate gagcccatgt acttgccgcc tgatgacccc agcctggagg cctga

SAMHD1 - human; open reading frame; SAM domain and HD domaincontaining protein

1	attgcgcctg	cgcagggagc	ccaaggcaag	agccgctagg	ctgeeetgee	cgaagggctc
61	aactgtcagt	gagootgogo	aggaggccaa	taggetgeea	atactecttg	gacteccoge
121	cagggccctg	ctgtcagtgc	geetgegege	gggtcoggog	ccgaggttct	tgactgctgt
181	geoggaegee	aggtgtagec	atgeagegag	ocgattcoga	geageoetee	aagogtocco
241	gttgcgatga	cagcecgaga	accoccteaa	acaccectte	cgcagaggca	gactggteec
301	cqqqcctqqa	actecatece	gactacaaga	catggggtcc	ggagcaggtg	tgotecttee
361	tcaggegegg	tugetttgaa	gageoggtoe	tgetgaagaa	catecgagaa	aatgaaatca
421	caddodcatt	actoccttot	cttgatgagt	ctcattttaa	aaatcttoqa	gtaagtteet
481	taaaqaaaa	gaagaagetg	cttaqttata	tecaqegatt	quitcasate	cacattoata
541	caatgaaggt	aattaatuat	ectatecato	accacattaa	actocaccot	ctecteatee
601	gaatcattga	tacaceteaa	tttcaacotc	trogatacat	caaacagetg	agaagtagtt
661	actatotttt	tocaquaget	teseseate	gatttgagga	tagtetaggg	ataggatate
723	tagcaggatg	tctagttcac	acactagata	aaaaacaacc	agagetgeag	ataadtaaac
781	aanatattet	stataticaa	attortosac	tttatcatoa	teteooteat	gaaccatttt
841	ctcacatott	taataaacaa	tttattcrac	ttarteacec	ngagataaaa	tagacacata
961	aacaacacto	pattatasta	tttaaaracc	ttattaatte	teatoneatt	aarectotea
461	togeaceate	taateteste	cctgaggagaa	atatttartt	tatagancas	caaattotan
1621	naccanttra	atrarctoto	raaratteat	totoccata	taaannarat	ontrossona
1021	aaanottoot	tratranata	matriala	asagaaatga	caitastata	narasatinna
1131	attattttor	carogagaca	cotratetto	aaagaaa cgg	taattttat	taraadoort
1201	thattaantt	taccotata	totacotto	acastroatt	aontattat	antanguguu
1041			- vy vy say voy - montheatra	acaacyayee aarabaaraa	gegeareege	anan anananya.
1301	- aggaaggeegg	addoolgcao	attattata	nostrattor	anatootitic	ntroppareara
1003	obcochecob	ageoggeaas	actactigata	aaaaaaaba	togocyccicco togocyccicco	secondageng
7887	acyaccacac	agagattata	- AA cAccAAaA	yaaaaaayta hhhhhh	tataanantt	acaycaaccy.
18031	- acyacacyya	ageocacace	aagorgacay	ttttccccc.	asttanotae	ant at the state
1001	- tragatotaa	arrigadagar	- guaugagaga	- 20000aaaaaaaa	aattoooaa	ageaactice.
1001	-scaaycacyc	- Addradard	- cagooaacag - accostorto	-gauaaaaaaaaa	gallaaagg offarture	gayyaccacy. massoctas
1601		titistanta	entetteten	accelacage	tanostanco	gegaaacega maaaansotn
1781	-ayyooyaaya -rasttaatra	tettaeayug	-gaugecacea Fottatoono	acacygacca mhannana	- cyyaacycaa - commensite	gaaaayaaco
1005	- case cyacea - secoranat	- 69020032000 - ****	- saucycaaga	astttanens	acoactest	aggattatatta
10023	aaaaaaaayyu		- ant that at a	aaccegeaga ooxoooooo	gradercaare	agagiacacc.
1001	gcaagaaggc	yyarayaway tttoroonn	ayuuyuaty	- Cogcadyaca 	avacturger	tagtggtgtg
1001		atayaaaaaaa	- CCycayya Cy - Saf Saf Sasa	gegaegetae teensistee	agottotacto	acaacacocc
00.83		acygaacyac	ayuactucay Attactor			yayaayaay
2031	toreadageag	ageocageee	CCCadayary	acceaacycy	aacytotyta	geoageogee
- 2.2.V.2 - 3.2.C.2	- Labadactici:	- Subbabbabbab - Arthorna babbaba	- Caab Coat Club	ayayyuuuda taastatsaa	accalayaat	LULYLAAALL Sachaaataa
2101	aaugacaacu abbbbbcabaa	Catycuttaa	- LLLLGLALLL - startstart	Lgaacytata	- cgcauge cga	agoldagiaa
- 2222 - 2225	- LULULAALUA	aayaaacaay	alggiactag	guadaluura.	Claracially	addaycatta
- 62 6 2 - 9 2 2 3	an en ordere teter.	- UUUNAAUUUU	du lanagu lu.	- An An An An	geogeneare.	sestedydd. Taaraatte
2042	aacaaccee		tyddululau toosoffto	- CECELEELE	CCCCadgagaga	taagguttug
2402 2723	- cooligicoyou - analanataa	- cago cogyac - ataratata	artostatas	- cuyyyuccaa - orazoooo	guyauuuuu	- Seglere is to de iglierte - anne a antie to de iglierte -
24.01	coccaaycay		agecargege	Caccacacco.	agectactee	gaaceeeaa
- ఓచెచి. - నకరిగి	gacagacgac	testortege	or compared	yaaacyayac finanaacoon	aaggggtataa	attygaatta
2302	gaaacuyata	- Lugicia Luga	- Caadcacyyo	todddduddd	dudludddu	
20912	gaagageeee	- Cugudaduda - atagonationa	aayuuuyayy	acaycoccoc	dagagugacu	caaguacaag
2791 19965	-uyaaayaudo -ttaasootee	tagaaraa	-cossuagee	- uuuuauaagg - aacaabaata	- cyccaggreg	yyycaaayay
2/01		- cyyccayyay	- Sygunu Coda - Angerana	Largergeg anatamana	ayayyuuuya	trayyttty tatesta
LOZI 7001	yyyaayarct	-gagageeeeg	geessteege name#anam	gayuyyyaaca axxaabaaaa	caaagergee tatazzaten	- LULUGUUGUG ataaataaaa
2001	ayacoosacc	-coagegacag	aggaaroooo	taaattayyt taatta	- cy cy cc cc yy	concercited and
5003 5347	-yeucoayace -ttooootae	- ogggtægtgo		- uyaauduuud - aabaalaabaa	- CCCCCCCCGGC - A A A A A A A A A A A A A A A A A A A	tertereses
2022	- uuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuu	- ccaycecyda - accateoraa	gaaggeeree		- uuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuuu	erereddadad.
33.01	tugguuayay	guuacuacay atamatété	yearbailer Foresthere	cagageeeee	Layaccuyyyy	auceeyeece
الدائنة الداقية	- CACERATE COLOR	y systatety	- vy ca ce cady	aycaadtatt	ayrrayyryr	- gguggugugu

MxA - human; open reading frame; influenza virus resistance 1

1 cgagcagaaa tgaaaccgaa actgaatigt ccgggaaatt cgcggtgggg gcggagagcg 61 cagggagaag taagcccagt gcaggateet gaggecegtg titigcaggae cagggeegge 121 ettoogatto occatteatt ecaqaaqeas eqaaceaeqe tgtgseegga teecaagige 191 ageggéacee agegtggges tggggtfgee ggttgaceeg gtreteages tggtageaga 241 ggccaggcca gtgccacaag gcacctaagt ccacctgggc ctggagcagg acaggttgca 301 aaagaaaata totogggaco oocaaactoo ttatgotaag ggasacatog agootgggaa 361 elgagecate accycloces thelittee caacegaac cetyligtes gaggiaesee 421 cadageaact ceacaceggg tgeatgeeae ageaacteea tettaaatag gagetggtaa 481 aacqaqqctq atacctactq qqctqcattc ccaqacqqca taqcqaqqaq qtqctqaaqa 541 gegéaggtti ggagaatgat éacciggatt ggaaccatag cictaccaat atggaaccea 601 geteeltagu eeleggtett eteatggaga acatggigig alaateetae teeteiggga 661 gggtggetgt taageettyg acegeagttg coggeeagga ateceagtgt caeggtggae 721 acycotocot cgogocottg cogoccacot gotcaccoag ctcaggggot ttggaattot 781 giggccacae tocoaggaga toggitetog gieggagget acaggaagae toccaeteee 941 tgaaatotgg agtgaagaac geegecatee agecaceatt ccaaggaggt geaggagaac 901 agetetetea taccalltaa ettetteaca tiaettttat tteaaggaac etalattaga 961 gottactting caaagaagga agatggtigt thoogaagig gacatogcaa aagotgatoo 1021 agetgetgea teocaccete tattaetgaa tggagatget aetgtggeee agaaaaatee 1081 aggeteggtg getgagaaca acctgtgeag ceagtatgag gagaaggtge geeeetgeat 1141 caacotcatt gactecetge gggetetagg tgtggageag gacetggeee tgecageeat 1201 cycogicate gyggaecaga getegggeaa gageleegty liggaggeae igteaggagt 1261 tyccotteec agaggeageg ggategtgat cagatgeeeg etggtgetga aaetgaagaa 1321 acttgtgaac gaagataagt ggagaggcaa ggtcagttac caggactacg agattgagat 1391 tteggatget teagaggtag aasaggaaat taataaagee cagaatgeea tegeegggga 1441 aggaatggga atcagicatg agctaatcac octggagatc agctocogag atgtocogga 1501 telgacteta atagaeette etggeataae cagagtgget gtgggeaate ageetgetga 1561 cattyygtat aagatcaaga cactcatcaa gaagtacatc cagaggcagg agacaatcag 1621 cotggtggtg gtocccagta atgtggacat ogccaccaca gaggototca gcatggocca 1681 ggaggtggac coogagggag acaggaccat oggaatottg acgaagootg atotggtgga 1741 caaaggaact gaagacaagg tigtggacgt ggtgeggaac clegtgttee acctgaagaa 1801 gggttacatg attgtcaagt geoggggeca geoggagate eaggaceage tgageetgte 1861 cysagoocty cagagagaga agatettett tyagaaccae ceatatttea gggatetget 1921 qqaqqaaqqa aaggeesegg tteeetgeet qqeagaasaa ettaccageg ageteateae 1981 acatatetyt aaatetetye eeetyttaga aaateaaate aaggagaete aeeagagaat 2041 aacagaggag ctacaaaagt atggtgtoga catacoggaa gaogaaaatg aaaaaatgtt 2101 ottootgata gatasagtha atgootttaa toaggacato actgototoa tgoaaggaga 2161 ggaaactgta ggggaggaag acattegget gtttaccaga cteegacaeg agttecacaa 2221 atggaytaca ataattgaaa acaattttca agaayyccat aaaattttga gtagaaaaat 2281 ccagaaattt gaaaatcagt atcgtggtag agagetgeca ggetttgtga attacaggae 2341 attigagada aleqigaaad aqeaaateaa ggeaelqgaa gaqooqqeig iqqataiqei 2401 acaceecgty acggatatyg tooggettge ttteacagat gtttegatea aaaattiiga 2461 agagtttttt aacotocaca gaacogocaa gtocaaaatt gaagacatta gagcagaaca 2521 agagagagaa gutgagaago tgatcogoot coacttocag atggaacaga ttgtotactg 2581 ccaqqaccaq qtatacaqqq qtqcattqca qaaqqtcaqa qaqaaqqaqc tqqaaqaaqa 2641 aaagaaqaag aaateetggg attitigggge titecaqtee ageteggeaa caqaetette 2701 catggaggag atotticago acotgatggo clatcaccag gaggocagoa agogcatoto 2761 cagecacate cettigates tecagitett catgetecag acgiacggee ageagettes 282) gaaggocatg ctgcagetee tgcaggacaa ggacacetae agetggetee tgaaggageg 2881 gagegacade agegacaage ggaagtteet gaaggagegg ettgeacgge tgaegeagge 2941 trggegergg ettgercagt terreggtta accaractet gteragreer gtagaogtge 3001 acquaeacty totycccccq ttoccqqqta gccactqqac tgacqactty aqtyctcagt 3061 agreagacty gatagreegt storgettat cogttageog tggtgattta geaggaaget 3121 gtgagagcag titiggtitet ageatgaaga cagageeeea eeeteagatg cacatgaget 3191 ggegggatty aaggatgetg tettegtaet gggaaaggga tttteayeee teagaatege 3241 Eccacettge agetelecce ttetetgtat tectagaaae tgacacatge tgaacateae

1	aagagatgat	ttotocatec	tgaacgtgca	gegagettgt	caggaagate	ggaggtgeea
51	agtagcagag	asagcateee	ccagetetga	cagggagaca	geacatgtet	aaggeeeaca
121	ageottggco	ctaccggagg	agaagtcaat	tttetteteg	aaaataootg	aaaaaagaaa
181	tgastteett	coagcaacag	ccacogocat	toggcacagt	gccaccacaa	atgatgtttc
241	otocaaactg	geagggggea	gagaaggacg	otgettteet	ogocaaggac	ttcaactttc
301	teactttgaa	caatcagcca	ccaccaggaa	acaggageea	accaaggggca	stggggcccg
361	agaacaaeet.	gtacagecag	tacgagcaga	aggtgegeee	ctgeattgac	ctcatcgact
421	ccatgoggga	totgggtgtg	gagcaggacc	tggcootgaa	agocatogeo	gteategggg
481	accagagoto	gggcaagagc	totgtgetgg	aggeactgte	aggagtogog	otteccagag
541	gcagcggaat	cgtaaccagg	tgteegetgg	tgetgaaact	gaaaaagcag	occtgtgagg
601	catgggoogg	aaggatcage	taccygaaca	cogagotaga	getteaggae	cctggccagg
661	tggagaaaga	gatacacaaa	geccagaaeg	teatggccgg	gaatggoogg	ggcateagee
721	atgageteat	cageetggag	atosoctocc	etgaggttee	agacetgaec	atcattgacc
781	ttoccggcat	caccagggtg	getgtggaea	accagecceg	agacatogga	ctgcagatca
841	aggeteteat	caagaagtac	atceagagge	agcagacyat	caacttggtg	gtggtteect
901	gtaacgtgga	cattgecacc	acggaggege	tgagcatggc	ccatgaggtg	gacccggaag
961	gggacaggac	catoggtate	ctgaccaaac	cagatetaat	ggacaggggc	actgagaaaa
1021	gegteatgaa	tatagtacaa	aacctcacqt	accocctcaa	gaagggctac	atgattgtga
1081	agtgcegggg	ccagoaggag	atcacaasoa	ggetgagett	ggoagaggca	accaagaaag
1141	aaattacatt	ctttcaaaca	catccatatt	tcagagttet	cctggaggag	gggtcageca
1201	aggtteeeag	actggcagaa	agacttacca	ctgaactcat	catgeatate	caaaaatege
1261	tcecgttgtt	agaaggacaa	ataagggaga	gecaccagaa	ggegaeegag	gagetgegge
1321	gttgegggge	tgacatecce	agcosggagg	cogacaagat	gttettteta	attgagaaaa
1381	teaagatgtt	taatcaggac	atogaaaagt	tagtagaagg	agaagaagtt	gtaagggaga
1441	atgagacceg	tttatacaac	aaaatoagag	aggattttaa	aaactgggta	ggcatacttg
1501	caactaatac	ccaaaaagtt	aaaaatatta	tecacgaaga	agttyaaaaa	tatgaaaago
1561	agtatogagg	caaggagett	ctgggatttg	tcaactacaa	gacatttgag	ateategtge
1621	atcagtaoat	ccagoagetg	gtggagcoog	occttageat	geteeagaaa	gccatggaaa
1681	ttatecagea	agetttcatt	aacgtggooa	aaaaacattt	tggogaattt	tteaacetta
1741	accasactgt	tcagagcacy	attgaagaca	taasagtgaa	acacacagea	aaggcagaaa
1801	acatgateca	acttcagttc	agaatggagc	agatggtttt	ttgtcaagat	cagatttaca
1861	gtgttgttet	gaagaaagto	cgagaagaga	tttttaacce	tatggggaog	cottoacaga
1921	atatgaagtt	gaactctoat	tttcocagta	atgagtette	ggttteetee	tttaotgaaa
1981	taggoateea	cotgaatgoo	tacttettgg	aaaccagcaa	acgletogee	aaccagatee
2041	catttataat	teagtatttt	atgeteegag	agaatggtga	ctccttgcag	aaagccatga
2101	tgcagatact	acaggaaaaa	aatogotatt	octggotgat	tcaagagcag	agtgagacog
2161	ctaccaagag	aagaatoott	aaggagagaa	tttaccggct	caotoaggeg	ogacacigoao
2223	totytoaatt	ctocagcaaa	gagatecaot	gaagggogge	dardoordra	gttgttttot
2281	tgtgcgtact	cattoattet	aaggggagtc	ggtgeaggat	geogettetg	otttggggoo
2341	aaactettet	gtcactatea	gtgtccatct	ctactgtact	ccatcagcat	cagageatge
2401	ateaggggte	cacacagget	cagetetete	ceccaccag	etetteeetg	acctteacga
2461	agggatgget	otocagtoot	rdddroeddr	ageacacagt	tacagtgtoc	taagatactg
2521	ctatcattet	togetaattt	gtatttgtat	teeetteece	otacaagatt	atgagacccc
2581	agagggggaa	ggtetgggte	aaattottot	tttgtatgtc	cagtetootg	cacageacet
2641	gcagcattgt	aactgettaa	taaatgacat	otcactgaac	gaatgagtgo	tgtgtaagtg
2701	atggagatac	etgaggetat	tyctcaagoo	eaggeettgg	acatttagtg	actgttagco
2761	ggtecettte	agatecagtg	gocatgeoco	otgetteeca	tggttcactg	teattgtgtt
2821	teccagoote	teosotecce	ogccagasag	gagcotgagt	gattetettt	tertettett
2831	recetgatta	tgatgagett	coartgttot	gttaagtott	gaagaggaat	ttaataaago
2941 aaag	aaactt tttaaaaa	acg t				

MxB - human; open reading frame

APOBEC3G - human; open reading frame; apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3G (APOBEC3G)

1 gtudictiget geoteageot ggtgtggadd caeoleecgd gegetggetg caatgaettt 61 etetttoeet tigeaatige etigggteet geegeacaga geggeeigte titateagag 121 gtccctctgc cagggggggg gccccagaga aaaccagaaa gagggtgaga gactgaggaa 181 gataaagogt cccagggcot ootacaccag ogoctgagoa ggaagoggga ggggocatga 241 otacqaqqoo ctqqqaqqtc actitaqqqa qqqctqtoot aaaaccaqaa qottqqaqca 301 gaaagtgaaa cootggtgot coagacaaag atottagtog ggactagoog gocaaggatg 361 aagootoact toagaaacao agtggagoga atgtatogag acacattoto otacaacttt 421 tataatagac ceateettte tegteggaat accgtetgge tgtgetaega agtgaaaaca 481 aagggteeet caaggecooc tttggacgea aagatettte gaggecaggt gtatteegaa 541 ottaaqtace acceaqaqat qaqattotte cactqqttea qeaaqtqqaq qaaqetqeat 601 egtgaceagy agtatgaggt eacetggtae atateetgga geeeetgeae aaagtgtaea 661 agggatatgg ccacgtteet ggeegaggae eegaaggtta ceetgaecat etttgttgee 721 egectetaet acttetggga eccagattae eaggaggege ttegeagect styteagaaa 761 agagaoggic cgcgtgccac calgaagate atgaattalg acgaalitea geactgttgg 841 ageaagtteg tytacageea aagagageta tityageett gyaataatet geetaaatat 901 tatatattac tycacatcat gotgggggag atteteagac actegatgga teeaceeaca 961 tteaetttea aetttaacaa tgaacettgg gteagaggae ggeatgagae ttaeetgtgt 1021 tatgaggtgg agogcatgoa caatgacacc tgggloctgo tgaaccagog caggggottt 1081 ctatocaaco aqqctcoaca taaacacqqt ttoottgaaq qooqccatqo aqaqctqtqc 1141 ttootggacy tgatteeett ttggaagetg gaeetggace aggactaeag ggttaeetge 1201 theaceteet ggageceetg etteagetgt geecaggaaa tggetaaatt eattteaaaa 1261 aacaaacacg tgagcotgtg catottcact goocgcatot atgatgatca aqqaagatgt 1321 caggagggge tgcgcaccet ggccgagget ggggccaaaa tttcaataat gacatacagt 1361 gaatttaage actgetggga caeetttgtg gaccaceagg gatgteeett eeageeetgg 1441 gatguactag atgagcacag ccaagacetg agtgggagge tgegggeeat tetecagaat 1501 caggaaaact gaaggatggg ootcagtoto taaggaaggo agagacotgg gttgagooto 1561 agaalaaaag alottettoo aagaaatgea aacaggetgt teaccaceal eteeagetga 1621 teacagacac cagcaaagea atgcacteet gaceaagtag attettttaa aaattagagt 1681 goattacttt gaatcaasaa tttatttata tttcaagaat saagtactaa gattgtgete 1741 aatacacaga aaagttteaa acctactaat eeagegacaa tttgaategg ttttgtaggt

1801 agaggaataa aatgaaatac taaatctttc tgtaaaaaaa aaaaaaaa

TRIM5 alpha - human; open reading frame; tripartite motif containing 5 (TRIM5)

ž.	agtttatctt	teacttteet	gccctgagtg	tgagcaagaa	tttcctgcgg	ttoctotagg
61	assatteett	tgtgcagate	aggcccgtgg	attggtgagt	gaateetaae	cacgtottcc
121	ctggcctgtc	ttcactcttc	tecceagaat	caccacttet	gcactggtgt	ctgaaggtgt
181	attgagtgat	tttgtggagg	gcagaagtag	gaagtetttg	ggacaaaact	gtatttacct
241	tgggatetgt	gaacaagagg	aacctcagca	gccaggacag	gcaggagcag	tggaataget
301	actatggett	ctggaatect	ggttaatgta	aaqqaqqaqq	tgacetgeec	catctgcctg
361	gaacteetga	cacaaccoct	gageetggae	tgeggeeaca	gettetgeca	ageatgeete
421	actocaaaco	acaaqaaqto	catgetagae	aaaqqaqaqa	gtagetgeee	tatatacega
481	atcaqttacc	agectgagaa	catacqueet	aatoggcatg	tagécaacat	aqtqqaqaaq
541	cteadqqaqq	teaagttgag	cccadadqqq	cagaaagttq	ateattqtqc	acqccatqqa
601	gagaaactte	tactetteta	tcaggaggac	gggaaggtca	tttggtaget	ttataaacaa
661	teteaggage	accotostca	ccacacotto	ctcacagagg	aggitaccog	ggagtaccaa
721	atuaaoctee	aggeagetet	ggagatocto	aggeagaage	adcaggaadc	tgaagagtta
781	gaagetgaca	tcagagaaga	gaaagettee	togaagacte	aaatacacta	tuscasasco
843	sacotettoo	cagattttga	acsactoana	gacatectog	actosoadda	asocastaso
901	otocaseseo	tagagagagaga	ggaggaagaa	attetosasa	accttacaaa	ctctraaact
961	gagatogtoc	adescaceca	atecetaada	aarcteatet	cagatotogaa	acategadate
1821	09000000000000000000000000000000000000	tostugaact	acttragaat	atastaaca	tcataaaaaa	
1021	atasettas	sessareses sessareses	aarttttora	asssatrasa	aaanantatt	tensactort
1141	astetessa	agaagaaga asstartsas	antotttana	nadotroran	atotecosco	etacteentt
3701	gatetegaaag	gaacystaga toorteess	rearetttre	gagorgacag tatantatas	tttrtassas	taaraaaaaa
2043	gaugegauag ataraatata	oggesseaaa	A A A A A A A A A A A A A A A A A A A	agago sy csa agago sy csa	aanaanaanaa	
1001	gegageeeee	- uyaaaccaca	gacaacacac	gggggcacgag	ggacaagaca	00003000000
1001	yuyaaseesa taabamasaa	accacogcac	- ogguarde og	ggott caaa	gua cuacaeco transmototra	ayyyaaacac
4301	-cacegggagg	Lagacytytt.	Ladyaaaauu Anabhaaaaa	georgaacee	-ugggggtatg	- Cycliggescu
1441	Caaccugaty	caauguguaa	lattgaaaaa	aatgaaaatt	accaacctaa	atacggctac
2001	- Egggelalag	ggilagga	aggagutaaa	tgtagtgott	tocaggatag	LUCCLUCCAL
1061	actected	teccecteat	racaccece	ccegegacea	rtrgscorga	regrarrada
1021	gttttcctag	actatgagge	ergeacegie	Cattorica	atatcacaaa	ccatggattt
1001	createrata	agtiticea	cugrucereur	rereageerg	tatttccata	tttaaateet
1/41	agaaaatgtg	gagueccoat	gactotgtgc	teaceaaget	ettgaacett	cttacacact
1901	cagoocerte	tgracageae	ctcttgtcca	ggtgcatete	atacacctga	actestinge
1861	atcatttaa	coatettte	errgergrer	contettte	tarttgaacg	rectteacte
1921	atcagtaaaa	tgtaataatt	geettgtgee	atattgtccc	caatatttta	ttgacatttg
1981	atagcaattt	ttttcatcat	tttccgtact	cctaaggaaa	actgacetat	acctcataaa
2041	atgagacogc	tatttaggta	ttacttctge	cagatattta	tcacccaatt	geetetgaca
2101	otgactaaga	agatgaagaa	aagettttea	acagcottto	tatateateg	tgtgataatt
2161	gttcaccaat	gaatgagtee	ttageeetgt	gtcagtttac	cetegatgee	cttatttgtg
2221	agttaaagag	aaaatatcat	aaatggtata	ctettaagta	tagaggtttt	gtatctagag
2281	gatetcagtt	esacteetgt	ctctccatat	accagcagtg	taactgtgaa	taacatactt
2341	aaatggctgt	gettatttee	ttttotttc	tttttttttt	tttttttt	ttgagatgaa
2401	gttttgctct	tgttccccag	getggagtge	aatggcacga	totoggttoa	ctgcaacetc
2461	cacctctcag	attcaagcaa	tteteetgee	teagectece	aagtagetgg	gattacaggt
2521	geccaccacc	accoctgget	aaatttgtat	tttcagtaga	gacggggttt	ccccatgttg
2581	gttaggeteg	tctagaacct	otgacctcag	gtgatecace	agaataygaa	teccaaagtg
2641	ctgggattac	aggogtgagc	cacggegeee	agcetgtget	tattttctta	aaataatttt
2701	tgtattaaaa	acttoacatt	aaataagtgo	taatgtttta	ttgcatagta	gggtgactag
2761	agttaacaat	aacctattge	atatattttg	aaatagctag	aagagaggat	tttgaaagtt
2821	ctoaacacaa	agaaatgaca	catatttgag	gtgatggata	tgetaattae	cctggttegg
2881	ttattacqca	atgtatacat	gtatcasaac	atcacactgt	accacataaa	tatgtatatt
2941	tattattiqt	caattaaaaq	caaaatsaaa	casaaaacct	toatctaata	ctttqqatca
3001	ttatqaaaaa	ataaatteet	gaagtataaa	geatetatet	aaqtqtcttq	atctaataag
3061	tactfottet	acaaattatt	gaaaaacata	aactototta	atqtctcatq	gaacaggttg
3121	tgeetteagg	qaaactaqqa	ttogatttac	taaattetea	tttttadat	ctcagatact
3181	actotcaasa	tgactteaat	tetacettet	atatataata	cacacatata	tttaggattt
		-				

Tetherin (BST2) - human; open reading frame 1 ataaaggggt ggeeegtaga agatteeage acceteeeet aasteeagge cagacteett 61 tragetasag gggagatotg gatggcatot acttegtatg actattgcag agtgeccatg 121 gaagacgggg ataagegetg taagettetg etggggalag gaattelggt geteetgate 181 atogtgatte tgggggtgee ettgattate tteaceatea aggeeaacag egaggeetge 241 egggaeggee tregggeagt gatggagtgt egeaatgtea eccateteet geaacaagag 301 styaccogage sceagaagge stttcaggat gtggaggsoc aggesgecas stgcaascas 361 actgigatgg cootaatggo ticcotggat goagaaagg cocaaggaca aaagaaagtg 421 gaggagettg agggagagat cactacatta aaccataage tteaggaege gtetgeagag 481 gtggagogac tgagaagaga aaaccaggto ttaagogtga gaatcgogga caagaagtac 541 taccoraget eccaggacte rageteeget geggegeeee agetgetgat tgtgetgetg 601 ggootcagog ctotgotgoa gtgagatooc aggaagotgg cacatottgg aaggtoogto 661 etgetegget titegettga acatteeett gateteatea gitetgageg ggteatgggg 721 caacacqqtt aqcqqqqaga qoacqqqqta gccqqaqaaq qqcctotqqa qoaqqtctqq 761 aggggocalg gggaagteet ggglgtgggg acacagtegg gttgacecag ggelgtetee 841 otocagagee tecotorgga caatgagtee eccetettgt eteocaceet gagattggge 901 atggggtgcg gtgtgggggg catgtgctgc ctgttgttat gggttttttt tgcgggggg 961 gttgettttt tolggggtot ttgageteca aaaaataaae actteetttg agggagagea 1021 cacctgaaaa aaaaaaaaaa aaaaaaaa

HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY

CROSS REFERENCE TO RELATED APPLICATION

[0001] The present application claims priority to U.S. Provisional Application No. 62/190,139, filed on Jul. 8, 2015, the disclosure of which is specifically incorporated by reference.

FIELD OF THE INVENTION

[0002] The present invention relates generally to the field of immunization and immunotherapy for the treatment and prevention of HIV. In particular, the disclosed methods of treatment and prevention relate to the administration of viral vectors and system for the delivery of genes and other therapeutic or diagnostic compositions.

BACKGROUND OF THE INVENTION

[0003] Combination antiretroviral therapy (cART) (also known as Highly Active Antiretroviral Therapy or HAART) limits HIV-1 replication and retards disease progression, but drug toxicities and the emergence of drug-resistant viruses are challenges for long-term control in HIV-infected persons. Additionally, traditional antiretroviral therapy, while successful at delaying the onset of AIDS or death, has yet to provide a functional cure. Alternative treatment strategies are clearly needed.

[0004] Intense interest in immunotherapy for HIV infection has been precipitated by emerging data indicating that the immune system has a major, albeit usually insufficient, role in limiting HIV replication. Virus-specific T-helper cells, which are critical to maintenance of cytolytic T cell (CTL) function, likely have a role. Viremia is also influenced by neutralizing antibodies, but they are generally low in magnitude in HIV infection and do not keep up with evolving viral variants in vivo.

[0005] Together these data indicate that increasing the strength and breadth of HIV-specific cellular immune responses might have a clinical benefit through so-called HIV immunotherapy. Some studies have tested vaccines against HIV, but success has been limited to date. Additionally, there has been interest in augmenting HIV immunotherapy by utilizing gene therapy techniques, but as with other immunotherapy approaches, success has been limited. One such method of implementing an HIV-specific immunotherapy or gene therapy could be via specially designed viral vectors.

[0006] Viral vectors can be used to transduce genes into target cells owing to specific virus envelope-host cell receptor interactions and viral mechanisms for gene expression. As a result, viral vectors have been used as vehicles for the transfer of genes into many different cell types including whole T cells or other immune cells as well as embryos, fertilized eggs, isolated tissue samples, tissue targets in situ and cultured cells. The ability to introduce and express foreign or altered genes in a cell is useful for therapeutic interventions such as gene therapy, somatic cell reprogramming of induced pluripotent stem cells, and various types of immunotherapy.

[0007] Gene therapy is one of the ripest areas of biomedical research with the potential to create new therapeutics that may involve the use of viral vectors. In view of the wide variety of potential genes available for therapy, an efficient means of delivering these genes is needed to fulfill the promise of gene therapy as a means of treating infectious and non-infectious diseases. Several viral systems including murine retrovirus, adenovirus, parvovirus (adeno-associated virus), vaccinia virus, and herpes virus have been developed as therapeutic gene transfer vectors.

[0008] There are many factors that must be considered when developing viral vectors, including tissue tropism, stability of virus preparations, stability and control of expression, genome packaging capacity, and construct-dependent vector stability. In addition, in vivo application of viral vectors is often limited by host immune responses against viral structural proteins and/or transduced gene products.

[0009] Thus, toxicity and safety are key hurdles that must be overcome for viral vectors to be used in vivo for the treatment of subjects. There are numerous historical examples of gene therapy applications in humans that have met with problems associated with the host immune responses against the gene delivery vehicles or the therapeutic gene products. Viral vectors (e.g., adenovirus) which co-transduce several viral genes together with one or more therapeutic gene(s) are particularly problematic.

[0010] Although lentiviral vectors do not generally induce cytotoxicity and do not elicit strong host immune responses, some lentiviral vectors such as HIV-1, which carry several immunostimulatory gene products, have the potential to cause cytotoxicity and induce strong immune responses in vivo. However, this may not be a concern for lentiviral derived transducing vectors that do not encode multiple viral genes after transduction. Of course, this may not always be the case, as sometimes the purpose of the vector is to encode a protein that will provoke a clinically useful immune response.

[0011] Another important issue related to the use of lentiviral vectors is that of possible cytopathogenicity upon exposure to some cytotoxic viral proteins. Exposure to certain HIV-1 proteins may induce cell death or functional unresponsiveness in T cells. Likewise, the possibility of generating replication-competent, virulent virus by recombination is often a concern.

[0012] Clearly, there is a need for improved treatment of HIV, and the present invention satisfies this need.

SUMMARY OF THE INVENTION

[0013] Disclosed herein are therapeutic immunization strategies and methods as well as highly effective therapeutic lentiviruses and other vectors capable of inhibiting HIV and reducing or altering expression of specific targets. The methods and compositions of the disclosed invention are useful for achieving a functional cure of HIV. More specifically, the invention includes methods for a functional cure of HIV that optimally combine therapeutic immunization of a patient, ex vivo re-stimulation of the patient's CD4 T cells, ex vivo lentivirus transduction of the enriched T cells, ex vivo culture of the cells, and reinfusion of the enriched, genetically modified cells. Additionally, the invention includes bioassays to measure treatment efficacy, sequential changes in therapeutic drug administration, monitoring intervals following withdrawal of HAART, and methods of diagnosis of a functional HIV cure.

[0014] In one aspect, the disclosed invention relates to a method of treating an HIV infection, comprising: (a) iden-

tifying a subject in need of treatment of HIV infection; (b) immunizing the subject with a therapeutically effective amount of an HIV vaccine; (c) removing lymphocytes from the subject and purifying peripheral blood mononuclear cells (PBMC); (d) contacting the PBMC ex vivo with a therapeutically effective amount of an HIV vaccine (which can be the same as or different from the HIV vaccine used in step (b)); (e) transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; (f) culturing the transduced PBMC for about 1 to about 21 or up to about 35 days (or any time frame in between these parameters, such as about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, or about 35 days); and (g) infusing the transduced PBMC into the subject.

[0015] In some embodiments, step (b) and step (d) utilize the same HIV vaccine, while in other embodiments, step (b) and step (d) utilize different HIV vaccines. In some embodiments, a patient may not require step (b) and/or step (d). Accordingly, in some embodiments, the disclosed methods may only comprise steps (a), (c), (d), (e), (f), and (g), or some combination thereof.

[0016] In some embodiments, the subject received cART or HAART prior to infusing the transduced PBMC into the subject. In some embodiments, the subject receives a cyclophosphamide pre-treatment prior to infusing the transduced PBMC into the subject.

[0017] In some embodiments, at least one genetic element is selected from the group consisting of a small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences. In some embodiments, the small RNA molecules targeting HIV RNA sequences are directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

[0018] In some embodiments, the transduced PBMC are cultured for about 1 to about 7 or up to about 10 days (or for any time frame in between these parameters, such as about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, or about 10 days) prior to infusing the transduced PBMC into the subject.

[0019] In another aspect, the disclosed invention relates to a viral vector for transducing HIV-specific CD4 T cells, wherein the viral vector encodes at least one genetic element selected from the group consisting of small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences.

[0020] In some embodiments, the vector is a lentivirus, but in other embodiments the vector is a DNA plasmid, adenoassociated virus, or other integrating or non-integrating vector systems for gene delivery.

[0021] In some embodiments, the small RNA molecules targeting HIV RNA sequences are directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

[0022] In another aspect, the disclosed invention relates to a bioassay for determining whether a HIV+ subject is functionally cured. Such a bioassay comprises determining the number of HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus, wherein the subject is functionally cured if the number of HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus is above a threshold value after a specified time following treatment according to disclosed methods.

[0023] In some embodiments, the threshold value is about 1×10^8 HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus, but the threshold value may be determined to be higher or lower than this value.

[0024] In some embodiments, the specified time following treatment is about 30 to about 60 days (or any time frame in between these two values), while in other embodiments the specified time following treatment is about 12 to about 26 weeks (or any time frame in between these two values).

[0025] In yet another aspect, the disclosed invention relates to a method of achieving a functional cure for HIV in a HIV+ subject. The method comprises: (a) identifying a subject that is HIV+; (b) immunizing the subject with a therapeutically effective amount of an HIV vaccine; (c) removing lymphocytes from the subject and purifying peripheral blood mononuclear cells (PBMC); (d) contacting the PBMC ex vivo with a therapeutically effective amount of an HIV vaccine; (f) culturing the transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; (f) culturing the transduced PBMC for about 1 to about 21 or up to 35 days (or any timeframe in between these values); and (g) infusing the transduced PBMC into the subject, wherein the HIV+ subject achieves a functional cure.

[0026] In some embodiments step (b) and step (d) comprise the same HIV vaccine, while in other embodiments, step (b) and step (d) comprise different HIV vaccines.

[0027] In some embodiments, the subject received cART or HAART prior to infusing the transduced PBMC into the subject. In some embodiments, the subject receives a cyclophosphamide pre-treatment or alternative conditioning therapies to improve T cell engraftment prior to infusing the transduced PBMC into the subject.

[0028] In some embodiments, at least one genetic element is selected from the group consisting of a small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences. In some embodiments, the small RNA molecules targeting HIV RNA sequences are directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

[0029] In some embodiments, the transduced PBMC are cultured for about 1 to about 7 or up to about 12 days (or any timeframe in between these two values, or other time periods described herein) prior to infusing the transduced PBMC into the subject.

[0030] The foregoing general description and following brief description of the drawings and detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed. Other objects, advantages, and novel features will be readily apparent to those skilled in the art from the following brief description of the drawings and detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0031] FIG. **1** shows a flow chart diagram of possible clinical therapy incorporating vaccination, cell collection to

obtain PBMC and CD4 T cells, ex vivo T cell culture after stimulating with vaccine immunogens and/or CD3/CD28 stimulation and/or mitogen stimulation in the presence of supporting growth cytokines (including, but not limited to, interleukin-2, interleukin-12, interleukin-15, interleukin-6, interleukin-7, and interleukin-23), lentivirus transduction to deliver anti-HIV genetic constructs, brief culture of transduced cells, and infusion back into the original subject.

[0032] FIG. **2** shows how CD4 T cells may be altered using gene therapy to prevent other cells from becoming infected or to prevent viral replication.

[0033] FIG. **3** shows a schematic of an exemplary therapeutic lentiviral construct. Therapeutic lentivirus constructs may substitute alternate sequences for the promoter region, targeting of regulatory RNA, and types of regulatory RNA. Further, the plasmids used for packaging the lentivirus particles can be altered according to production needs.

[0034] FIG. **4** shows exemplary vector sequences. Positive (genomic) strand sequence of the promoter and miR cluster were developed for inhibiting the spread of CCR5-tropic HIV strains. Sequences that are not underlined comprise the EF-1alpha promoter of transcription that was selected as best for this miR cluster. Sequences that are underlined show the miR cluster consisting of miR30 CCR5 (a modification of the natural human miR30 that redirects to CCR5 mRNA), miR21 Vif (redirects to Vif RNA sequence) and miR185 Tat (redirects to Tat RNA sequence). The sequences that are not underlined and in smaller font are restriction endonuclease cleavage sites that were incorporated into the oligonucle-otide primers for each of the miRNA constructs.

[0035] FIG. **5** shows that knockdown of CCR5 by an experimental vector prevents R5-tropic HIV infection in AGTc120 cells. Panel (A) shows CCR5 expression in AGTc120 cells with or without AGT103 lentivirus vector. Panel (B) shows the sensitivity of transduced AGTc120 cells to infection with a HIV BaL virus stock that was expressing green fluorescent protein (GFP) fused to the Nef gene of HIV.

[0036] FIG. **6** shows AGT103 reduces expression of Tat protein expression in cells transfected with HIV expression plasmid.

[0037] FIG. **7** shows AGT103 reduces levels of Vif protein expression in cells transfected with a full-length HIV expression plasmid. Cells were untreated (left lane and center lane) or transduced with AGT103 (left lane).

[0038] FIG. 8 shows generating a CD4+ T cell population that is highly enriched for HIV-specific, AGT103-transduced CD4 T cells. Panel (A) shows that therapeutic vaccination against HIV has minimal effect on the distribution of CD4+, CD8+ and CD4+/CD8+ T cells. An important CD4 T cell population is shown in the upper left quadrant of these analytical flow cytometry dot plots, and changes from 52% to 57% of total T cells after the vaccination series. These are representative data. Panel (B) shows the expression of CD4 and CD8 in a CD3+ population of peripheral blood mononuclear cells from a participant in an HIV therapeutic vaccine trial that were cultured for 12 days in medium+/interleukin-2/interleukin-12 or +/-interleukin-7/interleukin-15. Some cultures were stimulated with overlapping peptides representing the entire p55 Gag protein of HIV-1 (JPT PepMix) as a source of epitope peptides for T cell stimulation. Panel (C) shows that a combination of PepMix and interleukin-2/interleukin-12 provides for optimal expansion of antigen-specific CD4 T cells. The upper panels show the increase in cytokine (interferon-gamma) secreting cells in post-vaccination specimens exposed to PepMix. Panel (D) shows AGT103 transduction of antigen-expanded CD4 T cells can produce HIV-specific and HIV-resistant helper CD4 T cells for infusion into patients as part of a functional cure for HIV. The upper panels contain the results of analyzing the CD4+ T cell population in culture. The x axis is Green Fluorescent Protein (GFP) emission indicating that individual cells were transduced with the AGT103.

[0039] FIG. **9** shows the sequences of various exemplary cellular elements known to restrict HIV replication that may be incorporated into the disclosed vectors.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0040] Disclosed herein are methods and compositions for treating and/or preventing human immunodeficiency virus (HIV) disease to achieve a functional cure. A functional cure is defined as a condition resulting from the disclosed treatments and methods that reduces or eliminates the need for cART and may or may not require supporting adjuvant therapy. The methods of the invention include gene delivery by integrating lentivirus, non-integrating lentivirus, and related viral vector technology as described below.

[0041] Disclosed herein are therapeutic viral vectors (e.g., lentivirus vectors), immunotherapies, and methods for their use in a strategy to achieve a functional cure for HIV infection. The general strategy can include first therapeutic immunization with vaccines intended to produce strong immune responses against HIV in HIV-infected patients with stable suppression of viremia due to daily administration of HAART, for the purpose of enriching the fraction of HIV-specific CD4 T cells. This is followed by (1) isolating peripheral leukocytes by leukapheresis or purifying PBMC from venous blood, (2) re-stimulating CD4 T cells ex vivo with HIV vaccine proteins, (3) performing therapeutic lentivirus transduction, ex vivo T cell culture, and (4) reinfusion back into the original donor.

[0042] Previous efforts to achieve a cure for HIV have fallen short principally due to the failure to obtain sufficient numbers of HIV-specific CD4 T cells with protective genetic modifications. When this value is below a critical threshold, removing antiretroviral therapy allows HIV re-emergence, followed by rapid destruction of HIV-specific CD4 T cells, and also followed by return to progression of disease despite prior genetic therapy. By employing therapeutic immunization in the strategy described herein and providing highly effective therapeutic lentiviruses capable of inhibiting HIV, a new strategy for achieving a functional cure of HIV has been developed.

[0043] Also disclosed herein are novel viral vectors for enhancing HIV-specific CD4 T cells including lentiviral vectors and non-integrating, episomally replicating viral vectors and methods of using the same. Episomally replicating vectors like the present invention can comprise viral components from viruses like Papovaviridae (e.g. bovine papillomavirus or BPV) or Herpesviridae (e.g. Epstein Barr Virus or EBV) or Hepadnaviridae (e.g. Hepatitis B Virus or HBV). Episomal replicating vectors derived from these viruses may comprise a replication origin and at least one viral trans-acting factor, e.g., an initiator protein, such as E1 for BPV and EBNA-1 for EBV or HBV polymerase or terminus binding protein of Adenovirus. The process of 4

episomal replication typically incorporates both host cell replication machinery and viral trans-acting factors.

I. Human Immunodeficiency Virus (HIV)

[0044] HIV is a retrovirus that causes acquired immunodeficiency syndrome (AIDS) in humans. AIDS is a condition in which progressive failure of the immune system allows life-threatening opportunistic infections and cancers to thrive. Without treatment, average survival time after infection with HIV is estimated to be 9 to 11 years, depending upon the HIV subtype. Infection with HIV occurs by the transfer of bodily fluids, including but not limited to blood, semen, vaginal fluid, pre-ejaculate, saliva, tears, lymph or cerebro-spinal fluid, or breast milk. HIV may be present in an infected individual as both free virus particles and within infected immune cells.

[0045] HIV infects vital cells in the human immune system such as helper T cells, although tropism can vary among HIV subtypes. Immune cells that may be specifically susceptible to HIV infection include but are not limited to CD4+ T cells, macrophages, and dendritic cells. HIV infection leads to low levels of CD4+ T cells through a number of mechanisms, including but not limited to apoptosis of uninfected bystander cells, direct viral killing of infected cells, and killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections and cancer.

[0046] Structurally, HIV is distinct from many other retroviruses. The RNA genome consists of at least seven structural landmarks (LTR, TAR, RRE, PE, SLIP, CRS, and INS), and at least nine genes (gag, pol, env, tat, rev, nef, vif, vpr, vpu, and sometimes a tenth tev, which is a fusion of tat, env and rev), encoding 19 proteins. Three of these genes, gag, pol, and env, contain information needed to make the structural proteins for new virus particles.

[0047] HIV replicates primarily in CD4 T cells, and causes cellular destruction or dysregulation to reduce host immunity. Because HIV establishes infection as an integrated provirus and may enter a state of latency wherein virus expression in a particular cell decreases below the level for cytopathology affecting that cell or detection by the host immune system, HIV is difficult to treat and has not been eradicated even after prolonged intervals of highly active antiretroviral therapy (HAART). In the vast majority of cases, HIV infection causes fatal disease although survival may be prolonged by HAART.

[0048] A major goal in the fight against HIV is to develop strategies for curing disease. Prolonged HAART has not accomplished this goal, so investigators have turned to alternative procedures. Early efforts to improve host immunity by therapeutic immunization (using a vaccine after infection has occurred) had marginal or no impact. Likewise, treatment intensification had moderate or no impact.

[0049] Some progress has been made using genetic therapy, but positive results are sporadic and found only among rare human beings carrying defects in one or both alleles of the gene encoding CCR5 (chemokine receptor), which plays a critical role in viral penetration of host cells. However, many investigators are optimistic that genetic therapy holds the best promise for eventually achieving an HIV cure.

[0050] As disclosed herein, the methods and compositions of the invention are able to achieve a functional cure that may or may not include complete eradication of all HIV from the body. A functional cure is defined as a state or condition wherein HIV+ individuals who previously required HAART, may survive with low or undetectable virus replication and using lower or intermittent doses of HAART, or are potentially able to discontinue HAART altogether. As used herein, a functional cure may still possibly require adjunct therapy to maintain low level virus replication and slow or eliminate disease progression. A possible outcome of a functional cure is the eventual eradication of HIV to prevent all possibility of recurrence.

[0051] The primary obstacles to achieving a functional cure lie in the basic biology of HIV itself. Virus infection deletes CD4 T cells that are critical for all immune functions. Most importantly, HIV infection and depletion of CD4 T cells requires activation of individual cells. Activation is a specific mechanism for individual CD4 T cell clones that recognize pathogens or other molecules, using a rearranged T cell receptor.

[0052] In the case of HIV, infection activates a population of HIV-specific T cells that become infected and are consequently depleted before other T cells that are less specific for the virus, effectively crippling the immune system's defense against the virus. The capacity for HIV-specific T cell responses is rebuilt during prolonged HAART; however, when HAART is interrupted the rebounding virus infection repeats the process and again deletes the virus-specific cells, resetting the clock on disease progression.

[0053] Clearly, a functional cure is only possible if enough HIV-specific CD4 T cells are protected to allow for a host's native immunity to confront and control HIV once HAART is interrupted. In one embodiment, the present invention provides methods and compositions for improving the effectiveness of genetic therapy to provide a functional cure of HIV disease. In another embodiment, the present invention provides methods and compositions for enhancing host immunity against HIV to provide a functional cure. In yet another embodiment, the present invention provides methods and compositions for enhancing host immunity against HIV to provide a functional cure. In yet another embodiment, the present invention provides methods and compositions for enriching HIV-specific CD4 T cells in a patient to achieve a functional cure.

[0054] In one embodiment of the invention, treatment results in enriching a subject's HIV-specific CD4 T cells by about 100%, about 200%, about 300%, about 400%, about 500%, about 600%, about 700%, about 800%, about 900%, about 1000%, about 1500%, about 2000%, about 2500%, about 3000%, about 3500%, about 4000%, about 4500%, about 5000%, about 5500%, about 6000%, about 6500%, about 7000%, about 7500%, about 8000%, about 8500%, about 9000%, about 9500%, about 10000%, about 11000%, about 12000%, about 13000%, about 14000%, about 15000%, about 16000%, about 17000%, about 18000%, about 19000%, about 20000%, about 25000%, about 30000%, about 35000%, about 40000%, about 45000%, about 50000%, about 55000%, about 60000%, about 65000%, about 70000%, about 75000%, about 80000%, about 85000%, about 90000%, about 95000%, about 100000%, or any value in between.

II. Gene Therapy

[0055] Viral vectors are used to deliver genetic constructs to host cells for the purposes of disease therapy or prevention.

[0056] Genetic constructs can include, but are not limited to, functional genes or portions of genes to correct or complement existing defects, DNA sequences encoding regulatory proteins, DNA sequences encoding regulatory RNA molecules including antisense, short homology RNA, long non-coding RNA, small interfering RNA or others, and decoy sequences encoding either RNA or proteins designed to compete for critical cellular factors to alter a disease state. Gene therapy involves delivering these therapeutic genetic constructs to target cells to provide treatment or alleviation of a particular disease.

[0057] There are multiple ongoing efforts to utilize genetic therapy in the treatment of HIV disease, but thus far, the results have been poor. A small number of treatment successes were obtained in rare HIV patients carrying a spontaneous deletion of the CCR5 gene (an allele known as CCR5delta32).

[0058] Lentivirus-delivered nucleases or other mechanisms for gene deletion/modification may be used to lower the overall expression of CCR5 and/or help to lower HIV replication. At least one study has reported having success in treating the disease when lentivirus was administered in patients with a genetic background of CCR5delta32. However, this was only one example of success, and many other patients without the CCR5delta32 genotype have not been treated as successfully. Consequently, there is a substantial need to improve the performance of viral genetic therapy against HIV, both in terms of performance for the individual viral vector construct and for improved use of the vector through a strategy for achieving functional HIV cure.

[0059] For example, some existing therapies rely on zinc finger nucleases to delete a portion of CCR5 in an attempt to render cells resistant to HIV infection. However, even after optimal treatment, only 30% of T cells had been modified by the nuclease at all, and of those that were modified, only 10% of the total CD4 T cell population had been modified in a way that would prevent HIV infection. In contrast, the disclosed methods result in virtually every cell carrying a lentivirus transgene having a reduction in CCR5 expression below the level needed to allow HIV infection.

[0060] For the purposes of the disclosed methods, gene therapy can include, but is not limited to, affinity-enhanced T cell receptors, chimeric antigen receptors on CD4 T cells (or alternatively on CD8 T cells), modification of signal transduction pathways to avoid cell death cause by viral proteins, increased expression of HIV restriction elements including TREX, SAMHD1, MxA or MxB proteins, APOBEC complexes, TRIM5-alpha complexes, tetherin (BST2), and similar proteins identified as being capable of reducing HIV replication in mammalian cells.

[0061] For example, in some embodiments the disclosed vectors may include, but are not limited to, the restriction elements found in Table 1 below. The sequences of these exemplary restriction elements are further disclosed in FIG. 9.

TABLE 1

Gene	Accession Number
TREX1	NM_016381 (human)
TREX2	XM_015128506.1 (<i>Macaca mulatta</i>) NM_080701/NM_007205 (human
1161/12	XM 015128506.1 (Macaca mulatta)

TABLE 1-continued

Gene	Accession Number
SAMHD1	NM_015474 (human) IN036895 1 (Magaga mulatta)
MxA	NM_001144925 (human)
	JX297237.1 (Macaca mulatta)
MxB	NM_002463 (human)
APOBEC3G	NM_021822 (human)
	XM_015150306 (Macaca mulatta)
TRIM5-alpha	NM_033034 (human)
-	NM_001032910.1 (Macaca mulatta)
Tetherin	NM_004335 (human)
	FJ943432.1 (Macaca mulatta)

III. Immunotherapy

[0062] Historically, vaccines have been a go-to weapon against deadly infectious diseases, including smallpox, polio, measles, and yellow fever. Unfortunately, there is no currently approved vaccine for HIV. The HIV virus has unique ways of evading the immune system, and the human body seems incapable of mounting an effective immune response against it. As a result, scientists do not have a clear picture of what is needed to provide protection against HIV. [0063] However, immunotherapy may provide a solution that was previously unaddressed by conventional vaccine approaches. Immunotherapy, also called biologic therapy, is a type of treatment designed to boost the body's natural defenses to fight infections or cancer. It uses materials either made by the body or in a laboratory to improve, target, or restore immune system function.

[0064] In some embodiments of the disclosed invention, immunotherapeutic approaches may be used to enrich a population of HIV-specific CD4 T cells for the purpose of increasing the host's anti-HIV immunity. In some embodiments of the disclosed invention, integrating or non-integrating lentivirus vectors may be used to transduce a host's immune cells for the purposes of increasing the host's anti-HIV immunity. In yet another embodiment of the invention, a vaccine comprising HIV proteins including but not limited to a killed particle, a virus-like particle, HIV peptides or peptide fragments, a recombinant viral vector, a recombinant bacterial vector, a purified subunit or plasmid DNA combined with a suitable vehicle and/or biological or chemical adjuvants to increase a host's immune responses may be used to enrich the population of virus-specific T cells or antibodies, and these methods may be further enhanced through the use of HIV-targeted genetic therapy using lentivirus or other viral vector.

IV. Methods According to the Invention

[0065] In one aspect, the disclosed invention provides methods for using viral vectors to achieve a functional cure for HIV disease. The methods further include immuno-therapy to enrich the proportion of HIV-specific CD4 T cells, followed by lentivirus transduction to deliver inhibitors of HIV and CCR5 and CXCR4 as required.

[0066] In one embodiment, the methods include therapeutic immunization as a method for enriching the proportion of HIV-specific CD4 T cells. Therapeutic immunization can include purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, virus-like particles (VLPs), biological or chemical adjuvants including cytokines and/or chemokines, vehicles, and methods for immunization.

[0067] Therapeutic vaccines can include one or more HIV protein with protein sequences representing the predominant viral types of the geographic region where treatment is occurring. Therapeutic vaccines will include purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, virus-like particles (VLPs), biological or chemical adjuvants including cytokines and/or chemokines, vehicles, and methods for immunization. Vaccinations may be administered according to standard methods known in the art and HIV patients may continue antiretroviral therapy during the interval of immunization and subsequent ex vivo lymphocyte culture including lentivirus transduction.

[0068] In some embodiments, HIV+ patients can be immunized with an HIV vaccine, increasing the frequency of HIV-specific CD4 T cells by about 2, about 25, about 250, about 500, about 750, about 1000, about 1250, or about 1500-fold (or any amount in between these values). The vaccine may be any clinically utilized or experimental HIV vaccine, including the disclosed lentiviral, other viral vectors or other bacterial vectors used as vaccine delivery systems. For instance, the disclosed vector may comprise a recombinant Bacille Calmette Guerin (BCG) strain expressing HIV VLP. BCG is mycobacterium bovis attenuated for use as a human vaccine against tuberculosis. In another embodiment, the vectors can encode virus-like particles (VLPs) to induce higher titers of neutralizing antibodies. In another embodiment, the vectors can encode peptides or peptide fragments associated with HIV including but not limited to gag, pol, and env, tat, rev, nef, vif, vpr, vpu, and tev, as well as LTR, TAR, RRE, PE, SLIP, CRS, and INS. Alternatively, the HIV vaccine used in the disclosed methods may comprise purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, virus-like particles (VLPs), or biological or chemical adjuvants including cytokines and/or chemokines.

[0069] In one embodiment, the methods include ex vivo re-stimulation of CD4 T cells from persons or patients previously immunized by therapeutic vaccination, using purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, biological or chemical adjuvants including cytokines and/or chemokines, vehicles, and methods for re-stimulation. Ex vivo re-stimulation may be performed using the same vaccine or immune stimulating compound used for in vivo immunization, or it may be performed using a different vaccine or immune stimulating compound than those used for in vivo immunization. Moreover, in some embodiments, the patient may not require prior therapeutic vaccination or re-stimulation of CD4 T cells if the individual has sufficiently high antigen-specific CD4 T cell responses to HIV proteins. In these embodiments, such a patient may only require administration of the disclosed viral vectors to achieve a functional cure.

[0070] For example, peripheral blood mononuclear cells (PBMCs) can be obtained by leukapheresis and treated ex vivo to obtain about 1×10^{10} CD4 T cells of which about 0.1%, about 1%, about 5% or about 10% or about 30% are both HIV-specific in terms of antigen responses, and HIV-resistant by virtue of carrying the therapeutic transgene delivered by the disclosed lentivirus vector. Alternatively, about 1×10^7 , about 1×10^8 , about 1×10^9 , about 1×10^{10} , about

 1×10^{11} , or about 1×10^{12} CD4 T cells may be isolated for re-stimulation. Any suitable amount of CD4 T cells can be isolated for ex vivo re-stimulation.

[0071] The isolated CD4 T cells can be cultured in appropriate medium throughout re-stimulation with HIV vaccine antigens, which may include antigens present in the prior therapeutic vaccination. Antiretroviral therapeutic drugs including inhibitors of reverse transcriptase, protease or integrase may be added to prevent virus re-emergence during prolonged ex vivo culture. CD4 T cell re-stimulation is used to enrich the proportion of HIV-specific CD4 T cells in culture. The same procedure may also be used for analytical objectives wherein smaller blood volumes with peripheral blood mononuclear cells obtained by purification, are used to identify HIV-specific T cells and measure the frequency of this sub-population.

[0072] The PBMC fraction may be enriched for HIV-specific CD4 T cells by contacting the cells with HIV proteins matching or complementary to the components of the vaccine previously used for in vivo immunization. Ex vivo re-stimulation can increase the relative frequency of HIV-specific CD4 T cells by about 25, about 50, about 75, about 100, about 125, about 150, about 175, or about 200-fold.

[0073] The methods additionally include combining in vivo therapeutic immunization and ex vivo re-stimulation of CD4 T cells with ex vivo lentiviral transduction and culturing.

[0074] Thus, in one embodiment, the re-stimulated PBMC fraction that has been enriched for HIV-specific CD4 T cells can be transduced with therapeutic anti-HIV lentivirus or other vectors and maintained in culture about 1 to about 21 days or up to about 35 days. Alternatively, the cells may be cultured for about 1-about 18 days, about 1-about 15 days, about 1-about 7 days. Thus, the transduced cells may be cultured for about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, or about 35 days.

[0075] Once the transduced cells have been sufficiently cultured, transduced CD4 T cells are infused back into the original patient. Infusion can be performed using various machines and methods known in the art. In some embodiments, infusion may be accompanied by pre-treatment with cyclophosphamide or similar compounds to increase the efficiency of re-engraftment.

[0076] In some embodiments, a CCR5-targeted therapy may be added to a subject's antiretroviral therapy regimen, which was continued throughout the treatment process. Examples of CCR5-targeted therapies include but are not limited to Maraviroc (a CCR5 antagonist) or Rapamycin (immunosuppressive agent that lowers CCR5). In some embodiments, the antiretroviral therapy may be ceased and the subject can be tested for virus rebound. If no rebound occurs, adjuvant therapy can also be removed and the subject can be tested again for virus rebound.

[0077] Continued virus suppression with reduced or no antiretroviral therapy include cART or HAART, and reduced or no adjuvant therapy for about 26 weeks can be considered a functional cure for HIV. Other definitions of a functional cure are described herein.

[0078] The lentiviral and other vectors used in the disclosed methods may encode at least one, at least two, at least three, at least four, or at least five genes of interest. Given the versatility and therapeutic potential of HIV-targeted gene therapy, a viral vector of the invention may encode genes or nucleic acid sequences that include but are not limited to (i) an antibody directed to an antigen associated with an infectious disease or a toxin produced by the infectious pathogen, (ii) cytokines including interleukins that are required for immune cell growth or function and may be therapeutic for immune dysregulation encountered in HIV and other chronic or acute human viral or bacterial pathogens, (iii) factors that suppress the growth of HIV in vivo including CD8 suppressor factors, (iv) mutations or deletions of chemokine receptor CCR5, mutations or deletions of chemokine receptor CXCR4, or mutations or deletions of chemokine receptor CXCR5, (v) antisense DNA or RNA against specific receptors or peptides associated with HIV or host protein associated with HIV, (vi) small interfering RNA against specific receptors or peptides associated with HIV or host protein associated with HIV, or (vii) a variety of other therapeutically useful sequences that may be used to treat HIV or AIDS.

[0079] Additional examples of HIV-targeted gene therapy that can be used in the disclosed methods include, but are not limited to, affinity-enhanced T cell receptors, chimeric antigen receptors on CD4 T cells (or alternatively on CD8 T cells), modification of signal transduction pathways to avoid cell death cause by viral proteins, increased expression of HIV restriction elements including TREX, SAMHD1, MxA or MxB proteins, APOBEC complexes, TRIM5-alpha complexes, tetherin (BST2), and similar proteins identified as being capable of reducing HIV replication in mammalian cells.

[0080] In some embodiment, a patient may be undergoing cART or HAART concurrently while being treated according to the methods of the invention. In other embodiments, a patient may undergo cART or HAART before or after being treated according to the methods of the invention. In some embodiments, cART or HAART is maintained throughout treatment according to the methods of the invention and the patient may be monitored for HIV viral burden in blood and frequency of lentivirus-transduced CD4 T cells in blood. Preferably, a patient receiving cART or HAART prior to being treated according to the methods of the invention is able to discontinue or reduce cART or HAART following treatment according to the methods of the invention.

[0081] For efficacy purposes, the frequency of transduced, HIV-specific CD4 T cells, which is a novel surrogate marker for gene therapy effects, may be determined, as discussed in more detail in Section VI.

V. Compositions According to the Invention

[0082] In one aspect, the disclosed invention provides lentiviral vectors capable of delivering genetic constructs to inhibit HIV penetration of susceptible cells. For instance, one mechanism of action is to reduce mRNA levels for CCR5 and/or CXCR4 chemokine receptors and thus reduce the rates for viral entry into susceptible cells.

[0083] Alternatively, the disclosed lentiviral vectors may be capable of inhibiting the formation of DNA and HIVinfected cells by reducing the stability of incoming HIV genomic RNA. And in yet another embodiment, the disclosed lentivirus vectors are capable of preventing HIV production from a latently infected cell, wherein the mechanism of action is to cause instability of viral RNA sequences through the action of inhibitory RNA including short-homology, small-interfering or other regulatory RNA species. [0084] The therapeutic lentiviruses disclosed in this application generally comprise at least one of two types of genetic cargo. First, the lentiviruses may encode genetic elements that direct expression of small RNA capable of inhibiting the production of chemokine receptors CCR5 and/or CXCR4 that are important for HIV penetration of susceptible cells. The second type of genetic cargo includes constructs capable of expressing small RNA molecules targeting HIV RNA sequences for the purpose of preventing reverse transcription, RNA splicing, RNA translation to produce proteins, or packaging of viral genomic RNA for particle production and spreading infection. An exemplary structure is diagrammed in FIG. 3.

[0085] As shown in FIG. **3**, an exemplary construct may comprise numerous sections or components. For example, in one embodiment, an exemplary LV construct may comprise the following sections or components:

- [0086] RSV—a Rous Sarcoma virus long terminal repeat;
- [0087] 5'LTR—a portion of an HIV long terminal repeat that can be truncated to prevent replication of the vector after chromosomal integration;
- [0088] Psi—a packaging signal that allows for incorporation of the vector RNA genome into viral particles during packaging;
- [0089] RRE—a Rev Responsive element can be added to improve expression from the transgene by mobilizing RNA out of the nucleus and into the cytoplasm of cells;
- [0090] c PPT—a Poly purine tract that facilitates second strand DNA synthesis prior to integration of the transgene into the host cell chromosome;
- [0091] Promoter—a promoter initiates RNA transcription from the integrated transgene to express micro-RNA clusters (or other genetic elements of the construct), and in some embodiments, the vectors may use an EF-1 promoter;
- **[0092]** Anti-CCR5—a micro RNA targeting messenger RNA for the host cell factor CCR5 to reduce its expression on the cell surface;
- [0093] Anti-Rev/Tat—a micro RNA targeting HIV genomic or messenger RNA at the junction between HIV Rev and Tat coding regions, which is sometimes designated miRNA Tat or a\given a similar description in this application;
- [0094] Anti-Vif—a micro RNA targeting HIV genomic or messenger RNA within the Vif coding region;
- [0095] WPRE—a woodchuck hepatitis virus post-transcriptional regulatory element is an additional vector component that can be used to facilitate RNA transport of the nucleus; and
- [0096] deltaU3 3'LTR—a modified version of a HIV 3-prime long terminal repeat where a portion of the U3 region has been deleted to improve safety of the vector.

One of skill in the art will recognize that the above components are merely examples, and that such components may be reorganized, substituted with other elements, or 8

otherwise changed, so long as the construct is able to prevent expression of HIV genes and decrease the spread of infection.

[0097] Vectors of the invention may include either or both of the types of genetic cargo discussed above (i.e. genetic elements that direct expression of a gene or small RNAs, such as siRNA, shRNA, or miRNA that can prevent translation or transcription), and the vectors of the invention may also encode additionally useful products for the purpose of treatment or diagnosis of HIV. For instance, in some embodiments, these vectors may also encode green fluorescent protein (GFP) for the purpose of tracking the vectors or antibiotic resistance genes for the purposes of selectively maintaining genetically-modified cells in vivo.

[0098] The combination of genetic elements incorporated into the disclosed vectors is not particularly limited. For example, a vector may encode a single small RNA, two small RNAs, three small RNA, four small RNAs, five small RNAs, six small RNAs, seven small RNAs, eight small RNAs, nine small RNAs, or ten small RNAs. Such vectors may additionally encode other genetic elements to function in concert with the small RNAs to prevent expression and infection of HIV.

[0099] Those of skill in the art will understand that the therapeutic lentivirus may substitute alternate sequences for the promoter region, targeting of regulatory RNA, and types of regulatory RNA. Further, the therapeutic lentivirus of the invention may comprise changes in the plasmids used for packaging the lentivirus particles; these changes are required to increase levels of production in vitro.

[0100] In some embodiments, the vector used in the disclosed methods may be a DNA plasmid, adeno-associated virus, or other integrating or non-integrating vector systems for gene delivery.

VI. Bioassays

[0101] In one aspect, the present invention includes bioassays for determining the success of HIV treatment for achieving a functional cure. These assays will provide a method for measuring the efficacy of the disclosed methods of immunization and treatment by measuring the frequency of transduced, HIV specific CD4 T cells in a patient. HIV-specific CD4 T cells are recognizable because they proliferate, change the composition of cell surface markers, induce signaling pathways including phosphorylation, or express specific marker proteins that may be cytokines, chemokines, caspases, phosphorylated signaling molecules or other cytoplasmic and/or nuclear components. Specific responding CD4 T cells are recognized for example, using labeled monoclonal antibodies or specific in situ amplification of mRNA sequences, that allow sorting of HIV-specific cells using flow cytometry sorting, magnetic bead separation or other recognized methods for antigen-specific CD4 T cell isolation. The isolated CD4 T cells are tested to determine the frequency of cells carrying integrated therapeutic lentivirus. Single cell testing methods may also be used including microfluidic separation of individual cells, that are coupled with mass spectrometry, PCR, ELISA or antibody staining to confirm responsiveness to HIV and presence of integrated therapeutic lentivirus.

[0102] Thus, in one embodiment, following application of a treatment according to the invention (e.g., (a) immunization, (b) ex vivo lymphocyte culture; (c) re-stimulation with purified proteins, inactivated viruses, virally vectored pro-

teins, bacterially vectored proteins, biological or chemical adjuvants including cytokines and/or chemokines, vehicles; and (d) infusion of the enriched, transduced T cells), a patient may be subsequently assayed to determine the efficacy of the treatment. A threshold value of target T cells in the body may be established to measure a functional cure at, for instance, about 1×10^8 HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus. The value threshold value of cells refers to the total body content. It may not be measured directly, but instead may be extrapolated from blood CD4 T cell counts using a standard correction. For example, it is common in the art to assume that 90% of CD4 T cells are present in tissues and only 10% are found in blood.

[0103] Alternatively, the threshold value may be about 1×10^5 , about 1×10^6 , about 1×10^7 , about 1×10^9 , or about 1×10^{10} CD4 T cells in the body of the patient.

[0104] HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus can be determined using any suitable method, such as but not limited to flow cytometry, cell sorting, FACS analysis, DNA cloning, PCR, RT-PCR or Q-PCR, ELISA, FISH, western blotting, southern blotting, high throughput sequencing, RNA sequencing, oligonucleotide primer extension, or other methods known in the art.

[0105] Methods for defining antigen specific T cells with genetic modifications are known in the art. However, utilizing such methods to combine identifying HIV-specific T cells with integrated or non-integrated gene therapy constructs as a standard measure for efficacy is a new concept in the field of HIV treatment.

VII. Doses and Dosage Forms

[0106] The disclosed methods and compositions can be used for treating HIV+ patients during various stages of their disease. Accordingly, dosing regimens may vary based upon the condition of the patient and the method of administration.

[0107] In one embodiment, HIV-specific vaccines for the initial in vivo immunization may be administered to a subject in need in varying doses. In general, vaccines delivered by intramuscular injection include about 10 µg to about 300 µg, about 25 µg to about 275 µg, about 50 µg to about 250 µg, about 75 µg to about 225, or about 100 µg to about 200 µg of HIV protein, either total virus protein prepared from inactivated virus particles, virus-like particles or purified virus protein from recombinant systems or purified from virus preparations. Recombinant viral or bacterial vectors may be administered by any and all of the routes described. Intramuscular vaccines will include about 1 µg to about 100 µg, about 10 µg to about 90 µg, about 20 µg to about 80 µg, about 30 µg to about 70 µg, about 40 µg to about 60 µg, or about 50 µg of suitable adjuvant molecules and be suspended in oil, saline, buffer or water in volumes of 0.1 to 5 ml per injection dose, and may be soluble or emulsion preparations. Vaccines delivered orally, rectally, bucally, at genital mucosal or intranasally, including some virallyvectored or bacterially-vectored vaccines, fusion proteins, liposome formulations or similar preparations, may contain higher amounts of virus protein and adjuvant. Dermal, sub-dermal or subcutaneous vaccines utilize protein and adjuvant amounts more similar to oral, rectal or intranasaldelivered vaccines. Depending on responses to the initial immunization, vaccination may be repeated 1-5 times using

the same or alternate routes for delivery. Intervals may be of 2-24 weeks between immunizations. Immune responses to vaccination are measured by testing HIV-specific antibodies in serum, plasma, vaginal secretions, rectal secretions, saliva or bronchoalveolar lavage fluids, using ELISA or similar methodology. Cellular immune responses are tested by in vitro stimulation with vaccine antigens followed by staining for intracellular cytokine accumulation followed by flow cytometry or similar methods including lymphoproliferation, expression of phosphorylated signaling proteins or changes in cell surface activation markers. Upper limits of dosing may be determined based on the individual patient and will depend on toxicity/safety profiles for each individual product or product lot.

[0108] Immunization may occur once, twice, three times, or repeatedly. For instance, an agent for HIV immunization may be administered to a subject in need once a week, once every other week, once every three weeks, once a month, every other month, every three months, every six months, every nine months, once a year, every eighteen months, every two years, every 36 months, or every three years.

[0109] Immunization will occur at least once before ex vivo expansion and enrichment of CD4 T cells, and immunization may occur once, twice, three times, or more after ex vivo lymphocyte culture/re-stimulation and infusion.

[0110] In one embodiment, HIV-vaccines for immunization are administered as a pharmaceutical composition. In one embodiment, the pharmaceutical composition comprising an HIV vaccine can be formulated in a wide variety of nasal, pulmonary, oral, topical, or parenteral dosage forms for clinical application. Each of the dosage forms can comprise various disintegrating agents, surfactants, fillers, thickeners, binders, diluents such as wetting agents or other pharmaceutically acceptable excipients. The pharmaceutical composition comprising an HIV vaccine can also be formulated for injection.

[0111] HIV vaccine compositions for the purpose of immunization can be administered using any pharmaceutically acceptable method, such as intranasal, buccal, sublingual, oral, rectal, ocular, parenteral (intravenously, intradermally, intramuscularly, subcutaneously, intracisternally, intraperitoneally), pulmonary, intravaginal, locally administered, topically administered, topically administered after scarification, mucosally administered, via an aerosol, or via a buccal or nasal spray formulation.

[0112] Further, the HIV vaccine compositions can be formulated into any pharmaceutically acceptable dosage form, such as a solid dosage form, tablet, pill, lozenge, capsule, liquid dispersion, gel, aerosol, pulmonary aerosol, nasal aerosol, ointment, cream, semi-solid dosage form, and a suspension. Further, the composition may be a controlled release formulation, sustained release formulation, immediate release formulation, or any combination thereof. Further, the composition may be a transdermal delivery system.

[0113] In another embodiment, the pharmaceutical composition comprising an HIV vaccine can be formulated in a solid dosage form for oral administration, and the solid dosage form can be powders, granules, capsules, tablets or pills. In yet another embodiment, the solid dosage form can include one or more excipients such as calcium carbonate, starch, sucrose, lactose, microcrystalline cellulose or gelatin. In addition, the solid dosage form can include, in addition to the excipients, a lubricant such as talc or magnesium stearate. In some embodiments, the oral dosage form can be

immediate release or a modified release form. Modified release dosage forms include controlled or extended release, enteric release, and the like. The excipients used in the modified release dosage forms are commonly known to a person of ordinary skill in the art.

[0114] In a further embodiment, the pharmaceutical composition comprising a HIV vaccine can be formulated as a sublingual or buccal dosage form. Such dosage forms comprise sublingual tablets or solution compositions that are administered under the tongue and buccal tablets that are placed between the cheek and gum.

[0115] In yet a further embodiment, the pharmaceutical composition comprising an HIV vaccine can be formulated as a nasal dosage form. Such dosage forms of the present invention comprise solution, suspension, and gel compositions for nasal delivery.

[0116] In one embodiment, the pharmaceutical composition can be formulated in a liquid dosage form for oral administration, such as suspensions, emulsions or syrups. In other embodiments, the liquid dosage form can include, in addition to commonly used simple diluents such as water and liquid paraffin, various excipients such as humectants, sweeteners, aromatics or preservatives. In particular embodiments, the composition comprising HIV vaccine or a pharmaceutically acceptable salt thereof can be formulated to be suitable for administration to a pediatric patient.

[0117] In one embodiment, the pharmaceutical composition can be formulated in a dosage form for parenteral administration, such as sterile aqueous solutions, suspensions, emulsions, non-aqueous solutions or suppositories. In other embodiments, the non-aqueous solutions or suspensions can include propyleneglycol, polyethyleneglycol, vegetable oils such as olive oil or injectable esters such as ethyl oleate. As a base for suppositories, witepsol, macrogol, tween 61, cacao oil, laurin oil or glycerinated gelatin can be used.

[0118] The dosage of the pharmaceutical composition can vary depending on the patient's weight, age, gender, administration time and mode, excretion rate, and the severity of disease.

[0119] For the purposes of re-stimulation, lymphocytes, PBMCs, and/or CD4 T cells are removed from a patient and isolated for re-stimulation and culturing. The isolated cells may be contacted with the same HIV vaccine or activating agent used for immunization or a different HIV vaccine or activating agent. In one embodiment, the isolated cells are contacted with about 10 ng to 5 μ g of an HIV vaccine or activating agent per about 10⁶ cells in culture (or any other suitable amount). More specifically, the isolated cells may be contacted with about 50 ng, about 100 ng, about 200 ng, about 300 ng, about 400 ng, about 500 ng, about 200 ng, about 700 ng, about 400 ng, about 500 ng, about 3.5 μ g, about 4.5 μ g, or about 5 μ g of an HIV vaccine or activating agent per about 10⁶ cells in culture.

[0120] Activating agents or vaccines are generally used once for each in vitro cell culture but may be repeated after intervals of about 15 to about 35 days. For example, a repeat dosing could occur at about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, or about 35 days. **[0121]** For transduction of the enriched, re-stimulated cells, the cells may be transduced with lentiviral vectors or

with other known vector systems as disclosed in Section V and FIG. **3**. The cells being transduced may be contacted with about 1-1,000 viral genomes (measured by RT-PCR assay of culture fluids containing lentivirus vector) per target cell in culture (or any other suitable amount). Lentivirus transduction may be repeated 1-5 times using the same range of 1-1,000 viral genomes per target cell in culture.

VIII. Definitions

[0122] As used herein, the term "about" will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

[0123] "A treatment" is intended to target the disease state and combat it, i.e., ameliorate or prevent the disease state. The particular treatment thus will depend on the disease state to be targeted and the current or future state of medicinal therapies and therapeutic approaches. A treatment may have associated toxicities.

[0124] The terms "administration of" or "administering" an active agent should be understood to mean providing an active agent of the invention to the subject in need of treatment in a form that can be introduced into that individual's body in a therapeutically useful form and therapeutically effective amount.

[0125] The term "therapeutically effective amount" refers to a sufficient quantity of the active agents of the present invention, in a suitable composition, and in a suitable dosage form to treat or prevent the symptoms, progression, or onset of the complications seen in patients suffering from a given ailment, injury, disease, or condition. The therapeutically effective amount will vary depending on the state of the patient's condition or its severity, and the age, weight, etc., of the subject to be treated. A therapeutically effective amount can vary, depending on any of a number of factors, including, e.g., the route of administration, the condition of the subject, as well as other factors understood by those in the art.

[0126] The term "treatment" or "treating" generally refers to an intervention in an attempt to alter the natural course of the subject being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects include, but are not limited to, preventing occurrence or recurrence of disease, alleviating symptoms, suppressing, diminishing or inhibiting any direct or indirect pathological consequences of the disease, ameliorating or palliating the disease state, and causing remission or improved prognosis.

[0127] The term "functional cure" refers to a state or condition wherein HIV+ individuals who previously required cART or HAART, may survive with low or undetectable virus replication using lower doses, intermittent doses, or discontinued dosing of cART or HAART. An individual may be said to have been "functionally cured" while still requiring adjunct therapy to maintain low level virus replication and slow or eliminate disease progression. A possible outcome of a functional cure is the eventual eradication of HIV to prevent all possibility of recurrence. [0128] The term "HIV vaccine" encompasses immunogens plus vehicle plus adjuvant intended to elicit HIVspecific immune responses. Vaccine may include purified or whole inactivated virus particles that may be HIV or a recombinant virus vectors capable of expressing HIV proteins, protein fragments or peptides, glycoprotein fragments or glycopeptides, in addition to recombinant bacterial vectors, plasmid DNA or RNA capable of directing cells to producing HIV proteins, glycoproteins or protein fragments able to elicit specific immunity. Alternately, specific methods for immune stimulation including anti-CD3/CD28 beads, T cell receptor-specific antibodies, mitogens, superantigens and other chemical or biological stimuli may be used to activate dendritic, T or B cells for the purposes of enriching HIV-specific CD4 T cells prior to transduction or for in vitro assay of lentivirus-transduced CD4 T cells. Activating substances may be soluble, polymeric assemblies, liposome or endosome-based or linked to beads. Cytokines including interleukin-2, 6, 7, 12, 15, 23 or others may be added to improve cellular responses to stimuli and/or improve the survival of CD4 T cells throughout the culture and transduction intervals.

[0129] The terms "individual," "host," "subject," and "patient" are used interchangeably herein.

[0130] As used herein, "expression," "expressed," or "encodes" refers to the process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. Expression may include splicing of the mRNA in a eukaryotic cell or other forms of post-transcriptional modification or post-translational modification.

[0131] As used herein, "small RNA" refers to non-coding RNA that are generally less than about 200 nucleotides or less in length and possess a silencing or interference function. In other embodiments, the small RNA is about 175 nucleotides or less, about 150 nucleotides or less, about 125 nucleotides or less, about 100 nucleotides or less, or about 75 nucleotides or less in length. Such RNAs include micro-RNA (miRNA), small interfering RNA (siRNA), double stranded RNA (dsRNA), and short hairpin RNA (shRNA). "Small RNA" of the disclosure should be capable of inhibiting or knocking-down gene expression of a target gene, generally through pathways that result in the destruction of the target gene mRNA.

[0132] The following examples are given to illustrate the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples. All printed publications referenced herein are specifically incorporated by reference.

EXAMPLES

Example 1-Clinical Study for Treatment of HIV

[0133] Screening and Informed Consent.

[0134] Select HIV+ participants receiving combination antiretroviral therapy (cART) with stable suppression of virus burden are chosen to participate.

[0135] Immunization with a Therapeutic HIV Vaccine.

[0136] A vaccine that already has IND status and has been used in a clinical trial involving HIV+ participants is administered to participants. This step will increase the relative frequency of HIV-specific CD4 T cells by approximately 1,000-fold.

[0137] Next, blood lymphocytes are removed by leukapheresis and further purified to the peripheral blood mononuclear cell (PBMC) fraction. Alternately, cells may be purified from venous blood by column or density gradient methods.

[0138] Cultured PBMC are stimulated with HIV proteins or peptides matching or complementing the components in the therapeutic vaccine (perhaps using the same vaccine depending on its composition). This step will increase the relative frequency of HIV-specific CD4 T cells by approximately 100-fold.

[0139] Cultured PBMC cells are infected with therapeutic lentivirus or other disclosed vectors, for instance, one that encodes small RNA to interfere with the translation of CCR5 and viral replication proteins. After transduction, the cells are maintained in culture 3-7 days.

[0140] Transduced CD4 T cells are infused back into the original participant. Infusion can be performed according to methods known in the art. This may require pre-treatment with cyclophosphamide to increase the efficiency of reengraftment.

[0141] For participants receiving cART, cART will be maintained throughout. Following infusion, monitor HIV viral burden in blood and frequency of lentivirus-transduced CD4 T cells in blood.

[0142] If participants meet qualifying criteria (including having $>10^{\circ}$, $>10^{7}$, $>10^{8}$ Lentivirus-transduced, HIV-specific CD4 T cells in the total of blood and tissue compartments), they may move to an efficacy study. If participants miss the qualifying criterion they may be eligible to receive a second dose of therapeutic lentivirus using the same protocol.

[0143] Further Study and Bioassay.

[0144] For qualified participants at 30-60 days after transduced T cell infusion, begin testing the efficacy of gene therapy. First, add a CCR5-targeted treatment to their existing CART regimen. This may be the CCR5-blocking drug Maraviroc or the immune suppressing drug Rapamycin that lowers CCR5 receptor density on T cells. The therapeutic lentivirus also targets CCR5 and the combined effects of lentivirus plus Maraviroc or Rapamycin should decrease CCR5 below levels needed to sustain HIV replication.

[0145] Two weeks after adding Maraviroc or Rapamycin, the cART therapy is stopped and participants are monitored closely for HIV virus rebound. If they rebound, cART is reintroduced and managed by their regular physician.

[0146] If HIV virus does not rebound, step-down of the Maraviroc or Rapamycin is started in 2 week intervals until treatment is stopped.

[0147] If viremia does not return within 12-26 weeks, the participant has achieved a functional cure of HIV.

[0148] A database of values relating the number of HIVspecific, lentivirus-transduced CD4 T cells to treatment efficacy (including the frequency of cells with latent HIV infection or other markers) will establish a Gold Standard by which other gene therapy protocols could be judged.

Example 2—Developing an Anti-HIV Lentivirus Vector

[0149] The purpose of this example was to develop an anti-HIV lentivirus vector.

[0150] Inhibitory RNA Designs:

[0151] The sequence of *Homo sapiens* chemokine C-C motif receptor **5** (CCR5) (GC03P046377) mRNA was used to search for potential siRNA or shRNA candidates to knockdown CCR5 levels in human cells. Potential RNA

interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Broad Institute or the BLOCK-iT RNAi Designer from Thermo Scientific. Individual selected shRNA sequences were inserted into lentiviral vectors immediately 3' to a RNA polymerase III promoter such as H1, U6, or 7SK to regulate shRNA expression. These lentivirus-shRNA constructs were used to transduce cells and measure the change in specific mRNA levels. The shRNA most potent for reducing mRNA levels were embedded individually within a microRNA backbone to allow for expression by either the CMV or EF-1alpha RNA polymerase II promoters. The microRNA backbone was selected from mirbase.org/. RNA sequences were also synthesized as synthetic siRNA oligonucleotides and introduced directly into cells without using a lentiviral vector.

[0152] The genomic sequence of Bal strain of human immunodeficiency virus type 1 (HIV-1 85US_BaL, accession number AY713409) was used to search for potential siRNA or shRNA candidates to knockdown HIV replication levels in human cells. Based on sequence homology and experience, the search focused on regions of the Tat and Vif genes of HIV although an individual of skill in the art will understand that use of these regions is non-limiting and other potential targets might be selected. Highly conserved regions of Gag or Polymerase genes could not be targeted by shRNA because these same sequences were present in the packaging system complementation plasmids needed for vector manufacturing. As with the CCR5 (NM 000579.3, NM 001100168.1-specific RNAs, potential HIV-specific RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Gene-E Software Suite hosted by the Broad Institute (broadinstitute.org/mai/public) or the BLOCK-iT RNAi Designer from Thermo Scientific (madesigner.thermofisher. com/rnaiexpress/setOption.

do?designOption=shrna&pid=6712627360706061801).

Individual selected shRNA sequences were inserted into lentiviral vectors immediately 3' to a RNA polymerase III promoter such as H1, U6, or 7SK to regulate shRNA expression. These lentivirus-shRNA constructs were used to transduce cells and measure the change in specific mRNA levels. The shRNA most potent for reducing mRNA levels were embedded individually within a microRNA backbone to allow for expression by either the CMV or EF-1alpha RNA polymerase II promoters

[0153] Vector Constructions:

[0154] For CCR5, Tat or Vif shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by Eurofins MWG Operon, LLC. Overlapping sense and antisense oligonucleotide sequences were mixed and annealed during cooling from 70 degrees Celsius to room temperature. The lentiviral vector was digested with the restriction enzymes BamHI and EcoRI for one hour at 37 degrees Celsius. The digested lentiviral vector was purified by agarose gel electrophoresis and extracted from the gel using a DNA gel extraction kit from Invitrogen. The DNA concentrations were determined and vector to oligo (3:1 ratio) were mixed, allowed to anneal, and ligated. The ligation reaction was performed with T4 DNA ligase for 30 minutes at room temperature. 2.5 microliters of the ligation mix were added to 25 microliters of STBL3 competent bacterial cells. Transformation was achieved after heatshock at 42 degrees Celsius. Bacterial cells were spread on agar plates containing ampicillin and drug-resistant colonies (indicating the presence of ampicillin-resistance plasmids) were recovered, purified and expanded in LB broth. To check for insertion of the oligo sequences, plasmid DNA were extracted from harvested bacteria cultures with the Invitrogen DNA mini prep kit. Insertion of the shRNA sequence in the lentiviral vector was verified by DNA sequencing using a specific primer for the promoter used to regulate shRNA expression. Exemplary vector sequences and cellular elements known to restrict HIV replication can be found in FIGS. **4** and **9**, respectively.

[0155] For example, the shRNA sequences with the highest activity against CCR5, Tat or Vif gene expression were then assembled into a microRNA (miR) cluster under control of the EF-1alpha promoter. The promoter and miR sequences are depicted in FIG. **4**.

[0156] Functional Assays:

[0157] Individual lentivirus vectors containing CCR5, Tat or Vif shRNA sequences and, for experimental purposes, expressing green fluorescent protein (GFP) under control of the CMV Immediate Early Promoter, and designated AGT103/CMV-GFP were tested for their ability to knockdown CCR5, Tat or Vif expression. Mammalian cells were transduced with lentiviral particles either in the presence or absence of polybrene. Cells were collected after 2-4 days; protein and RNA were analyzed for CCR5, Tat or Vif expression. Protein levels were tested by Western blot assay or by labeling cells with specific fluorescent antibodies (CCR5 assay), followed by analytical flow cytometry comparing modified and unmodified cell fluorescence using either the CCR5-specific or isotype control antibodies.

[0158] Starting Testing of Lentivirus:

[0159] T cell culture medium was made using RPMI 1640 supplemented with 10% FBS and 1% penicillin-streptomycin. Cytokine stocks of IL2 10000 units/ml, IL12 1 μ g/ml, IL7 1 μ g/ml, IL15 1 μ g/ml were also prepared in advance. **[0160]** Prior to transduction with the lentivirus, an infectious viral titer was determined and used to calculate the amount of virus to add for the proper multiplicity of infection (MOI).

[0161] Day 0-12: Antigen-Specific Enrichment:

[0162] On day 0, cryopreserved PBMC were thawed, washed with 10 ml 37° C. medium at 1200 rpm for 10 minutes and resuspended at a concentration of 2×10^{6} /ml in 37° C. medium. The cells were cultured at 0.5 ml/well in a 24-well plate at 37° C. in 5% CO₂. To define the optimal stimulation conditions, cells were stimulated with combinations of reagents as listed in Table 2 below:

TABLE 2

 1	2	3	4	5	6
 IL2 + IL12	IL7 + IL15	Peptides + IL2 + IL12	Peptides + IL7 + IL15	MVA + IL2 + IL12	MVA + IL7 + IL15

[0163] Final concentrations: IL2=20 units/ml, IL12=10 ng/ml, IL7=10 ng/ml, IL15=10 ng/ml, peptides=5 µg/ml individual peptide, MVA MOI=1.

[0164] On days 4 and 8, 0.5 ml fresh medium and cytokine at listed concentrations (all concentrations indicate the final concentration in the culture) were added to the stimulated cells.

[0165] Day 12-24: Non-Specific Expansion and Lentivirus Transduction:

[0166] On day 12, the stimulated cells were removed from the plate by pipetting and resuspended in fresh T cell culture medium at a concentration of 1×10^6 /ml. The resuspended cells were transferred to T25 culture flasks and stimulated with DYNABEADS® Human T-Activator CD3/CD28 following the manufacturer's instruction plus cytokine as listed above; flasks were incubated in the vertical position.

[0167] On day 14, AGT103/CMV-GFP was added at MOI 20 and cultures were returned to the incubator for 2 days. At this time, cells were recovered by pipetting, collected by centrifugation at 1300 rpm for 10 minutes, resuspended in the same volume of fresh medium, and centrifuged again to form a loose cell pellet. That cell pellet was resuspended in fresh medium with the same cytokines used in previous steps, with cells at 0.5×10^6 viable cells per ml.

[0168] From days 14 to 23, the number of the cells was evaluated every 2 days and the cells were diluted to 0.5×10^6 /ml with fresh media. Cytokines were added every time.

[0169] On day 24, the cells were collected and the beads were removed from the cells. To remove the beads, cells were transferred to a suitable tube that was placed in the sorting magnet for 2 minutes. Supernatant containing the cells was transferred to a new tube. Cells were then cultured for 1 day in fresh medium at 1×10^6 /ml. Assays were performed to determine the frequencies of antigen-specific T cells and lentivirus transduced cells.

[0170] To prevent possible viral outgrowth, amprenavir (0.5 ng/ml) was added to the cultures on the first day of stimulation and every other day during the culture.

[0171] Examine Antigen-Specific T Cells by Intracellular Cytokine Staining for IFN-Gamma:

[0172] Cultured cells after peptide stimulation or after lentivirus transduction at 1×10^6 cells/ml were stimulated with medium alone (negative control), Gag peptides (5 µg/ml individual peptide), or PHA (5 µg/ml, positive control). After 4 hours, BD GOLGIPLUGTM (1:1000, BD Biosciences) was added to block Golgi transport. After 8 hours, cells were washed and stained with extracellular (CD3, CD4 or CD8; BD Biosciences) and intracellular (IFN-_gamma; BD Biosciences) and intracellular (IFN-_gamma; BD Biosciences) antibodies with BD CYTOFIX/CYTOPERMTM kit following the manufacturer's instruction. Samples were analyzed on a BD FAC-SCALIBURTM Flow Cytometer. Control samples labeled with appropriate isotype-matched antibodies were included in each experiment. Data were analyzed using Flowjo software.

[0173] Lentivirus transduction rate was determined by the frequency of GFP+ cells. The transduced antigen-specific T cells are determined by the frequency of CD3+CD4+GFP+ IFN gamma+ cells; tests for CD3+CD8+GFP+IFN gamma+ cells are included as a control.

[0174] These results indicate that CD4 T cells, the target T cell population, can be transduced with lentiviruses that are designed to specifically knock down the expression of HIV-specific proteins, thus producing an expandable population of T cells that are immune to the virus. This example serves as a proof of concept indicating that the disclosed lentiviral constructs can be used in combination with vaccination to produce a functional cure in HIV patients.

Example 3—CCR5 Knockdown with Experimental Vectors

[0175] AGTc120 is a Hela cell line that stably expresses large amounts of CD4 and CCR5. AGTc120 was transduced with or without LV-CMV-mCherry (the red fluorescent protein mCherry expressed under control of the CMV Immediate Early Promoter) or AGT103/CMV-mCherry. Gene expression of the mCherry fluorescent protein was controlled by a CMV (cytomegalovirus immediate early promoter) expression cassette. The LV-CMV-mCherry vector lacked a microRNA cluster, while AGT103/CMV-mCherry expressed therapeutic miRNA against CCR5, Vif, and Tat. [0176] As shown in FIG. 5A, transduction efficiency was >90%. After 7 days, cells were collected and stained with fluorescent monoclonal antibody against CCR5 and subjected to analytical flow cytometry. Isotype controls are shown in gray on these histograms plotting Mean Fluorescence Intensity of CCR5 APC (x axis) versus cell number normalized to mode (y axis). After staining for cell surface CCR5, cells treated with no lentivirus or control lentivirus (expressing only the mCherry marker) showed no changes in CCR5 density while AGT103 (right section) reduced CCR5 staining intensity to nearly the levels of isotype control. After 7 days, cells were infected with or without R5-tropic HIV reporter virus Bal-GFP. 3 days later, cells were collected and analyzed by flow cytometry. More than 90% of cells were transduced. AGT103-CMV/CMVmCherry reduced CCR5 expression in transduced AGTc120 cells and blocked R5-tropic HIV infection compared with cells treated with the Control vector.

[0177] FIG. **5**B shows the relative insensitivity of transfected AGTc120 cells to infection with HIV. As above, the lentivirus vectors express mCherry protein and a transduced cell that was also infected with HIV (expressing GFP) would appear as a double positive cell in the upper right quadrant of the false color flow cytometry dot plots. In the absence of HIV (upper panels), there were no GFP+ cells under any condition. After HIV infection (lower panels), 56% of cells were infected in the absence of lentivirus transduction and 53.6% of cells became infected in AGTc120 cells transduced with the LV-CMV-mCherry. When cells were transduced with the therapeutic AGT103/CMV-mCherry vector, only 0.83% of cells appeared in the double positive quadrant indicating they were transduced and infected.

[0178] Dividing 53.62 (proportion of double positive cells with control vector) by 0.83 (the proportion of double positive cells with the therapeutic vector) shows that AGT103 provided greater than 65-fold protection against HIV in this experimental system.

Example 4—AGT103 Decreases Expression of Tat and Vif

[0179] Cells were transfected with exemplary vector AGT103/CMV-GFP. AGT103 and other exemplary vectors are defined in Table 3 below.

TABLE 3

Vector Designation	Composition
AGT103	EF1-miR30CCR5-miR21Vif-miR185-Tat-WPRE
Control-mCherry	CMV-mCherry
AGT103/CMV-	CMV-mCherry-EF1-miR30CCR5-miR21Vif-
mCherry	miR185-Tat-WPRE-

TABLE 3-continued

Vector Designation	Composition
Control-GFP AGT103/CMV-GFP	CMV-mCherry CMV-GFP-EF1-miR30CCR5-miR21Vif-miR185- Tat-WPRE-

Abbreviations:

EF-1: elongation factor 1 transcriptional promoter

miR30CCR5—synthetic microRNA capable of reducing CCR5 protein on cell surfaces miR21Vif—synthetic microRNA capable of reducing levels of HIV RNA and Vif protein expression miR185Tat—synthetic micro RNA capable of reducing levels of HIV RNA and Tat protein

expression CMV—Immediate early transcriptional promoter from human cytomegalovirus

mCherry—coding region for the mCherry red fluorescent protein

GFP-coding region for the green fluorescent protein

WPRE-Woodchuck hepatitis virus post transcriptional regulatory element

[0180] A T lymphoblastoid cell line (CEM; CCRF-CEM; American Type Culture Collection Catalogue number CCL119) was transduced with AGT103/CMV-GFP. 48 hours later the cells were transfected with an HIV expression plasmid encoding the entire viral sequence. After 24 hours, RNA was extracted from cells and tested for levels of intact Tat sequences using reverse transcriptase polymerase chain reaction. Relative expression levels for intact Tat RNA were reduced from approximately 850 in the presence of control lentivirus vector, to approximately 200 in the presence of AGT103/CMV-GFP for a total reduction of >4 fold, as shown in FIG. **6**.

[0181] In a similar experiment, HEK 293T cells (Human Embryonic Kidney 293T; American Type Culture Collection Catalogue number CRL-3216) cells were transduced with AGT103/CMV-GFP and then an HIV expression plasmid was transfected into cells 7 days after transduction (control was not transfected with HIV). 24 hours after transfection cells were lysed and analyzed by Western blot using antibodies specific for Actin (cellular loading control) or HIV Vif protein. The presence of AGT103/CMV-GFP (right lane) caused a dramatic reduction in Vif protein expression levels as shown in FIG. **7**.

Example 5—Generating a Population of CD4+ T Cells Enriched for HIV-Specificity and Transduced with AGT103/CMV-GFP

[0182] Therapeutic vaccination against HIV had minimal effect on the distribution of CD4+, CD8+ and CD4+/CD8+ T cells. As shown in FIG. **8**A, the CD4 T cell population is shown in the upper left quadrant of the analytical flow cytometry dot plots, and changes from 52% to 57% of total T cells after the vaccination series. These are representative data.

[0183] Peripheral blood mononuclear cells from a participant in an HIV therapeutic vaccine trial were cultured for 12 days in medium+/-interleukin-2/interleukin-12 or +/-interleukin-7/interleukin-15. Some cultures were stimulated with overlapping peptides representing the entire p55 Gag protein of HIV-1 (JPT PepMix) as a source of epitope peptides for T cell stimulation. These peptides are 10-20 amino acids in length and overlap by 20-50% of their length to represent the entire Gag precursor protein (p55) from HIV-1 BaL strain. The composition and sequence of individual peptides can be adjusted to compensate for regional variations in the predominant circulating HIV sequences or when detailed sequence information is available for an individual patient receiving this therapy. At culture end, cells were recovered

and stained with anti-CD4 or anti-CD8 monoclonal antibodies and the CD3+ population was gated and displayed here. The PepMix stimulation for either pre- or post-vaccination samples was similar to the medium control indicating that PepMix was not toxic to cells and was not acting as a polyclonal mitogen. The results of this analysis can be found in FIG. **8**B.

[0184] PepMix and interleukin-2/interleukin-12 provided for optimal expansion of antigen-specific CD4 T cells. As shown in the upper panels of FIG. **8**C, there was an increase in cytokine (interferon-gamma) secreting cells in post-vaccination specimens exposed to PepMix. In the pre-vaccination sample, cytokine secreting cells increased from 0.43 to 0.69% as a result of exposure to antigenic peptides. In contrast, the post-vaccination samples showed an increase of cytokine secreting cells from 0.62 to 1.76% of total CD4 T cells as a result of peptide stimulation. These data demonstrate the strong impact of vaccination on the CD4 T cell responses to HIV antigen.

[0185] Finally, AGT103/CMV-GFP transduction of antigen-expanded CD4 T cells produced HIV-specific and HIVresistant helper CD4 T cells that are needed for infusion into patients as part of a functional cure for HIV (in accordance with other various aspects and embodiments, AGT103 may be used alone or without further additional elements; for example, clinical embodiments may not include the CMV-GFP segment). The upper panels of FIG. 8D show the results of analyzing the CD4+ T cell population in culture. The x axis of FIG. 8D shows Green Fluorescent Protein (GFP) emission indicating that individual cells were transduced with the AGT103/CMV-GFP. In the post-vaccination samples 1.11% of total CD4 T cells that were both cytokine secreting was recovered, indicating that the cells are responding specifically to HIV antigen, and transduced with AGT103/CMV-GFP. This is the target cell population and the clinical product intended for infusion and functional cure of HIV. With the efficiency of cell expansion during the antigen stimulation and subsequent polyclonal expansion phases of ex vivo culture, 4×10^8 antigen-specific, lentivirus transduced CD4 T cells can be produced. This exceeds the target for cell production by 4-fold and will allow achievement of a count of antigen-specific and HIV-resistant CD4 T cells of approximately 40 cells/microliter of blood or around 5.7% of total circulating CD4 T cells.

[0186] Table 4 below shows the results of the ex vivo production of HIV-specific and HIV-resistant CD4 T cells using the disclosed vectors and methods.

TABLE 4

Material/manipulation	Total CD4 T cells	Percentage HIV- specific	Percentage HIV- specific and HIV-resistant
Leukapheresis pack	\sim 7 × 10 ⁸	~0.12	N/A
Peptide expansion ex	$\sim 8 \times 10^8$	~2.4	N/A
Mitogen expansion Lentivirus transduction	$\sim 1.5 \times 10^{10}$ $\sim 1.5 \times 10^{10}$	~2.4 ~2.4	N/A ~1.6

[0187] While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be

limited to such embodiments. Various modifications may be made thereto without departing from the scope and spirit of the present invention.

What is claimed is:

- 1. A method of treating HIV infection, comprising:
- (a) identifying a subject in need of treatment of HIV infection;
- (b) immunizing the subject with a therapeutically effective amount of an HIV vaccine;
- (c) removing lymphocytes from the subject and purifying peripheral blood mononuclear cells (PBMC);
- (d) contacting the PBMC ex vivo with a therapeutically effective amount of an HIV vaccine;
- (e) transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element;
- (f) culturing the transduced PBMC for about 1 to about 35 days; and
- (g) infusing the transduced PBMC into the subject.

2. The method of claim **1**, wherein step (b) and step (d) comprise the same HIV vaccine.

3. The method of claim **1**, wherein step (b) and step (d) comprise different HIV vaccines.

4. The method of claim **1**, wherein the subject was receiving cART or HAART prior to infusing the transduced PBMC into the subject.

5. The method of claim **1**, wherein the subject receives a cyclophosphamide pre-treatment prior to infusing the transduced PBMC into the subject.

6. The method of claim 1, wherein the at least one genetic element is selected from the group consisting of small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences.

7. The method of claim 6, wherein the small RNA molecules targeting HIV RNA sequences is directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

8. The method of claim **1**, wherein the transduced PBMC are cultured for about 1 to about 10 days prior to infusing the transduced PBMC into the subject.

9. A viral vector for transducing HIV-specific CD4 T cells, wherein the viral vector encodes at least one genetic element selected from the group consisting of small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences.

10. The viral vector of claim 9, wherein the vector is a lentivirus.

11. The viral vector of claim **9**, wherein the vector is a vector-in-vector system.

12. The viral vector of claim **9**, wherein the small RNA molecules targeting HIV RNA sequences is directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

13. A bioassay for determining whether a HIV+ subject is functionally cured, comprising determining the number of HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus, wherein the subject is functionally cured if the number of HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus is above a threshold value after a specified time following treatment according to claim **1**.

14. The method of claim 13, wherein the threshold value is about 1×10^8 HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus.

15. The method of claim **13**, wherein the specified time following treatment is about 30-about 60 days.

16. The method of claim **13**, wherein the specified time following treatment is about 12-about 26 weeks.

17. A method of achieving a functional cure for HIV in a HIV+ subject, comprising:

(a) identifying a subject that is HIV+;

- (b) immunizing the subject with a therapeutically effective amount of an HIV vaccine;
- (c) removing lymphocytes from the subject and purifying peripheral blood mononuclear cells (PBMC);
- (d) contacting the PBMC ex vivo with a therapeutically effective amount of an HIV vaccine;
- (e) transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element;
- (f) culturing the transduced PBMC for about 1-about 21 days; and

(g) infusing the transduced PBMC into the subject, wherein the HIV+ subject achieves a functional cure.

18. The method of claim **18**, wherein step (b) and step (d) comprise the same HIV vaccine.

19. The method of claim **18**, wherein step (b) and step (d) comprise different HIV vaccines.

20. The method of claim **18**, wherein the subject was receiving cART or HAART prior to infusing the transduced PBMC into the subject.

21. The method of claim **18**, wherein the subject receives a cyclophosphamide pre-treatment prior to infusing the transduced PBMC into the subject.

22. The method of claim **18**, wherein the at least one genetic element is selected from the group consisting of small RNA capable of inhibiting the production of chemokine receptor CCR5, small RNA capable of inhibiting the production of chemokine receptor CXCR4, and small RNA molecules targeting HIV RNA sequences.

23. The method of claim **22**, wherein the small RNA molecules targeting HIV RNA sequences is directed to gag, pol, env, tat, rev, nef, vif, vpr, vpu, tev, LTR, TAR, RRE, PE, SLIP, CRS, or INS.

24. The method of claim **18**, wherein the transduced PBMC are cultured for about 1 to about 7 days prior to infusing the transduced PBMC into the subject.

* * * * *