



US 20190201523A1

(19) **United States**(12) **Patent Application Publication****Pauza et al.**(10) **Pub. No.: US 2019/0201523 A1**(43) **Pub. Date: Jul. 4, 2019**(54) **HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY**

8, 2016, provisional application No. 62/385,864, filed on Sep. 9, 2016, provisional application No. 62/409,270, filed on Oct. 17, 2016.

(71) Applicant: **AMERICAN GENE TECHNOLOGIES INTERNATIONAL INC.**, Rockville, MD (US)(72) Inventors: **Charles David Pauza**, Rockville, MD (US); **Haishan Li**, Rockville, MD (US); **Tyler Lahusen**, Rockville, MD (US); **Gary Mansfield**, Rockville, MD (US)(21) Appl. No.: **16/312,056**(22) PCT Filed: **Jul. 7, 2017**(86) PCT No.: **PCT/US2017/041168**

§ 371 (c)(1),

(2) Date: **Dec. 20, 2018****Related U.S. Application Data**

(63) Continuation-in-part of application No. PCT/US2017/013019, filed on Jan. 11, 2017.




(60) Provisional application No. 62/360,185, filed on Jul.

Publication Classification(51) **Int. Cl.****A61K 39/21** (2006.01)**A61P 31/18** (2006.01)**C12N 5/0783** (2006.01)**C12N 15/113** (2006.01)**C12N 15/86** (2006.01)(52) **U.S. Cl.**CPC **A61K 39/21** (2013.01); **A61P 31/18** (2018.01); **C12N 5/0636** (2013.01); **A61K 2039/5156** (2013.01); **C12N 15/86** (2013.01); **C12N 15/1132** (2013.01); **C12N 15/1138** (2013.01)

(57)

ABSTRACT

The present invention relates generally to immunization and immunotherapy for the treatment or prevention of HIV. In particular, the methods include in vivo and/or ex vivo enrichment of HIV-specific CD4+ T cells.

Specification includes a Sequence Listing.**Therapeutic Vector****Helper Plasmid****Envelope Plasmid**




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

Elongation Factor-1 alpha (EF1-alpha) promoter

CCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCTT
TTTCCCAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTTCGCAA
CGGGTTTGCCGCCAGAACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGG
GTTATGGCCCTTGCGTGCCCTGAATTACTTCCACGCCCCCTGGCTGCAGTACGTGATTCTTGATCC
CGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCGCTTAAGGAGCCCCCTTCGCCTC
GTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGCCGCGTGCGAATCTGGTGGCACCTTCGC
GCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAAATTTTGTATGACCTGCTGCGACGCT
TTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTCCGGTTTTTG
GGCCCGCGGGCGGCGACGGGGCCCGTGCGTCCCAGCGCACATGTTCCGGCGAGGCGGGCCGCGA
GCGCGGCCACCGAGAATCGGACGGGGGTAGTCTCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCT
CGCGCCGCGTGTATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCGGTCCGCACCAAGTTGCGTGAG
CGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCGCTCGGGA
GAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTTCCGTCCTCAGCCGTCGCTTCATG
TGACTCCACGGAGTACCGGGCGCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGT
CGTCTTTAGGTTGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACT
GAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCTTTTGTAGTTTGGATC
TTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTTTTTTCTCCATTCAGGTGTCGTGAT
GTACA

miR30 CCR5

AGGTATATTGCTGTTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAGCCACAGATG
GGTAGAGCAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAAGGGGCTT

miR21 Vif


CCCCGGCATCTCCATGGCTGTACCACCTTGTCGGGGGATGTGTACTTCTGAACTTGTGTTGAATC
TCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGGTATCTTTCATCTGACCA

miR185 Tat


GCTAGCGGGCCTGGCTCGAGCAGGGGGCGAGGGATTCCGCTTCTTCCTGCCATAGCGTGGTCCCC
TCCCCTATGGCAGGCAGAAGCGGCACCTTCCCTCCCAATGACCGCGTCTTCGTC

FIG. 7


Vector 1




Vector 2




Vector 3




Vector 4




Vector 5




Vector 6



Vector 7



Vector 8



Vector 9




FIG. 8




FIG. 9A






FIG. 9B

**FIG. 10A****FIG. 10B**




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 14





FIG. 15



FL4-H :: CCR5 APC

FIG. 16




FIG. 17




FIG. 18




FIG. 19A




FIG. 19B




FIG. 19C




FIG. 19C CONTINUED




FIG. 19D




FIG. 19E




FIG. 20A




FIG. 20B




FIG. 20C




FIG. 21A




FIG. 21B

C8166 cells infected by supernatant from HIV-infected Primary CD4 T cells




FIG. 22




FIG. 23




FIG. 24A




FIG. 24B




FIG. 24C




FIG. 24D

HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to: U.S. Provisional Patent Application No. 62/360,185 filed on Jul. 8, 2016 entitled "HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY", U.S. Provisional Patent Application No. 62/385,864 filed on Sep. 9, 2016 entitled "HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY", U.S. Provisional Patent Application No. 62/409,270 filed on Oct. 17, 2016 entitled "HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY," and PCT/US17/13019 filed on Jan. 11, 2017 entitled "HIV PRE-IMMUNIZATION AND IMMUNOTHERAPY", the disclosures of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention relates generally to the field of immunization and immunotherapy for the treatment and prevention of HIV. In particular, the disclosed methods relate to obtaining and processing leukocytes from HIV+ individuals seeking a functional cure to prepare a cell product suitable for infusion to such HIV+ individuals.

BACKGROUND OF THE INVENTION

[0003] Combination antiretroviral therapy (cART) (also known as Highly Active Antiretroviral Therapy or HAART) limits HIV-1 replication and slows disease progression, but drug toxicities and the emergence of drug-resistant viruses are challenges for long-term control in HIV-infected persons. Additionally, traditional antiretroviral therapy, while successful at delaying the onset of AIDS or death, has yet to provide a functional cure. Alternative treatment strategies are needed.

[0004] Intense interest in immunotherapy for HIV infection has been precipitated by emerging data indicating that the immune system has a major, albeit usually insufficient, role in limiting HIV replication. Virus-specific T-helper cells, which are critical to maintenance of cytolytic T cell (CTL) function, likely play a role. Viremia is also influenced by neutralizing antibodies, but they are generally low in magnitude in HIV infection and do not keep up with evolving viral variants in vivo.

[0005] Together these data indicate that increasing the strength and breadth of HIV-specific cellular immune responses may have a clinical benefit through so-called HIV immunotherapy. Some studies have tested vaccines against HIV, but success has been limited to date. Additionally, there has been interest in augmenting HIV immunotherapy by utilizing gene therapy techniques, but as with other immunotherapy approaches, success has been limited.

[0006] Viral vectors can be used to transduce genes into target cells owing to specific virus-host cell interactions and mechanisms for expressing therapeutic gene constructs. As a result, viral vectors have been used as vehicles for the transfer of genes into many different cell types including whole T cells or other immune cells as well as embryos, fertilized eggs, isolated tissue samples, tissue targets in situ and cultured cells. The ability to introduce and express foreign or altered genes in a cell is useful for therapeutic

interventions such as gene therapy, somatic cell reprogramming of induced pluripotent stem cells, and various types of immunotherapy.

[0007] Gene therapy is one of the ripest areas of biomedical research with the potential to create new therapeutics that may involve the use of viral vectors. In view of the wide variety of potential genes available for therapy, an efficient means of delivering these genes is needed to fulfill the promise of gene therapy as a means of treating infectious and non-infectious diseases. Several viral systems including murine retrovirus, adenovirus, parvovirus (adeno-associated virus), coxsackie virus, vaccinia virus, and herpes virus have been proposed as therapeutic gene transfer vectors.

[0008] There are many factors that must be considered when developing viral vectors including tissue tropism, stability of virus preparations, stability and control of expression, genome packaging capacity, construct-dependent vector stability, and whether or not the desired outcome is to have stable gene integration into the host genome. In addition, in vivo application of viral vectors is often limited by host immune responses against viral structural proteins and/or transduced gene products.

[0009] Thus, toxicity and safety are key hurdles that must be overcome for viral vectors to be used in vivo for the treatment of subjects. There are numerous historical examples of gene therapy applications in humans that have met with problems associated with the host immune responses against the gene delivery vehicles or the therapeutic gene products. Viral vectors (e.g., adenovirus) which co-transduce several viral genes together with one or more therapeutic gene(s) are particularly problematic.

[0010] Although lentiviral vectors do not generally induce cytotoxicity and do not elicit strong host immune responses, some lentiviral vectors such as HIV-1, which encode several immunostimulatory gene products, have the potential to cause cytotoxicity and induce strong immune responses in vivo. However, this may not be a concern for lentivirus-derived transducing vectors that do not encode viral genes after transduction. Of course, this may not always be the case, as sometimes the purpose of the vector is to encode a protein that will provoke a clinically useful immune response.

[0011] Another important issue related to the use of lentiviral vectors is that of possible cytopathogenicity upon exposure to some cytotoxic viral proteins. Exposure to certain HIV-1 proteins may induce cell death or functional unresponsiveness in T cells. Likewise, the possibility of generating replication-competent, virulent virus by recombination is often a concern. Accordingly, there remains a need for improved treatments of HIV.

SUMMARY OF THE INVENTION

[0012] In one aspect, a method of treating cells infected with HIV is provided. The method variously includes contacting peripheral blood mononuclear cells (PBMC) isolated from a subject infected with HIV with a therapeutically effective amount of a stimulatory agent, wherein the contacting is carried out ex vivo; transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; and culturing the transduced PBMC for a sufficient period of time to ensure adequate transduction. In embodiments, the transduced PBMC are cultured from about 1 to about 35 days. In embodiments, the method further includes infusing the transduced PBMC into a subject. In embodi-

ments, the method further includes positively selecting HIV-specific CD4⁺ T cells from the PBMC. In further embodiments, the HIV-specific CD4⁺ T cells are positively selected using at least one physical method of selection. In embodiments, the subject is a human. In embodiments, the stimulatory agent includes any agent suitable for stimulating a T cell response in a subject. In embodiments, the stimulatory agent is a peptide or mixture of peptides, and in embodiments includes a gag peptide. In further embodiments, the stimulatory agent is a vaccine. In embodiments, the vaccine is a HIV vaccine, and in embodiments, the HIV vaccine is a MVA/HIV62B vaccine or a variant thereof. In embodiments, the viral delivery system includes a lentiviral particle. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5. In further embodiments, the at least one genetic element includes at least one small RNA capable of targeting a HIV RNA sequence. In further embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. The HIV RNA sequence includes any HIV sequence suitable for targeting by a viral delivery system. In embodiments, the HIV RNA sequence includes one or more of a HIV Vif sequence, a HIV Tat sequence, or a variant thereof. The at least one genetic element includes any genetic element capable of being expressed by a viral delivery system. In embodiments, the at least one genetic element includes a microRNA or a shRNA. In further embodiments, the at least one genetic element comprises a microRNA cluster.

[0013] In another aspect, a method is disclosed that includes obtaining peripheral blood from HIV⁺ individuals; fractionating the blood to obtain a PBMC population; contacting the PBMC population with purified antigen-presenting cells or peptides or proteins representing components of HIV; culturing the contacted PBMC population for about 1 to about 35 days to increase the number of antigen-specific T cells; positively selecting cells that respond to peptide stimulation to produce an enriched cell fraction; transducing the enriched cell fraction ex vivo with a viral delivery system as detailed herein, and culturing the transduced cell fraction for a period of time sufficient to ensure adequate transduction. The order of the method steps disclosed herein can be changed. As a non-limiting example, the steps of positively selecting cells and transducing cells may be reversed and polyclonal stimulation may also be added to improve transduction efficiency.

[0014] In embodiments, the PBMC population is further purified to produce a purified fraction of PBMC. In embodiments, further purified fractions of PBMC are contacted with peptides or proteins representing components of HIV.

[0015] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTG-TAAACTGAGCTTGCTCTACTGTGAAG CCACA-GATGGGTAGAGCAAGCACAGTTTACCGCTGC-CTACTGCCTCGGACTTCAA GGGGCTT (SEQ ID NO: 1). In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0016] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with

CATCTCCATGGCTGTACCACCTTGTCGGGGGATGT-GTACTTCTGAACCTTGTTGA ATCTCATGGAGT-TCAGAAGAACACATCCGCACTGACATTTTGG-TATCTTTCATCTG ACCA (SEQ ID NO: 2); or at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with GGGCCTGGCTCGAGCAGGGGGC-GAGGGATTCCGCTTCTTCCTGCCATAGCGTGG TCCCCTCCCCTATGGCAGGCAGAAGCGGCACCTTC-CCTCCCAATGACCGCGTCTTC GTCG (SEQ ID NO: 3). In embodiments, the at least one genetic element includes SEQ ID NO: 2; or SEQ ID NO: 3.

[0017] In another aspect, the microRNA cluster includes a sequence having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTGTAAACTGAGCTTGCTCTACTGTGAAG CCACAGATGGGTAGAGCAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAA GGGGCTTCCCGGGCATCTCCATGGCTGTAC-CACCTTGTCGGGGGATGTGTACTTCT GAACTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGG TATCTTTCATCTGACCAGCTAGCGGGCCTGGCTC-GAGCAGGGGGCGAGGGATTCC GCTCTTCTCTGC-CATAGCGTGGTCCCCTCCCCTATGGCAGGCA-GAAGCGGCACCTT CCCTCCCAATGACCGCGTCTTCGTC (SEQ ID NO: 31). In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0018] In another aspect, a method of treating HIV infection in a subject is disclosed. The method variously includes immunizing the subject with an effective amount of a first stimulatory agent; removing leukocytes from the subject and obtaining peripheral blood mononuclear cells (PBMC). The method further includes contacting the PBMC ex vivo with a therapeutically effective amount of a second stimulatory agent; transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; and culturing the transduced PBMC for a sufficient period of time to ensure adequate transduction. In embodiments, the transduced PBMC may be cultured from about 1 to about 35 days. In embodiments, the method further includes positively selecting HIV-specific CD4⁺ T cells from the PBMC. In further embodiments, the HIV-specific CD4⁺ T cells are positively selected using at least one physical method of selection. In embodiments, the method further involves infusing the transduced PBMC into a subject. The subject may be a human. The first and second stimulatory agents may be the same or different. The first and second stimulatory agents may include one or more of a peptide or mixture of peptides. In embodiments, at least one of the first and second stimulatory agents includes a gag peptide. In embodiments, at least one of the first and second stimulatory agents includes a mixture of gag peptides that are recognized by immune cells resident in the PBMC or in the purified fractions of PBMC. The at least one of the first and second stimulatory agents may include a vaccine. In embodiments, the vaccine is a HIV vaccine; in further embodiments, the HIV vaccine is a MVA/HIV62B vaccine or a variant thereof. In embodiments, the viral delivery system includes a lentiviral particle. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5. In embodiments, the at least one genetic element includes at least one small RNA capable of targeting an HIV RNA sequence. In embodi-

ments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. The HIV RNA sequence may include a HIV Vif sequence, a HIV Tat sequence, or a variant thereof. The at least one genetic element may include a microRNA or a shRNA. In embodiments, the at least one genetic element comprises a synthetic microRNA cluster designed for simultaneous expression of all elements in the cluster under control of a single transcriptional promoter. In embodiments, the PBMC population is further purified to produce a purified fraction of PBMC.

[0019] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTG-TAAACTGAGCTTGCTCTACTGTGAAG CCACA-GATGGGTAGAGCAAGCACAGTTTACCGCTGC-CTACTGCCTCGGACTTCAA GGGGCTT (SEQ ID NO: 1). In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0020] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with CATCTCCATGGCTGTACCACCTTGTTCGGGGGATGT-GTACTTCTGAACTTGTGTTGA ATCTCATGGAGT-TCAGAAGAACACATCCGCACTGACATTTTGG-TATCTTTCATCTG ACCA (SEQ ID NO: 2); or at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with GGGCCTGGCTCGAGCAGGGGGC-GAGGGATTCCGCTTCTTCCTGCCATAGCGTGG TCCCCTCCCCTATGGCAGGCAGAAGCGGCACCTTC-CCTCCCAATGACCGCGTCTTC GTCG (SEQ ID NO: 3). In embodiments, the at least one genetic element includes SEQ ID NO: 2 or SEQ ID NO: 3.

[0021] In another aspect, the microRNA cluster includes a sequence having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAG CCACAGATGGGTAGAG-CAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAA GGGGCTTCCCGGGCATCTCCATGGCTGTAC-CACCTTGTTCGGGGGATGTGTACTTCT GAACTTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGG TATCTTTCATCTGACCAGCTAGCGGGCCTGGCTC-GAGCAGGGGGCGAGGGATTCC GCTTCTTCCTGC-CATAGCGTGGTCCCCTCCCCTATGGCAGGCA-GAAGCGGCACCTT CCCTCCCAATGACCGCGTCTTCGTC (SEQ ID NO: 31). In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0022] In another aspect, a lentiviral vector is disclosed. The lentiviral vector includes at least one encoded genetic element, wherein the at least one encoded genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5. The at least one encoded genetic element may also comprise at least one small RNA capable of targeting an HIV RNA sequence. In another aspect, the at least one encoded genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. The HIV RNA sequence may include a HIV Vif sequence, a HIV Tat sequence, or a variant

thereof. The at least one encoded genetic element may include a microRNA or a shRNA. The at least one encoded genetic element may include a synthetic microRNA cluster designed for simultaneous expression of all elements in the cluster under control of a single transcriptional promoter.

[0023] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTG-TAAACTGAGCTTGCTCTACTGTGAAG CCACA-GATGGGTAGAGCAAGCACAGTTTACCGCTGC-CTACTGCCTCGGACTTCAA GGGGCTT (SEQ ID NO: 1). In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0024] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with CATCTCCATGGCTGTACCACCTTGTTCGGGGGATGT-GTACTTCTGAACTTGTGTTGA ATCTCATGGAGT-TCAGAAGAACACATCCGCACTGACATTTTGG-TATCTTTCATCTG ACCA (SEQ ID NO: 2); or at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with GGGCCTGGCTCGAGCAGGGGGC-GAGGGATTCCGCTTCTTCCTGCCATAGCGTGG TCCCCTCCCCTATGGCAGGCAGAAGCGGCACCTTC-CCTCCCAATGACCGCGTCTTC GTCG (SEQ ID NO: 3). In embodiments, the at least one genetic element includes SEQ ID NO: 2; or SEQ ID NO: 3.

[0025] In another aspect, the microRNA cluster includes a sequence having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AGGTATATTGCTGTTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAG CCACAGATGGGTAGAG-CAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAA GGGGCTTCCCGGGCATCTCCATGGCTGTAC-CACCTTGTTCGGGGGATGTGTACTTCT GAACTTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGG TATCTTTCATCTGACCAGCTAGCGGGCCTGGCTC-GAGCAGGGGGCGAGGGATTCC GCTTCTTCCTGC-CATAGCGTGGTCCCCTCCCCTATGGCAGGCA-GAAGCGGCACCTT CCCTCCCAATGACCGCGTCTTCGTC (SEQ ID NO: 31). In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0026] In another aspect, a lentiviral vector system for expressing a lentiviral particle is disclosed. The system includes a lentiviral vector as described herein; an envelope plasmid for expressing an envelope protein preferably optimized for infecting a cell; and at least one helper plasmid for expressing genes of interest. In embodiments, the genes of interest include one or more of gag, pol, and rev genes. In embodiments, the lentiviral vector, the envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell line. In further embodiments, a lentiviral particle is produced by the packaging cell line. In embodiments, the lentiviral particle is capable of modulating production of a target of interest. In embodiments, the target of interest is any of chemokine receptor CCR5 or an HIV RNA sequence. The system may further include a first helper plasmid and a second helper plasmid. In embodiments, a first helper plasmid expresses the gag and pol genes, and a second helper plasmid expresses the rev gene.

[0027] In another aspect, a lentiviral particle capable of infecting a cell is provided. The lentiviral particle includes an envelope protein preferably optimized for infecting a cell, and a lentiviral vector as described herein. In embodiments, the envelope protein may be optimized for infecting a T cell. In embodiments, the envelope protein is optimized for infecting a CD4+ T cell.

[0028] In another aspect, a modified cell is provided. The modified cell includes any cell capable of being infected with a lentiviral vector system for use in accordance with present aspects and embodiments. In embodiments, the cell is a CD4+ T cell that is infected with a lentiviral particle. In embodiments, the CD4+ T cell also has been selected to recognize an HIV antigen. In embodiments, the HIV antigen includes a gag antigen. In embodiments, the CD4+ T cell expresses a decreased level of CCR5 following infection with the lentiviral particle.

[0029] In another aspect, a method of selecting a subject for a therapeutic treatment regimen is provided. The method variously includes immunizing the subject with an effective amount of a first stimulatory agent; removing leukocytes from the subject and purifying peripheral blood mononuclear cells (PBMC) and determining a first quantifiable measurement associated with at least one factor associated with the PBMC; contacting the PBMC ex vivo with a therapeutically effective amount of a second stimulatory agent, and determining a second measurement associated with the at least one factor associated with the PBMC, whereby when the second quantifiable measurement is higher than the first quantifiable measurement, the subject is selected for the treatment regimen. The at least one factor may include any factor associated with T cell proliferation or IFN gamma production.

[0030] The foregoing general description and following brief description of the drawings and detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed. Other objects, advantages, and novel features will be readily apparent to those skilled in the art from the following brief description of the drawings and detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0031] FIG. 1 depicts a flow diagram of an ex vivo treatment method of the present disclosure.

[0032] FIG. 2 depicts CD4+ T cell alteration and prevention of new infection in accordance with the present disclosure.

[0033] FIG. 3 depicts an exemplary lentiviral vector system comprised of a therapeutic vector, a helper plasmid, and an envelope plasmid. The therapeutic vector shown here is a preferred therapeutic vector, which is also referred to herein as AGT103, and contains miR30CCR5-miR21Vif-miR185-Tat.

[0034] FIG. 4 depicts an exemplary 3-vector lentiviral vector system in a circularized form.

[0035] FIG. 5 depicts an exemplary 4-vector lentiviral vector system in a circularized form.

[0036] FIG. 6 depicts a further exemplary 3-vector lentiviral vector system in a circularized form.

[0037] FIG. 7 depicts exemplary vector sequences. Positive (i.e., genomic) strand sequences of the promoter and miR cluster were developed for inhibiting the spread of CCR5-tropic HIV strains. Sequences that are not underlined

comprise the EF-1alpha promoter of transcription that was selected as being a preferable promoter for this miR cluster. Sequences that are underlined show the miR cluster consisting of miR30 CCR5, miR21 Vif, and miR185 Tat (as shown collectively in SEQ ID NO: 33).

[0038] FIG. 8 depicts exemplary lentiviral vector constructs according to various aspects of this disclosure.

[0039] FIG. 9 shows knockdown of CCR5 by an experimental vector and corresponding prevention of R5-tropic HIV infection in AGTc120 cells. (A) shows CCR5 expression in AGTc120 cells with or without AGT103 lentivirus vector. (B) shows the sensitivity of transduced AGTc120 cells to infection with a HIV BaL virus stock that was expressing green fluorescent protein (GFP) fused to the Nef gene of HIV.

[0040] FIG. 10 depicts data demonstrating regulation of CCR5 expression by shRNA inhibitor sequences in a lentiviral vector of the present disclosure. (A) Screening data for potential candidates is shown. (B) CCR5 knock-down data following transduction with CCR5 shRNA-1 (SEQ ID NO: 16) is shown.

[0041] FIG. 11 depicts data demonstrating regulation of HIV components by shRNA inhibitor sequences in a lentiviral vector of the present disclosure. (A) Knock-down data for the rev/tat target gene is shown. (B) Knock-down data for the gag target gene is shown.

[0042] FIG. 12 depicts data demonstrating that AGT103 reduces expression of Tat protein expression in cells transfected with an HIV expression plasmid, as described herein.

[0043] FIG. 13 depicts data demonstrating regulation of HIV components by synthetic microRNA sequences in a lentiviral vector of the present disclosure. (A) Tat knock-down data is shown. (B) Vif knock-down data is shown.

[0044] FIG. 14 depicts data demonstrating regulation of CCR5 expression by synthetic microRNA sequences in a lentiviral vector of the present disclosure.

[0045] FIG. 15 depicts data demonstrating regulation of CCR5 expression by synthetic microRNA sequences in a lentiviral vector of the present disclosure containing either a long or short WPRE sequence.

[0046] FIG. 16 depicts data demonstrating regulation of CCR5 expression by synthetic microRNA sequences in a lentiviral vector of the present disclosure with or without a WPRE sequence.

[0047] FIG. 17 depicts data demonstrating regulation of CCR5 expression by a CD4 promoter regulating synthetic microRNA sequences in a lentiviral vector of the present disclosure.

[0048] FIG. 18 depicts data demonstrating detection of HIV Gag-specific CD4 T cells.

[0049] FIG. 19 depicts data demonstrating HIV-specific CD4 T cell expansion and lentivirus transduction. (A) An exemplary schedule of treatment is shown. (B) IFN-gamma production in CD4-gated T cells is shown, as described herein. (C) IFN-gamma production and GFP expression in CD4-gated T cells is shown, as described herein. (D) Frequency of HIV-specific CD4+ T cells is shown, as described herein. (E) IFN-gamma production from PBMCs post-vaccination and following ex vivo peptide and expansion is shown, as described herein.

[0050] FIG. 20 depicts data demonstrating a functional assay for a dose response of increasing AGT103-GFP and inhibition of CCR5 expression. (A) Dose response data for increasing amounts of AGT103-GFP is shown. (B) Nor-

mally distributed populations in terms of CCR5 expression are shown and a left shift of the population average indicates decreasing CCR5 expression due to AGT103-GFP transduction. (C) Percentage inhibition of CCR5 expression with increasing doses of AGT103-GFP is shown.

[0051] FIG. 21 depicts data demonstrating AGT103 transduction efficiency for primary human CD4+ T cells. (A) Frequency of transduced cells (GFP-positive) is shown by FACS, as described herein. (B) Number of vector copies per cell is shown, as described herein.

[0052] FIG. 22 depicts data demonstrating AGT103 inhibition of HIV replication in primary CD4+ T cells, as described herein.

[0053] FIG. 23 depicts data demonstrating AGT103 protection of primary human CD4+ T cells from HIV-induced depletion.

[0054] FIG. 24 depicts data demonstrating generation of a CD4+ T cell population that is highly enriched for HIV-specific, AGT103-transduced CD4 T cells. (A) shows CD4 and CD8 expression profiles for cell populations, as described herein. (B) shows CD4 and CD8 expression profiles for cell populations, as described herein. (C) shows IFN-gamma and CD4 expression profiles for cell populations, as described herein. (D) shows IFN-gamma and GFP expression profiles for cell populations, as described herein.

DETAILED DESCRIPTION

Overview

[0055] Disclosed herein are methods and compositions for treating and/or preventing human immunodeficiency virus (HIV) disease to achieve a functional cure. The methods and compositions include integrating lentivirus, non-integrating lentivirus, and related viral vector technology as described below.

[0056] Disclosed herein are therapeutic viral vectors (e.g., lentiviral vectors), immunotherapies, and methods for their use for treating HIV infection. In embodiments, methods and compositions for achieving a functional cure for HIV infection are provided. As depicted in FIG. 1 herein, the various aspects and embodiments include a first stimulation event, for example a first therapeutic immunization with vaccines intended to produce strong immune responses against HIV in HIV-infected patients, for example with stable suppression of viremia due to daily administration of HAART. In embodiments, the first stimulation event enriches the fraction of HIV-specific CD4 T cells. This is followed by (1) isolating peripheral leukocytes by leukapheresis or purifying PBMC or purifying fractions of PBMC from venous blood, (2) a second stimulating event, for example stimulating CD4 T cells ex vivo with a suitable stimulatory agent, such as any vaccine or protein, for example, HIV or HIV-related peptides, (3) selecting an enriched cell population based on a biological response to peptide stimulation, (4) performing therapeutic lentivirus transduction, ex vivo T cell culture, and (5) re-infusion back into the original patient.

[0057] The various methods and compositions can be used to prevent new cells, such as CD4+ T cells, from becoming infected with HIV. For example as illustrated in FIG. 2, to prevent new cells from becoming infected, CCR5 expression can be targeted to prevent virus attachment. Further, destruction of any residual infecting viral RNA can also be targeted. In respect of the foregoing, and in reference to FIG.

2 herein, compositions and methods are provided to stop the HIV viral cycle in cells that have already become infected with HIV. To stop the HIV viral cycle, viral RNA produced by latently-infected cells, such as latently-infected CD4+ T cells, is targeted.

[0058] Previous efforts to achieve a cure for HIV have fallen short due to, among others, the failure to obtain sufficient numbers of HIV-specific CD4 T cells with protective genetic modifications. When this number is below a critical threshold, a functional cure as described herein is not achieved. For example, upon termination of antiretroviral therapy HIV re-emergence generally follows. Thereafter, patients often experience rapid destruction of HIV-specific CD4 T cells, followed by return to progression of disease despite prior genetic therapy. By employing selective enrichment for HIV-specific T cells in accordance with the compositions and methods described herein, a new HIV treatment regimen has been developed including, in various embodiments, a functional cure.

Definitions and Interpretation

[0059] Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. The methods and techniques of the present disclosure are generally performed according to conventional methods well-known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g.: Sambrook J. & Russell D. *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2000); Ausubel et al., *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*, Wiley, John & Sons, Inc. (2002); Harlow and Lane, *Using Antibodies: A Laboratory Manual*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1998); and Coligan et al., *Short Protocols in Protein Science*, Wiley, John & Sons, Inc. (2003). Any enzymatic reactions or purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art.

[0060] As used herein, the term "about" will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

[0061] As used herein, the terms "administration of" or "administering" an active agent means providing an active agent of the invention to the subject in need of treatment in

a form that can be introduced into that individual's body in a therapeutically useful form and therapeutically effective amount.

[0062] As used herein, the term “AGT103” refers to a particular embodiment of a lentiviral vector that contains a miR30-CCR5/miR21-Vif/miR185-Tat microRNA cluster sequence, as detailed herein.

[0063] As used herein, the term “AGT103T” refers to a cell that has been transduced with a lentivirus that contains the AGT103 lentiviral vector.

[0064] As used herein, the term “CCR5” refers to C-C chemokine receptor 5. Reference herein to “CCR5delta32” is reference to a mutant genotype in the CCR5 gene.

[0065] As used herein, the term “CD” refers to a particular cluster of differentiation protein. A non-limiting example of this terminology as used herein is CD4 protein expression. Examples of such proteins include, but are not limited to CD4.

[0066] As used herein, the term “cART” refers to combination antiretroviral therapy. The term “cART” may be used synonymously with HAART (Highly Active Antiretroviral Therapy).

[0067] Throughout this specification and claims, the word “comprise,” or variations such as “comprises” or “comprising,” will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers. Further, as used herein, the term “includes” means includes without limitation.

[0068] As used herein, the term “engraftment” refers to the ability for one skilled in the art to determine a quantitative level of sustained engraftment in a subject following infusion of a cellular source (see for e.g.: Rosenberg et al., *N Engl. J. Med.* 323:570-578 (1990); Dudley et al., *J. Immunother.* 24:363-373 (2001); Yee et al., *Curr. Opin. Immunol.* 13:141-146 (2001); Rooney et al., *Blood* 92:1549-1555 (1998)).

[0069] The terms, “expression,” “expressed,” or “encodes” refer to the process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. Expression may include splicing of the mRNA in a eukaryotic cell or other forms of post-transcriptional modification or post-translational modification.

[0070] The term “functional cure”, as referenced above, and further defined herein, refers to a state or condition wherein HIV+ individuals who previously required ongoing HIV therapies such as cART or HAART, may survive with low or undetectable virus replication using lower doses, intermittent doses, alternate drug combinations or single agents, or discontinued dosing of such HIV therapies. An individual may be said to have been “functionally cured” while still requiring adjunct therapy to maintain low level virus replication and slow or eliminate disease progression. A possible outcome of a functional cure is the eventual eradication of all or virtually all HIV such that no recurrence is detected within a specified time frame, for example, 1 month, 3 months, 6 months, 1 year, 3 years, and 5 years, and all other time frames as may be defined.

[0071] The term “HIV vaccine” encompasses immunogens plus vehicle plus adjuvant intended to elicit HIV-specific immune responses. The term “HIV vaccine” is within the meaning of the term “stimulatory agent” as described herein. A “HIV vaccine” may include purified or

whole inactivated virus particles that may be HIV or recombinant virus vectors capable of expressing HIV proteins, protein fragments or peptides, glycoprotein fragments or glycopeptides, in addition to recombinant bacterial vectors, plasmid DNA or RNA capable of directing cells to producing HIV proteins, glycoproteins or protein fragments able to elicit specific immunity. Alternately, specific methods for immune stimulation including anti-CD3/CD28 beads, T cell receptor-specific antibodies, mitogens, superantigens, cytokines and other chemical or biological stimuli may be used to activate dendritic, T or B cells for the purposes of enriching HIV-specific CD4 T cells prior to transduction or for in vitro assay of lentivirus-transduced CD4 T cells. Activating substances may be soluble, polymeric assemblies, liposome or endosome-based or linked to beads. Cytokines including interleukin-2, 6, 7, 12, 15, 23 or others may be added to improve cellular responses to stimuli and/or improve the survival of CD4 T cells throughout the culture and transduction intervals. Alternately, and without limiting any of the foregoing, the term “HIV vaccine” encompasses the MVA/HIV62B vaccine and variants thereof. The MVA/HIV62B vaccine is a known highly attenuated double recombinant MVA vaccine. The MVA/HIV62B vaccine was constructed through the insertion of HIV-1 gag-pol and env sequences into the known MVA vector (see: for e.g.: Goepfert et al. (2014) *J. Infect. Dis.* 210(1): 99-110, and see WO2006026667, both of which are incorporated herein by reference). The term “HIV vaccine” also includes any one or more vaccines provided in Table 1, below and in any similar tables contained in the priority documents (all of which are incorporated herein in their entirety).

TABLE 1

IAVI Clinical Trial ID*	Prime**
HVTN 704 AMP	VRC-HIVMAB060-00-AB
VAC89220HPX2004	Ad26.Mos.HIV Trivalent
01-I-0079	VRC4302
04/400-003-04	APL 400-003 GENEVAX-HIV
10-1074	10-1074
87 I-114	gp160 Vaccine (Immuno-AG)
ACTG 326; PACTG 326	ALVAC vCP1452
Ad26.ENVA.01	Ad26.EnvA-01
Ad5HVR48.ENVA.01	Ad5HVR48.ENVA.01
ANRS VAC 02	rgp 160 + peptide V3 ANRS VAC 02
ANRS VAC 04	LIPO-6
ANRS VAC 05	ALVAC vCP125
ANRS VAC 07	ALVAC vCP300
ANRS VAC 08	ALVAC-HIV MN120TMG strain (vCP205)
ANRS VAC 09 bis	LIPO-6
ANRS VAC 12	LPHIV1
ANRS VAC 14	gp160 MN/LAI
ANRS VAC 16	LPHIV1
ANRS VAC 18	LIPO-5
APL 400-003RX101	APL 400-003 GENEVAX-HIV
AVEG 002	HIVAC-1e
AVEG 003	VaxSyn gp160 Vaccine (MicroGeneSys)
AVEG 004	gp160 Vaccine (Immuno-AG)
AVEG 005A/B	Env 2-3
AVEG 006X; VEU 006	MN rgp120
AVEG 007A/B	rgp120/HIV-1 SF-2
AVEG 011	UBI HIV-1 Peptide Immunogen, Multivalent
AVEG 013A	gp160 Vaccine (Immuno-AG)
AVEG 014A/B	TBC-3B
AVEG 017	UBI HIV-1 Peptide Vaccine, Microparticulate Monovalent
AVEG 019	p17/p24:Ty-VLP
AVEG 020	gp120 C4-V3
AVEG 021	P3C541b Lipopeptide
AVEG 022	ALVAC-HIV MN120TMG strain (vCP205)

TABLE 1-continued

IAVI Clinical Trial ID*	Prime**
AVEG 028	Salmonella typhi CVD 908-HIV-1 LAI gp 120
AVEG 031	APL 400-047
AVEG 034/034A	ALVAC vCP1433
C060301	GTU-MultiHIV
C86P1	HIV gp140 ZM96
Cervico-vaginal CN54gp	CN54gp140
140-hsp70 Conjugate Vaccine (TL01)	
CM235 and SF2gp120	CM235 (ThaiE) gp120 plus SF2(B) gp120
CombiHIVvac (KombiVICHvak)	CombiHIVvac
CRC282	P2G12
CUTHIVAC002	DNA-C CN54ENV
DCVax-001	DCVax-001
DNA-4	DNA-4
DP67001	DP67001 DNA
DVP-1	EnvDNA
EN41-UGR7C	EN41-UGR7C
EnvPro	EnvPro
EuroNeut41	EN41-FPA2
EV01	NYVAC-C
EV02 (EuroVacc 02)	DNA-C
Extention HVTN 073E/SAAVI 102	Sub C gp140
F4/AS01	F4/AS01
FIT Biotech	GTU-Nef
Guangxi CDC DNA vaccine	Chinese DNA
HGP-30 memory responses	HGP-30
HIV-CORE002	ChAdV63.HIVconsV
HIV-POL-001	MVA-mBN32
HIVIS 01	HIVIS-DNA
HIVIS 02	MVA-CMDR
HVRF-380-131004	Vichrepol
HVTN 040	AVX101
HVTN 041	rgp120w61d
HVTN 044	VRV-HIVDNA009-00-VP
HVTN 045	pGA2/JS7 DNA
HVTN 048	EP HIV-1090
HVTN 049	Gag and Env DNA/PLG microparticles
HVTN 050/Merck 018	MRKAd5 HIV-1 gag
HVTN 052	VRV-HIVDNA009-00-VP
HVTN 054	VRV-HIVADV014-00-VP
HVTN 055	TBC-M335
HVTN 056	MEP
HVTN 059	AVX101
HVTN 060	HIV-1 gag DNA
HVTN 064	EP HIV-1043
HVTN 065	pGA2/JS7 DNA
HVTN 067	EP-1233
HVTN 070	PENNVAX-B
HVTN 072	VRV-HIVDNA044-00-VP
HVTN 073	SAAVI DNA-C2
HVTN 076	VRV-HIVDNA016-00-VP
HVTN 077	VRV-HIVADV027-00-VP
HVTN 078	NYVAC-B
HVTN 082	VRV-HIVDNA016-00-VP
HVTN 084	VRV-HIVADV054-00-VP
HVTN 086, SAAVI 103	SAAVI MVA-C
HVTN 087	HIV-MAG
HVTN 088	Oligomeric gp140/MF59
HVTN 090	VSV-Indiana HIV gag vaccine
HVTN 092	DNA-HIV-PT123
HVTN 094	GEO-D03
HVTN 096	DNA-HIV-PT123
HVTN 097	ALVAC-HIV vCP1521
HVTN 100	ALVAC-HIV-C (vCP2438)
HVTN 101	DNA-HIV-PT123
HVTN 104	VRV-HIVMAB060-00-AB
HVTN 105	AIDSVAX B/E
HVTN 106	DNA Nat-B env
HVTN 110	Ad4-mgag
HVTN 112	HIV-1 nef/tat/vif, env pDNA vaccine
HVTN 116	VRV-HIVMAB060-00-AB

TABLE 1-continued

IAVI Clinical Trial ID*	Prime**
HVTN 205	pGA2/JS7 DNA
HVTN 702	ALVAC-HIV-C (vCP2438)
HVTN 703 AMP	VRV-HIVMAB060-00-AB
HVTN 908	pGA2/JS7 DNA
IAVI 001	DNA.HIVA
IAVI 016	MVA.HIVA
IAVI A001	tgAAC09
IAVI A003	AAV1-PG9
IAVI B001	Ad35-GRIN/ENV
IAVI B002	Adjuvanted GSK investigational HIV vaccine formulation 1
IAVI B003	Ad26.EnvA-01
IAVI B004	HIV-MAG
IAVI C001	ADVAX
IAVI C002	ADMVA
IAVI D001	TBC-M4
IAVI N004	Ad35-GRIN
HIV-CORE 004	
IAVI R001	rcAd26.MOS1.HIVEnv
IAVI S001	SeV-G
IDEA EV06	DNA-HIV-PT123
IHV01	Full-Length Single Chain (FLSC)
IMPAACT P1112	VRV-HIVMAB060-00-AB
IPCAVD006	MVA mosaic
IPCAVD008	Trimeric gp140
IPCAVD009	Ad26.Mos.HIV Trivalent
ISS P-001	Tat vaccine
LFn-p24 vaccine	LFn-p24
MCA-0835	3BNC117
Mucovac2	CN54gp140
MV1-F4	Measles Vector - GSK
MYM-V101	Virosome-Gp41
NCHECR-AE1	pHIS-HIV-AE
PEACHI-04	ChAdV63.HIVconsV
PedVacc001 &	MVA.HIVA
PedVacc002	
PolyEnv1	PolyEnv1
PXVX-HIV-100-001	Ad4-mgag
RISVAC02	MVA-B
RV 151/WRAIR 984	LFn-p24
RV 158	MVA-CMDR
SG06RS02	HIV gp140 ZM96
TAB9	TAB9
TaMoVac II	HIVIS-DNA
UBI V106	UBI HIV-1 Peptide Vaccine, Microparticulate Monovalent
UCLA MIG-001	TBC-3B
UKHVCSpoke003	DNA - CN54ENV and ZM96GPN
V3-MAPS	V3-MAPS
VAX 002	AIDSVAX B/B
VAX 003	AIDSVAX B/E
VRV 602	VRV-HIVMAB060-00-AB
VRV 607	VRCHIVMAB080-00-AB

*IAVI is the International AIDS Vaccine Initiative, whose clinical trials database is publicly available at <http://www.iaivi.org/trials-database/trials>.

**As used herein, the term "Prime" refers to the composition initially used as an immunological inoculant in a given clinical trial as referenced in Table 1 herein.

[0072] The term "in vivo" refers to processes that occur in a living organism. The term "ex vivo" refers to processes that occur outside of a living organism. For example, in vivo treatment refers to treatment that occurs within a patient's body, while ex vivo treatment is one that occurs outside of a patient's body, but still uses or accesses or interacts with tissues from that patient. Thereafter, an ex vivo treatment step may include a subsequent in vivo treatment step.

[0073] The term "miRNA" refers to a microRNA, and also may be referred to herein as "miR". The term "microRNA cluster" refers to at least two microRNAs that are situated on a vector in close proximity to each other and are co-expressed.

[0074] The term "packaging cell line" refers to any cell line that can be used to express a lentiviral particle.

[0075] The term “PBMC” refers to peripheral blood mononuclear cells.

[0076] The term “percent identity,” in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (e.g., BLASTP and BLASTN or other algorithms available to persons of ordinary skill in the art) or by visual inspection. Depending on the application, the “percent identity” can exist over a region of the sequence being compared, e.g., over a functional domain, or, alternatively, exist over the full length of the two sequences to be compared. For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0077] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel et al., *infra*).

[0078] One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website.

[0079] The percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent identity between two nucleotide or amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

[0080] The nucleic acid and protein sequences of the present disclosure can further be used as a “query sequence” to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of

Altschul, et al. (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, word length=12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to the protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., (BLAST and NBLAST)) can be used. See <http://www.ncbi.nlm.nih.gov>.

[0081] As used herein, “pharmaceutically acceptable” refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues, organs, and/or bodily fluids of human beings and animals without excessive toxicity, irritation, allergic response, or other problems or complications commensurate with a reasonable benefit/risk ratio.

[0082] As used herein, a “pharmaceutically acceptable carrier” refers to, and includes, any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. The compositions can include a pharmaceutically acceptable salt, e.g., an acid addition salt or a base addition salt (see, e.g., Berge et al. (1977) *J Pharm Sci* 66:1-19).

[0083] As used herein, the term “physical method of selection” refers to any physical method that can be used to positively select for a cell type within a larger mixture of cells (e.g., PBMC). A non-limiting example of a physical method of selection is magnetic bead sorting.

[0084] As used herein, the term “SEQ ID NO” is synonymous with the term “Sequence ID No.”

[0085] As used herein, “small RNA” refers to non-coding RNA that are generally less than about 200 nucleotides or less in length and possess a silencing or interference function. In other embodiments, the small RNA is about 175 nucleotides or less, about 150 nucleotides or less, about 125 nucleotides or less, about 100 nucleotides or less, or about 75 nucleotides or less in length. Such RNAs include micro-RNA (miRNA), small interfering RNA (siRNA), double stranded RNA (dsRNA), and short hairpin RNA (shRNA). “Small RNA” of the disclosure should be capable of inhibiting or knocking-down gene expression of a target gene, for example through pathways that result in the destruction of the target gene mRNA.

[0086] As used herein, the term “stimulatory agent” refers to any exogenous agent that can stimulate an immune response, and includes, without limitation, a vaccine, a HIV vaccine, and HIV or HIV-related peptides. A stimulatory agent can preferably stimulate a T cell response.

[0087] As used herein, the term “subject” includes a human patient but also includes other mammals. The terms “subject,” “individual,” “host,” and “patient” may be used interchangeably herein.

[0088] The term “Tat” refers to the HIV tat gene and its gene product, and variants thereof.

[0089] The term “therapeutically effective amount” refers to a sufficient quantity of the active agents of the present invention, in a suitable composition, and in a suitable dosage

form to treat or prevent the symptoms, progression, or onset of the complications seen in patients suffering from a given ailment, injury, disease, or condition. The therapeutically effective amount will vary depending on the state of the patient's condition or its severity, and the age, weight, etc., of the subject to be treated. A therapeutically effective amount can vary, depending on any of a number of factors, including, e.g., the route of administration, the condition of the subject, as well as other factors understood by those in the art.

[0090] As used herein, the term "therapeutic vector" is synonymous with a lentiviral vector such as the AGT103 vector.

[0091] The term "treatment" or "treating" generally refers to an intervention in an attempt to alter the natural course of the subject being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects include, but are not limited to, preventing occurrence or recurrence of disease, alleviating symptoms, suppressing, diminishing or inhibiting any direct or indirect pathological consequences of the disease, ameliorating or palliating the disease state, and causing remission or improved prognosis.

[0092] The term "vaccine", which is used interchangeably with the term "therapeutic vaccine" refers to an exogenous agent that can elicit an immune response in an individual and includes, without limitation, purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, or virus-like particles (VLPs).

[0093] The term "Vif" refers to the HIV vif gene and its gene product, and variants thereof.

Description of Aspects of the Disclosure

[0094] As detailed herein, in one aspect, a method of treating cells infected with HIV is provided. The method generally includes contacting peripheral blood mononuclear cells (PBMC) isolated from a subject infected with HIV with a therapeutically effective amount of a stimulatory agent, wherein the contacting step is carried out ex vivo; transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; and culturing the transduced PBMC for a period of time sufficient to achieve such transduction. In embodiments, the transduced PBMC are cultured from about 1 to about 35 days. In embodiments, the method further includes infusing the transduced PBMC into a subject. In embodiments, the subject is a human. In embodiments, the stimulatory agent is a peptide or mixture of peptides, and in embodiments includes a gag peptide. In further embodiments, the stimulatory includes a vaccine. In embodiments, the vaccine is a HIV vaccine, and in further embodiments, the HIV vaccine is a MVA/HIV62B vaccine or a variant thereof. In embodiments, the viral delivery system includes a lentiviral particle. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5. In embodiments, the at least one genetic element includes at least one small RNA capable of targeting an HIV RNA sequence. In other embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. In embodiments, the HIV RNA sequence includes a HIV Vif sequence, a HIV Tat sequence, or variants thereof. In

embodiments, the at least one genetic element includes at least one of a microRNA or a shRNA. In further embodiments, the at least one genetic element comprises a microRNA cluster.

[0095] In another aspect, a method is disclosed which includes obtaining peripheral blood from HIV+ individuals; fractionating the blood to obtain a PBMC population; contacting the PBMC population with purified antigen-presenting cells or peptides or proteins representing components of HIV; culturing the contacted PBMC population for about 1 to about 12 days to expand an antigen-specific population; positively selecting cells that respond to peptide stimulation to produce an enriched cell fraction; transducing the enriched cell fraction ex vivo with a viral delivery system as detailed herein, and culturing the transduced cell fraction for a period of time sufficient to ensure adequate transduction.

[0096] In embodiments, the PBMC population is further purified to produce a purified fraction of PBMC. In embodiments, further purified fractions of PBMC are contacted with peptides or proteins representing components of HIV.

[0097] In another aspect, the at least one genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 1. In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0098] In another aspect, the at least one genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 2; or at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 3. In embodiments, the at least one genetic element includes SEQ ID NO: 2 or SEQ ID NO: 3.

[0099] In another aspect, the microRNA cluster includes a sequence having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 31. In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0100] In another aspect, a method of manufacturing a cell product for treating HIV infection in a subject is disclosed. The method generally includes obtaining blood leukocytes; removing leukocytes from the subject and purifying peripheral blood mononuclear cells (PBMC) or defined fractions of PBMC. The method further includes contacting the PBMC or purified fraction of PBMC ex vivo with a therapeutically effective amount of a stimulatory agent; positive selection based on response to stimulatory agent to enrich the proportion of antigen-specific T cells; transducing the PBMC or purified fraction of PBMC ex vivo with a viral delivery system encoding at least one genetic element; and culturing the transduced PBMC or a purified fraction of PBMC for a period of time sufficient to achieve transduction and growth of the modified cell population. The method may further include further enrichment of the PBMC, for example, by preferably enriching the PBMC for CD4+ T cells or select-

ing for antigen-specific cells based on cytokine expression or combinations of selection methods to enrich for the therapeutic fraction of cells. In embodiments, the transduced PBMC or purified fraction of PBMC are cultured from about 1 to about 35 days. The method may further involve infusing the transduced PBMC or purified fraction of PBMC into a subject. The subject may be a human. The at least one of the first stimulatory agents may include a peptide or mixture of peptides and may represent one, two, three or more of proteins encoded by the HIV genome. In embodiments, at least one of the first stimulatory agents includes a gag peptide. The at least one of the first stimulatory agents may include a vaccine. In embodiments, the vaccine is a HIV vaccine, and in further embodiments, the HIV vaccine is a MVA/HIV62B vaccine or a variant thereof. In embodiments, the first stimulatory agent is a mixture of gag peptides.

[0101] In embodiments, the viral delivery system includes a lentiviral particle. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5. In embodiments, the at least one genetic element includes at least one small RNA capable of targeting an HIV RNA sequence. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. The HIV RNA sequence may include a HIV Vif sequence, a HIV Tat sequence, or variants thereof. The at least one genetic element may include a microRNA or a shRNA, or a cluster thereof. In embodiments, the at least one genetic element comprises a synthetic microRNA cluster.

[0102] In another aspect, the at least one genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 1. In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0103] In another aspect, the at least one genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 2; or at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 3. In embodiments, the at least one genetic element includes SEQ ID NO: 2 or SEQ ID NO: 3.

[0104] In another aspect, the microRNA cluster includes a sequence having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 31. In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0105] In another aspect, a lentiviral vector is disclosed. The lentiviral vector includes at least one encoded genetic element, wherein the at least one encoded genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5 or at least one small RNA capable of targeting an HIV RNA sequence. In another aspect a lentiviral vector is disclosed in the at least one encoded genetic element comprises a small RNA capable of

inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence. The HIV RNA sequence may include a HIV Vif sequence, a HIV Tat sequence, or a variant thereof. The at least one encoded genetic element may include a microRNA or a shRNA. The at least one encoded genetic element may include a microRNA cluster.

[0106] In another aspect, the at least one genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 1. In embodiments, the at least one genetic element comprises SEQ ID NO: 1.

[0107] In another aspect, the at least one genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 2; or at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 3. In embodiments, the at least one genetic element includes SEQ ID NO: 2; or SEQ ID NO: 3.

[0108] In another aspect, the microRNA cluster includes a sequence having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 31. In embodiments, the microRNA cluster includes SEQ ID NO: 31.

[0109] In another aspect, a lentiviral vector system for expressing a lentiviral particle is provided. The system includes a lentiviral vector as described herein; at least one envelope plasmid for expressing an envelope protein preferably optimized for infecting a cell; and at least one helper plasmid for expressing a gene of interest, for example any of gag, pol, and rev genes, wherein when the lentiviral vector, the at least one envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell, wherein a lentiviral particle is produced by the packaging cell, wherein the lentiviral particle is capable of modulating a target sequence of interest, for example inhibiting production of chemokine receptor CCR5 or targeting an HIV RNA sequence.

[0110] In another aspect, a lentiviral particle capable of infecting a cell is disclosed. The lentiviral particle includes at least one envelope protein preferably optimized for infecting a cell, and a lentiviral vector as described herein. The envelope protein may be optimized for infecting a T cell. In embodiments, the envelope protein is optimized for infecting a CD4+ T cell.

[0111] In another aspect, a modified cell is disclosed. In embodiments, the modified cell is a CD4+ T cell. In embodiments, the CD4+ T cell is infected with a lentiviral particle as described herein. In embodiments, the CD4+ T cell also has been selected to recognize an HIV antigen based on the prior immunization with a stimulatory agent. In a further embodiment, the HIV antigen that is recognized by the CD4+ T cell includes a gag antigen. In a further embodi-

ment, the CD4+ T cell expresses a decreased level of CCR5 following infection with the lentiviral particle.

[0112] In another aspect, a method of selecting a subject for a therapeutic treatment regimen is disclosed. The method generally includes immunizing the subject with an effective amount of a first stimulatory agent; removing leukocytes from the subject and purifying peripheral blood mononuclear cells (PBMC) and determining a first quantifiable measurement associated with at least one factor associated with the PBMC; contacting the PBMC ex vivo with a therapeutically effective amount of a second stimulatory agent, and determining a second measurement associated with the at least one factor associated with the PBMC, whereby when the second quantifiable measurement is different (e.g., higher) than the first quantifiable measurement, the subject is selected for the treatment regimen. The at least one factor may be T cell proliferation or IFN gamma production.

Human Immunodeficiency Virus (HIV)

[0113] Human Immunodeficiency Virus, which is also commonly referred to as “HIV”, is a retrovirus that causes acquired immunodeficiency syndrome (AIDS) in humans. AIDS is a condition in which progressive failure of the immune system allows life-threatening opportunistic infections and cancers to thrive. Without treatment, average survival time after infection with HIV is estimated to be 9 to 11 years, depending upon the HIV subtype and genetics of the host population. Infection with HIV occurs by the transfer of bodily fluids, including but not limited to blood, semen, vaginal fluid, pre-ejaculate, saliva, tears, lymph or cerebro-spinal fluid, or breast milk, or use of contaminated blood or tissue products. HIV may be present in an infected individual as both free virus particles and within infected immune cells.

[0114] HIV infects vital cells in the human immune system such as helper T cells, although tropism can vary among HIV subtypes. Immune cells that may be specifically susceptible to HIV infection include but are not limited to CD4+ T cells, macrophages, and dendritic cells. HIV infection leads to low levels of CD4+ T cells through a number of mechanisms, including but not limited to apoptosis of uninfected bystander cells, direct viral killing of infected cells, and killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections and cancer.

[0115] Structurally, HIV is distinct from many other retroviruses. The RNA genome consists of at least seven structural landmarks (LTR, TAR, RRE, PE, SLIP, CRS, and INS), and at least nine genes (gag, pol, env, tat, rev, nef, vif, vpr, vpu, and sometimes a tenth tev, which is a fusion of tat, env and rev), encoding 19 proteins. Three of these genes, gag, pol, and env, contain information needed to make the structural proteins for new virus particles.

[0116] HIV replicates primarily in CD4 T cells, and causes cellular destruction or dysregulation to reduce host immunity. Because HIV establishes infection as an integrated provirus and may enter a state of latency wherein virus expression in a particular cell decreases below the level for cytopathology affecting that cell or detection by the host immune system, HIV is difficult to treat and has not been eradicated even after prolonged intervals of combination

antiretroviral therapy (cART). In most cases, HIV infection causes fatal disease although survival may be prolonged by cART.

[0117] A major goal in the fight against HIV is to develop strategies for curing disease. Prolonged cART has not accomplished this goal, so investigators have turned to alternative procedures. Early efforts to improve host immunity by therapeutic immunization (using a vaccine after infection has occurred) had marginal or no impact. Likewise, treatment intensification had moderate or no impact.

[0118] Some progress has been made using genetic therapy, but positive results are sporadic and found only among rare human beings carrying defects in one or both alleles of the gene encoding CCR5, which plays a critical role in viral penetration of host cells. However, many investigators are optimistic that genetic therapy holds the best promise for eventually achieving an HIV cure.

[0119] As disclosed herein, the methods and compositions of the invention are able to achieve a functional cure that may or may not include complete eradication of all HIV from the body. As mentioned above, a functional cure is defined as a state or condition wherein HIV+ individuals who previously required cART, may survive with low or undetectable virus replication and using lower or intermittent doses of cART, or are potentially able to discontinue cART altogether. As used herein, a functional cure may still possibly require adjunct therapy to maintain low level virus replication and slow or eliminate disease progression. A possible outcome of a functional cure is the eventual eradication of HIV to prevent all possibility of recurrence.

[0120] The primary obstacles to achieving a functional cure lie in the basic biology of HIV itself. Virus infection deletes CD4 T cells that are critical for nearly all immune functions. Most importantly, HIV infection and depletion of CD4 T cells requires activation of individual cells. Activation is a specific mechanism for individual CD4 T cell clones that recognize pathogens or other molecules, using a rearranged T cell receptor.

[0121] In the case of HIV, infection activates a population of HIV-specific T cells that become infected and are consequently depleted before other T cells that are less specific for the virus, which effectively cripples the immune system's defense against the virus. The capacity for HIV-specific T cell responses is rebuilt during prolonged cART; however, when cART is interrupted the rebounding virus infection repeats the process and again deletes the virus-specific cells, resetting the clock on disease progression.

[0122] Clearly, a functional cure is only possible if enough HIV-specific CD4 T cells are protected to allow for a host's native immunity to confront and control HIV once cART is interrupted. In one embodiment, the present invention provides methods and compositions for improving the effectiveness of genetic therapy to provide a functional cure of HIV disease. In another embodiment, the present invention provides methods and compositions for enhancing host immunity against HIV to provide a functional cure. In yet another embodiment, the present invention provides methods and compositions for enriching HIV-specific CD4 T cells in a patient to achieve a functional cure.

[0123] In one embodiment of the invention, treatment results in enriching a subject's HIV-specific CD4 T cells by about 100%, about 200%, about 300%, about 400%, about 500%, about 600%, about 700%, about 800%, about 900%, or about 1000%.

Gene Therapy

[0124] Viral vectors are used to deliver genetic constructs to host cells for the purposes of disease therapy or prevention.

[0125] Genetic constructs can include, but are not limited to, functional genes or portions of genes to correct or complement existing defects, DNA sequences encoding regulatory proteins, DNA sequences encoding regulatory RNA molecules including antisense, short homology RNA, long non-coding RNA, small interfering RNA or others, and decoy sequences encoding either RNA or proteins designed to compete for critical cellular factors to alter a disease state. Gene therapy involves delivering these therapeutic genetic constructs to target cells to provide treatment or alleviation of a particular disease.

[0126] There are multiple ongoing efforts to utilize genetic therapy in the treatment of HIV disease, but thus far, the results have been poor. A small number of treatment successes were obtained in rare HIV patients carrying a spontaneous deletion of the CCR5 gene (an allele known as CCR5delta32).

[0127] Lentivirus-delivered nucleases or other mechanisms for gene deletion/modification may be used to lower the overall expression of CCR5 and/or help to lower HIV replication. At least one study has reported having success in treating the disease when lentivirus was administered in patients with a genetic background of CCR5delta32. However, this was only one example of success, and many other patients without the CCR5delta32 genotype have not been treated as successfully. Consequently, there is a substantial need to improve the performance of viral genetic therapy against HIV, both in terms of performance for the individual viral vector construct and for improved use of the vector through a strategy for achieving functional HIV cure.

[0128] For example, some existing therapies rely on zinc finger nucleases to delete a portion of CCR5 in an attempt to render cells resistant to HIV infection. However, even after optimal treatment, only 30% of T cells had been modified by the nuclease at all, and of those that were modified, only 10% of the total CD4 T cell population had been modified in a way that would prevent HIV infection. In contrast, the disclosed methods result in virtually every cell carrying a lentivirus transgene having a reduction in CCR5 expression below the level needed to allow HIV infection.

[0129] For the purposes of the disclosed methods, gene therapy can include, but is not limited to, affinity-enhanced T cell receptors, chimeric antigen receptors on CD4 T cells (or alternatively on CD8 T cells), modification of signal transduction pathways to avoid cell death caused by viral proteins, increased expression of HIV restriction elements including TREX, SAMHD1, MxA or MxB proteins, APOBEC complexes, TRIMs-alpha complexes, tetherin (BST2), and similar proteins identified as being capable of reducing HIV replication in mammalian cells.

Immunotherapy

[0130] Historically, vaccines have been a go-to weapon against deadly infectious diseases, including smallpox, polio, measles, and yellow fever. Unfortunately, there is no currently approved vaccine for HIV. The HIV virus has unique ways of evading the immune system, and the human body seems incapable of mounting an effective immune

response against it. As a result, scientists do not have a clear picture of what is needed to provide protection against HIV.

[0131] However, immunotherapy may provide a solution that was previously unaddressed by conventional vaccine approaches. Immunotherapy, also called biologic therapy, is a type of treatment designed to boost the body's natural defenses to fight infections or cancer. It uses materials either made by the body or in a laboratory to improve, target, or restore immune system function.

[0132] In some embodiments of the disclosed invention, immunotherapeutic approaches may be used to enrich a population of HIV-specific CD4 T cells for the purpose of increasing the host's anti-HIV immunity. In some embodiments of the disclosed invention, integrating or non-integrating lentivirus vectors may be used to transduce a host's immune cells for the purposes of increasing the host's anti-HIV immunity. In yet another embodiment of the invention, a vaccine comprising HIV proteins including but not limited to a killed particle, a virus-like particle, HIV peptides or peptide fragments, a recombinant viral vector, a recombinant bacterial vector, a purified subunit or plasmid DNA combined with a suitable vehicle and/or biological or chemical adjuvants to increase a host's immune responses may be used to enrich the population of virus-specific T cells or antibodies, and these methods may be further enhanced through the use of HIV-targeted genetic therapy using lentivirus or other viral vector.

Methods

[0133] In one aspect, the disclosure provides methods for using viral vectors to achieve a functional cure for HIV disease. The methods generally include immunotherapy to enrich the proportion of HIV-specific CD4 T cells, followed by lentivirus transduction to deliver inhibitors of HIV and CCR5 and CXCR4 as required.

[0134] In one embodiment, the methods include a first stimulation event to enrich a proportion of HIV-specific CD4 T cells. The first stimulation can include administration of one or more of any agent suitable for enriching a patient's HIV-specific CD4+ T cells including but not limited to a vaccine.

[0135] Therapeutic vaccines can include one or more HIV protein with protein sequences representing the predominant viral types of the geographic region where treatment is occurring. Therapeutic vaccines will include purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, virus-like particles (VLPs), biological or chemical adjuvants including cytokines and/or chemokines, vehicles, and methods for immunization. Vaccinations may be administered according to standard methods known in the art and HIV patients may continue antiretroviral therapy during the interval of immunization and subsequent ex vivo lymphocyte culture including lentivirus transduction.

[0136] In some embodiments, HIV+ patients are immunized with an HIV vaccine, increasing the frequency of HIV-specific CD4 T cells by about 2, about 25, about 250, about 500, about 750, about 1000, about 1250, or about 1500-fold (or any amount in between these values). The vaccine may be any clinically utilized or experimental HIV vaccine, including the disclosed lentiviral, other viral vectors or other bacterial vectors used as vaccine delivery systems. In another embodiment, the vectors encode virus-like particles (VLPs) to induce higher titers of neutralizing

antibodies. In another embodiment, the vectors encode peptides or peptide fragments associated with HIV including but not limited to gag, pol, and env, tat, rev, nef, vif, vpr, vpu, and tev, as well as LTR, TAR, RRE, PE, SLIP, CRS, and INS. Alternatively, the HIV vaccine used in the disclosed methods may comprise purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, peptides or peptide fragments, virus-like particles (VLPs), or biological or chemical adjuvants including cytokines and/or chemokines.

[0137] In one embodiment, the methods include ex vivo re-stimulation of CD4 T cells from persons or patients previously immunized by therapeutic vaccination, using purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, biological or chemical adjuvants including cytokines and/or chemokines, vehicles, and methods for re-stimulation. Ex vivo re-stimulation may be performed using the same vaccine or immune stimulating compound used for in vivo immunization, or it may be performed using a different vaccine or immune stimulating compound than those used for in vivo immunization. Moreover, in some embodiments, the patient does not require prior therapeutic vaccination or re-stimulation of CD4 T cells if the individual has sufficiently high antigen-specific CD4 T cell responses to HIV proteins. In these embodiments, such a patient may only require ex vivo stimulation of CD4 T cells with viral antigens, vaccines or peptides followed by selection for HIV-specific T cells based on the response to stimulation. Enriched cell preparations may include 1%, 5%, 10%, 20%, 30%, 40%, 50% or more of the HIV-specific CD4⁺ T cells and are used for lentivirus transduction of genes able to protect from HIV-mediated depletion. Stimulation with polyclonal mitogen plus cytokines increases the number of enriched and transduced T cells until appropriate levels are reached for infusion back into the original patient.

[0138] In embodiments, peripheral blood mononuclear cells (PBMCs) are obtained by leukapheresis and treated ex vivo to obtain about 1×10^9 CD4 T cells of which about 0.1%, about 1%, about 5% or about 10% or about 30% or about 40% or about 50% are both HIV-specific in terms of antigen responses, and HIV-resistant by virtue of carrying the therapeutic transgene delivered by the disclosed lentivirus vector. Alternatively, about 1×10^7 , about 1×10^8 , about 1×10^9 , about 1×10^{10} , about 1×10^{11} , or about 1×10^{12} CD4 T cells may be isolated for re-stimulation. Any suitable amount of CD4 T cells are isolated for ex vivo re-stimulation.

[0139] The isolated CD4 T cells can be cultured in appropriate medium throughout re-stimulation with HIV vaccine antigens, which may include antigens present in the prior therapeutic vaccination. Antiretroviral therapeutic drugs including inhibitors of reverse transcriptase, protease or integrase may be added to prevent virus re-emergence during prolonged ex vivo culture. CD4 T cell re-stimulation is used to enrich the proportion of HIV-specific CD4 T cells in culture. The same procedure may also be used for analytical objectives wherein smaller blood volumes with peripheral blood mononuclear cells obtained by purification, are used to identify HIV-specific T cells and measure the frequency of this sub-population.

[0140] The PBMC fraction may be enriched for HIV-specific CD4 T cells by contacting the cells with HIV proteins matching or complementary to the components of the vaccine previously used for in vivo immunization. Ex

vivo re-stimulation can increase the relative frequency of HIV-specific CD4 T cells by about 2, about 5, about 10, about 25, about 50, about 75, about 100, about 125, about 150, about 175, or about 200-fold. Further enrichment is obtained by positive selection for cells responding to HIV antigens, vaccines or peptides. Positive selection is accomplished, for example, with the CliniMACS Cytokine Capture System (Miltenyi Biotec Product number 130-028-701, San Diego, Calif. 92121) or similar manual or automated system including the Miltenyi Prodigy System (Miltenyi Biotec Product number 200-075-301, San Diego, Calif. 92121) that is compatible with selecting viable cells based on expression of a cytokine (including but not limited to interferon gamma or tumor necrosis factor alpha) that is captured by a bi-specific reagent and labeled with a magnetic bead antibody to enable positive selection on a magnetic column. Enrichment may also be accomplished by labeling stimulated cells with antibodies capable of detecting cell surface markers expressed preferentially on activated T cells including CD45RO, MHC Class II and others known in the art. Purification of labeled cells may be by fluorescence activated cell sorting, magnetic bead sorting or other physical methods capable of purifying viable cells based on phenotypic characteristics.

[0141] The methods additionally include combining in vivo therapeutic immunization and ex vivo re-stimulation of CD4 T cells with ex vivo lentiviral transduction and culturing.

[0142] Thus, in one embodiment, the re-stimulated PBMC or fraction of PBMC that has been enriched for HIV-specific CD4 T cells can be positively selected as described above, cultured for 1, 2, 3, 4, 5 or up to 12, 20 or 30 days before activating again with a polyclonal mitogen such as Miltenyi GMP TransAct T cell reagent (Miltenyi Biotec Product number 170-076-156, San Diego, Calif. 92121) or other plant- or fungal based agglutinins or other reagents capable of recognizing cell surface CD3 and CD28 to cross link these molecules and cause polyclonal T cell activation. After polyclonal stimulation cells are transduced with therapeutic anti-HIV lentivirus or other vectors and maintained in culture for a sufficient period of time for such transduction, for example from about 1 to about 21 days, including up to about 35 days. Alternatively, the cells may be cultured for about 1-about 18 days, about 1-about 15 days, about 1-about 12 days, about 1-about 9 days, or about 3-about 7 days. Thus, the transduced cells may be cultured for about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, or about 35 days. Activation with a polyclonal mitogen may or may not be included in the cell product manufacturing process.

[0143] In further embodiments, once the transduced cells have been cultured for a sufficient period of time, transduced CD4 T cells are infused back into the original patient. Infusion can be performed using various devices and methods known in the art. In some embodiments, infusion may be accompanied by pre-treatment with cyclophosphamide or similar compounds to increase the efficiency of re-engraftment.

[0144] In some embodiments, a CCR5-targeted therapy may be added to a subject's antiretroviral therapy regimen,

which was continued throughout the treatment process. Examples of CCR5-targeted therapies include but are not limited to Maraviroc (a CCR5 antagonist) or Rapamycin (immunosuppressive agent that lowers CCR5). In some embodiments, the antiretroviral therapy may be ceased and the subject can be tested for virus rebound. If no rebound occurs, adjuvant therapy can also be removed and the subject can be tested again for virus rebound.

[0145] In various embodiments, continued virus suppression with reduced or no antiretroviral therapy including cART or HAART, and reduced or no adjuvant therapy for about 26 weeks can be considered a functional cure for HIV. Other definitions of a functional cure are described herein.

[0146] The lentiviral and other vectors used in the disclosed methods may encode at least one, at least two, at least three, at least four, or at least five genes, or at least six genes, or at least seven genes, or at least eight genes, or at least nine genes, or at least ten genes, or at least eleven genes, or at least twelve genes of interest. Given the versatility and therapeutic potential of HIV-targeted gene therapy, a viral vector of the invention may encode genes or nucleic acid sequences that include but are not limited to (i) an antibody directed to an antigen associated with an infectious disease or a toxin produced by the infectious pathogen, (ii) cytokines including interleukins that are required for immune cell growth or function and may be therapeutic for immune dysregulation encountered in HIV and other chronic or acute human viral or bacterial pathogens, (iii) factors that suppress the growth of HIV in vivo including CD8 suppressor factors, (iv) mutations or deletions of chemokine receptor CCR5, mutations or deletions of chemokine receptor CXCR4, or mutations or deletions of chemokine receptor CXCR5, (v) antisense DNA or RNA against specific receptors or peptides associated with HIV or host protein associated with HIV, (vi) small interfering RNA against specific receptors or peptides associated with HIV or host protein associated with HIV, or (vii) a variety of other therapeutically useful sequences that may be used to treat HIV or AIDS.

[0147] Additional examples of HIV-targeted gene therapy that can be used in the disclosed methods include, but are not limited to, affinity-enhanced T cell receptors, chimeric antigen receptors on CD4 T cells (or alternatively on CD8 T cells), modification of signal transduction pathways to avoid cell death cause by viral proteins, increased expression of HIV restriction elements including TREX, SAMHD1, MxA or MxB proteins, APOBEC complexes, TRIMS-alpha complexes, tetherin (BST2), and similar proteins identified as being capable of reducing HIV replication in mammalian cells.

[0148] In some embodiments, a patient may be undergoing cART or HAART concurrently while being treated according to the methods of the invention. In other embodiments, a patient may undergo cART or HAART before or after being treated according to the methods of the invention. In some embodiments, cART or HAART is maintained throughout treatment according to the methods of the invention and the patient may be monitored for HIV viral burden in blood and frequency of lentivirus-transduced CD4 T cells in blood. Preferably, a patient receiving cART or HAART prior to being treated according to the methods of the invention is able to discontinue or reduce cART or HAART following treatment according to the methods of the invention.

[0149] For efficacy purposes, the frequency of transduced, HIV-specific CD4 T cells, which is a novel surrogate marker for gene therapy effects, may be determined, as discussed in more detail herein.

Compositions

[0150] In various aspects, the disclosure provides lentiviral vectors capable of delivering genetic constructs to inhibit HIV penetration of susceptible cells. For instance, one mechanism of action in accordance herein is to reduce mRNA levels for CCR5 and/or CXCR4 chemokine receptors for reducing the rates for viral entry into susceptible cells.

[0151] Alternatively, the disclosed lentiviral vectors are capable of inhibiting the formation of HIV-infected cells by reducing the stability of incoming HIV genomic RNA. And in yet another embodiment, the disclosed lentivirus vectors are capable of preventing HIV production from a latently infected cell, wherein the mechanism of action is to cause instability of viral RNA sequences through the action of inhibitory RNA including short-homology, small-interfering or other regulatory RNA species.

[0152] The therapeutic lentiviruses disclosed generally comprise at least one of two types of genetic cargo. First, the lentiviruses may encode genetic elements that direct expression of small RNA capable of inhibiting the production of chemokine receptors CCR5 and/or CXCR4 that are important for HIV penetration of susceptible cells. The second type of genetic cargo includes constructs capable of expressing small RNA molecules targeting HIV RNA sequences for the purpose of preventing reverse transcription, RNA splicing, RNA translation to produce proteins, or packaging of viral genomic RNA for particle production and spreading infection. An exemplary structure is diagrammed in FIG. 3.

[0153] As shown in FIG. 3 (top panel), an exemplary construct may comprise numerous sections or components. For example, in one embodiment, an exemplary LV construct may comprise the following sections or components:

[0154] RSV—a Rous Sarcoma virus long terminal repeat;

[0155] 5'LTR—a portion of an HIV long terminal repeat that can be truncated to prevent replication of the vector after chromosomal integration;

[0156] Psi—a packaging signal that allows for incorporation of the vector RNA genome into viral particles during packaging;

[0157] RRE—a Rev Responsive element can be added to improve expression from the transgene by mobilizing RNA out of the nucleus and into the cytoplasm of cells;

[0158] cPPT—a Poly purine tract that facilitates second strand DNA synthesis prior to integration of the transgene into the host cell chromosome;

[0159] Promoter—a promoter initiates RNA transcription from the integrated transgene to express micro-RNA clusters (or other genetic elements of the construct), and in some embodiments, the vectors may use an EF-1 promoter;

[0160] Anti-CCR5—a micro RNA targeting messenger RNA for the host cell factor CCR5 to reduce its expression on the cell surface;

[0161] Anti-Rev/Tat—a micro RNA targeting HIV genomic or messenger RNA at the junction between

HIV Rev and Tat coding regions, which is sometimes designated miRNA Tat or given a similar description in this application;

[0162] Anti-Vif—a micro RNA targeting HIV genomic or messenger RNA within the Vif coding region;

[0163] WPRE—a woodchuck hepatitis virus post-transcriptional regulatory element is an additional vector component that can be used to facilitate RNA transport of the nucleus; and

[0164] deltaU3 3'LTR—a modified version of a HIV 3' long terminal repeat where a portion of the U3 region has been deleted to improve safety of the vector.

[0165] One of ordinary skill in the art will recognize that the above components are merely examples, and that such components may be reorganized, substituted with other elements, or otherwise changed, so long as the construct is able to prevent expression of HIV genes and decrease the spread of infection.

[0166] Vectors of the invention may include either or both of the types of genetic cargo discussed above (i.e., genetic elements that direct expression of a gene or small RNAs, such as siRNA, shRNA, or miRNA that can prevent translation or transcription), and the vectors of the invention may also encode additionally useful products for the purpose of treatment or diagnosis of HIV. For instance, in some embodiments, these vectors may also encode green fluorescent protein (GFP) for the purpose of tracking the vectors or antibiotic resistance genes for the purposes of selectively maintaining genetically-modified cells in vivo.

[0167] The combination of genetic elements incorporated into the disclosed vectors is not particularly limited. For example, a vector herein may encode a single small RNA, two small RNAs, three small RNA, four small RNAs, five small RNAs, six small RNAs, seven small RNAs, eight small RNAs, nine small RNAs, or ten small RNAs, or eleven small RNAs, or twelve small RNAs. Such vectors may additionally encode other genetic elements to function in concert with the small RNAs to prevent expression and infection of HIV.

[0168] Those of ordinary skill in the art will understand that the therapeutic lentivirus may substitute alternate sequences for the promoter region, targeting of regulatory RNA, and types of regulatory RNA. Further, the therapeutic lentivirus of the disclosure may comprise changes in the plasmids used for packaging the lentivirus particles; these changes are required to increase levels of production in vitro.

Lentiviral Vector System

[0169] A lentiviral virion (particle) in accordance with various aspects and embodiments herein is expressed by a vector system encoding the necessary viral proteins to produce a virion (viral particle). In various embodiments, one vector containing a nucleic acid sequence encoding the lentiviral pol proteins is provided for reverse transcription and integration, operably linked to a promoter. In another embodiment, the pol proteins are expressed by multiple vectors. In other embodiments, vectors containing a nucleic acid sequence encoding the lentiviral Gag proteins for forming a viral capsid, operably linked to a promoter, are provided. In embodiments, this gag nucleic acid sequence is on a separate vector than at least some of the pol nucleic acid

sequence. In other embodiments, the gag nucleic acid is on a separate vector from all the pol nucleic acid sequences that encode pol proteins.

[0170] Numerous modifications can be made to the vectors herein, which are used to create the particles to further minimize the chance of obtaining wild type revertants. These include, but are not limited to deletions of the U3 region of the LTR, tat deletions and matrix (MA) deletions. In embodiments, the gag, pol and env vector(s) do not contain nucleotides from the lentiviral genome that package lentiviral RNA, referred to as the lentiviral packaging sequence.

[0171] The vector(s) forming the particle preferably do not contain a nucleic acid sequence from the lentiviral genome that expresses an envelope protein. Preferably, a separate vector that contains a nucleic acid sequence encoding an envelope protein operably linked to a promoter is used. This env vector also does not contain a lentiviral packaging sequence. In one embodiment the env nucleic acid sequence encodes a lentiviral envelope protein.

[0172] In another embodiment the envelope protein is not from the lentivirus, but from a different virus. The resultant particle is referred to as a pseudotyped particle. By appropriate selection of envelopes one can “infect” virtually any cell. For example, one can use an env gene that encodes an envelope protein that targets an endocytic compartment such as that of the influenza virus, VSV-G, alpha viruses (Semliki forest virus, Sindbis virus), arenaviruses (lymphocytic choriomeningitis virus), flaviviruses (tick-borne encephalitis virus, Dengue virus, hepatitis C virus, GB virus), rhabdoviruses (vesicular stomatitis virus, rabies virus), paramyxoviruses (mumps or measles) and orthomyxoviruses (influenza virus). Other envelopes that can preferably be used include those from Moloney Leukemia Virus such as MLV-E, MLV-A and GALV. These latter envelopes are particularly preferred where the host cell is a primary cell. Other envelope proteins can be selected depending upon the desired host cell. For example, targeting specific receptors such as a dopamine receptor can be used for brain delivery. Another target can be vascular endothelium. These cells can be targeted using a Filovirus envelope. For example, the GP of Ebola, which by post-transcriptional modification become the GP₁ and GP₂ glycoproteins. In another embodiment, one can use different lentiviral capsids with a pseudotyped envelope (for example, FIV or SHIV [U.S. Pat. No. 5,654, 195]). A SHIV pseudotyped vector can readily be used in animal models such as monkeys.

[0173] Lentiviral vector systems as provided herein typically include at least one helper plasmid comprising at least one of a gag, pol, or rev gene. Each of the gag, pol and rev genes may be provided on individual plasmids, or one or more genes may be provided together on the same plasmid. In one embodiment, the gag, pol, and rev genes are provided on the same plasmid (e.g., FIG. 4). In another embodiment, the gag and pol genes are provided on a first plasmid and the rev gene is provided on a second plasmid (e.g., FIG. 5). Accordingly, both 3-vector (e.g., FIGS. 4 and 6) and 4-vector (e.g., FIG. 5) systems can be used to produce a lentivirus as described herein. In embodiments, the therapeutic vector, at least one envelope plasmid and at least one helper plasmid are transfected into a packaging cell, for example a packaging cell line. A non-limiting example of a packaging cell line is the 293T/17 HEK cell line. When the therapeutic vector, the envelope plasmid, and at least one helper plasmid

are transfected into the packaging cell line, a lentiviral particle is ultimately produced.

[0174] In another aspect, a lentiviral vector system for expressing a lentiviral particle is disclosed. The system includes a lentiviral vector as described herein; an envelope plasmid for expressing an envelope protein optimized for infecting a cell; and at least one helper plasmid for expressing gag, pol, and rev genes, wherein when the lentiviral vector, the envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell line, a lentiviral particle is produced by the packaging cell line, wherein the lentiviral particle is capable of inhibiting production of chemokine receptor CCR5 or targeting an HIV RNA sequence.

[0175] In another aspect, the lentiviral vector, which is also referred to herein as a therapeutic vector, includes the following elements: hybrid 5' long terminal repeat (RSV/5' LTR) (SEQ ID NOS: 34-35), Psi sequence (RNA packaging site) (SEQ ID NO: 36), RRE (Rev-response element) (SEQ ID NO: 37), cPPT (polypurine tract) (SEQ ID NO: 38), EF-1 α promoter (SEQ ID NO: 4), miR30CCR5 (SEQ ID NO: 1), miR21Vif (SEQ ID NO: 2), miR185Tat (SEQ ID NO: 3), Woodchuck Post-Transcriptional Regulatory Element (WPPE) (SEQ ID NOS: 32 or 80), and Δ U3 3' LTR (SEQ ID NO: 39). In another aspect, sequence variation, by way of substitution, deletion, addition, or mutation can be used to modify the sequences references herein.

[0176] In another aspect, a helper plasmid includes the following elements: CAG promoter (SEQ ID NO: 41); HIV component gag (SEQ ID NO: 43); HIV component pol (SEQ ID NO: 44); HIV Int (SEQ ID NO: 45); HIV RRE (SEQ ID NO: 46); and HIV Rev (SEQ ID NO: 47). In another aspect, the helper plasmid may be modified to include a first helper plasmid for expressing the gag and pol genes, and a second and separate plasmid for expressing the rev gene. In another aspect, sequence variation, by way of substitution, deletion, addition, or mutation can be used to modify the sequences references herein.

[0177] In another aspect, an envelope plasmid includes the following elements: RNA polymerase II promoter (CMV) (SEQ ID NO: 60) and vesicular stomatitis virus G glycoprotein (VSV-G) (SEQ ID NO: 62). In another aspect, sequence variation, by way of substitution, deletion, addition, or mutation can be used to modify the sequences references herein.

[0178] In various aspects, the plasmids used for lentiviral packaging are modified by substitution, addition, subtraction or mutation of various elements without loss of vector function. For example, and without limitation, the following elements can replace similar elements in the plasmids that comprise the packaging system: Elongation Factor-1 (EF-1), phosphoglycerate kinase (PGK), and ubiquitin C (UbC) promoters can replace the CMV or CAG promoter. SV40 poly A and bGH poly A can replace the rabbit beta globin poly A. The HIV sequences in the helper plasmid can be constructed from different HIV strains or clades. The VSV-G glycoprotein can be substituted with membrane glycoproteins from feline endogenous virus (RD114), gibbon ape leukemia virus (GALV), Rabies (FUG), lymphocytic choriomeningitis virus (LCMV), influenza A fowl plague virus (FPV), Ross River alphavirus (RRV), murine leukemia virus 10A1 (MLV), or Ebola virus (EboV).

[0179] Various lentiviral packaging systems can be acquired commercially (e.g., Lenti-vpak packaging kit from

OriGene Technologies, Inc., Rockville, Md.), and can also be designed as described herein. Moreover, it is within the skill of a person ordinarily skilled in the art to substitute or modify aspects of a lentiviral packaging system to improve any number of relevant factors, including the production efficiency of a lentiviral particle.

Bioassays

[0180] In various aspects, the present invention includes bioassays for determining the success of HIV treatment for achieving a functional cure. These assays provide a method for measuring the efficacy of the disclosed methods of immunization and treatment by measuring the frequency of transduced, HIV specific CD4 T cells in a patient. HIV-specific CD4 T cells are recognizable because, among others, they proliferate, change the composition of cell surface markers, induce signaling pathways including phosphorylation, and/or express specific marker proteins that may be cytokines, chemokines, caspases, phosphorylated signaling molecules or other cytoplasmic and/or nuclear components. Specific responding CD4 T cells are recognized for example, using labeled monoclonal antibodies or specific in situ amplification of mRNA sequences, that allow sorting of HIV-specific cells using flow cytometry sorting, magnetic bead separation or other recognized methods for antigen-specific CD4 T cell isolation. The isolated CD4 T cells are tested to determine the frequency of cells carrying integrated therapeutic lentivirus. Single cell testing methods may also be used including microfluidic separation of individual cells that are coupled with mass spectrometry, PCR, ELISA or antibody staining to confirm responsiveness to HIV and presence of integrated therapeutic lentivirus.

[0181] Thus, in various embodiments, following application of a treatment according to the invention (e.g., (a) immunization, (b) ex vivo leukocyte/lymphocyte culture; (c) re-stimulation with purified proteins, inactivated viruses, virally vectored proteins, bacterially vectored proteins, biological or chemical adjuvants including cytokines and/or chemokines, vehicles; and (d) infusion of the enriched, transduced T cells), a patient may be subsequently assayed to determine the efficacy of the treatment. A threshold value of target T cells in the body may be established to measure a functional cure at a determined value, for example, at about 1×10^8 HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus. Alternatively, the threshold value may be about 1×10^5 , about 1×10^6 , about 1×10^7 , about 1×10^8 , about 1×10^9 , or about 1×10^{10} CD4 T cells in the body of the patient.

[0182] HIV-specific CD4 T cells bearing genetic modification from therapeutic lentivirus can be determined using any suitable method, such as but not limited to flow cytometry, cell sorting, FACS analysis, DNA cloning, PCR, RT-PCR or Q-PCR, ELISA, FISH, western blotting, southern blotting, high throughput sequencing, RNA sequencing, oligonucleotide primer extension, or other methods known in the art.

[0183] While methods for defining antigen specific T cells with genetic modifications are known in the art, utilizing such methods to combine identifying HIV-specific T cells with integrated or non-integrated gene therapy constructs as a standard measure for efficacy is a novel concept in the field of HIV treatment, as described variously herein.

Doses and Dosage Forms

[0184] The disclosed methods and compositions can be used for treating HIV+ patients during various stages of their disease. Accordingly, dosing regimens may vary based upon the condition of the patient and the method of administration.

[0185] In various embodiments, HIV-specific vaccines for the initial in vivo immunization are administered to a subject in need in varying doses. In general, vaccines delivered by intramuscular injection include about 10 µg to about 300 µg, about 25 µg to about 275 µg, about 50 µg to about 250 µg, about 75 µg to about 225, or about 100 µg to about 200 µg of HIV protein, either total virus protein prepared from inactivated virus particles, virus-like particles or purified virus protein from recombinant systems or purified from virus preparations. Recombinant viral or bacterial vectors may be administered by any and all of the routes described. Intramuscular vaccines will include about 1 µg to about 100 µg, about 10 µg to about 90 µg, about 20 µg to about 80 µg, about 30 µg to about 70 µg, about 40 µg to about 60 µg, or about 50 µg of suitable adjuvant molecules and be suspended in oil, saline, buffer or water in volumes of 0.1 to 5 ml per injection dose, and may be soluble or emulsion preparations. Vaccines delivered orally, rectally, buccally, at genital mucosal or intranasally, including some virally-vectored or bacterially-vectored vaccines, fusion proteins, liposome formulations or similar preparations, may contain higher amounts of virus protein and adjuvant. Dermal, sub-dermal or subcutaneous vaccines utilize protein and adjuvant amounts more similar to oral, rectal or intranasal-delivered vaccines. Depending on responses to the initial immunization, vaccination may be repeated 1-5 times using the same or alternate routes for delivery. Intervals may be of 2-24 weeks between immunizations. Immune responses to vaccination are measured by testing HIV-specific antibodies in serum, plasma, vaginal secretions, rectal secretions, saliva or bronchoalveolar lavage fluids, using ELISA or similar methodology. Cellular immune responses are tested by in vitro stimulation with vaccine antigens followed by staining for intracellular cytokine accumulation followed by flow cytometry or similar methods including lymphoproliferation, expression of phosphorylated signaling proteins or changes in cell surface activation markers. Upper limits of dosing may be determined based on the individual patient and will depend on toxicity/safety profiles for each individual product or product lot.

[0186] Immunization may occur once, twice, three times, or repeatedly. For instance, an agent for HIV immunization may be administered to a subject in need once a week, once every other week, once every three weeks, once a month, every other month, every three months, every six months, every nine months, once a year, every eighteen months, every two years, every 36 months, or every three years.

[0187] Immunization will generally occur at least once before ex vivo expansion and enrichment of CD4 T cells, and immunization may occur once, twice, three times, or more after ex vivo leukocyte/lymphocyte culture/re-stimulation and infusion.

[0188] In one embodiment, HIV vaccines for immunization are administered as a pharmaceutical composition. In one embodiment, the pharmaceutical composition comprising an HIV vaccine is formulated in a wide variety of nasal, pulmonary, oral, topical, or parenteral dosage forms for clinical application. Each of the dosage forms can comprise

various disintegrating agents, surfactants, fillers, thickeners, binders, diluents such as wetting agents or other pharmaceutically acceptable excipients. The pharmaceutical composition comprising an HIV vaccine can also be formulated for injection.

[0189] HIV vaccine compositions for the purpose of immunization can be administered using any pharmaceutically acceptable method, such as intranasal, buccal, sublingual, oral, rectal, ocular, parenteral (intravenously, intradermally, intramuscularly, subcutaneously, intracisternally, intraperitoneally), pulmonary, intravaginal, locally administered, topically administered, topically administered after scarification, mucosally administered, via an aerosol, or via a buccal or nasal spray formulation.

[0190] Further, the HIV vaccine compositions can be formulated into any pharmaceutically acceptable dosage form, such as a solid dosage form, tablet, pill, lozenge, capsule, liquid dispersion, gel, aerosol, pulmonary aerosol, nasal aerosol, ointment, cream, semi-solid dosage form, and a suspension. Further, the composition may be a controlled release formulation, sustained release formulation, immediate release formulation, or any combination thereof. Further, the composition may be a transdermal delivery system.

[0191] In another embodiment, the pharmaceutical composition comprising an HIV vaccine is formulated in a solid dosage form for oral administration, and the solid dosage form can be powders, granules, capsules, tablets or pills. In yet another embodiment, the solid dosage form includes one or more excipients such as calcium carbonate, starch, sucrose, lactose, microcrystalline cellulose or gelatin. In addition, the solid dosage form can include, in addition to the excipients, a lubricant such as talc or magnesium stearate. In some embodiments, the oral dosage form is in immediate release or a modified release form. Modified release dosage forms include controlled or extended release, enteric release, and the like. The excipients used in the modified release dosage forms are commonly known to a person of ordinary skill in the art.

[0192] In a further embodiment, the pharmaceutical composition comprising a HIV vaccine is formulated as a sublingual or buccal dosage form. Such dosage forms comprise sublingual tablets or solution compositions that are administered under the tongue and buccal tablets that are placed between the cheek and gum.

[0193] In yet a further embodiment, the pharmaceutical composition comprising an HIV vaccine is formulated as a nasal dosage form. Such dosage forms of the present invention comprise solution, suspension, and gel compositions for nasal delivery.

[0194] In one embodiment, the pharmaceutical composition is formulated in a liquid dosage form for oral administration, such as suspensions, emulsions or syrups. In other embodiments, the liquid dosage form can include, in addition to commonly used simple diluents such as water and liquid paraffin, various excipients such as humectants, sweeteners, aromatics or preservatives. In particular embodiments, the composition comprising HIV vaccine or a pharmaceutically acceptable salt thereof is formulated to be suitable for administration to a pediatric patient.

[0195] In one embodiment, the pharmaceutical composition is formulated in a dosage form for parenteral administration, such as sterile aqueous solutions, suspensions, emulsions, non-aqueous solutions or suppositories. In other embodiments, the non-aqueous solutions or suspensions

includes propyleneglycol, polyethyleneglycol, vegetable oils such as olive oil or injectable esters such as ethyl oleate. As a base for suppositories, witpsol, macrogol, tween 61, cacao oil, laurin oil or glycerinated gelatin can be used.

[0196] The dosage of the pharmaceutical composition can vary depending on the patient's weight, age, gender, administration time and mode, excretion rate, and the severity of disease.

[0197] For the purposes of re-stimulation, lymphocytes, PBMCs, and/or CD4 T cells are generally removed from a patient and isolated for re-stimulation and culturing. The isolated cells may be contacted with the same HIV vaccine or activating agent used for immunization or a different HIV vaccine or activating agent. In one embodiment, the isolated cells are contacted with about 10 ng to 5 μ g of an HIV vaccine or activating agent per about 10^6 cells in culture (or any other suitable amount). More specifically, the isolated cells may be contacted with about 50 ng, about 100 ng, about 200 ng, about 300 ng, about 400 ng, about 500 ng, about 600 ng, about 700 ng, about 800 ng, about 900 ng, about 1 μ g, about 1.5 μ g, about 2 μ g, about 2.5 μ g, about 3 μ g, about 3.5 μ g, about 4 μ g, about 4.5 μ g, or about 5 μ g of an HIV vaccine or activating agent per about 10^6 cells in culture.

[0198] Activating agents or vaccines are generally used once for each in vitro cell culture but may be repeated after intervals of about 15 to about 35 days. For example, a repeat dosing could occur at about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, or about 35 days.

[0199] For transduction of the enriched, re-stimulated cells, the cells may be transduced with lentiviral vectors or with other known vector systems as disclosed, for example, in FIG. 4 or

[0200] FIG. 6. The cells being transduced may be contacted with about 1-1,000 viral genomes (measured by RT-PCR assay of culture fluids containing lentivirus vector) per target cell in culture (or any other suitable amount). Lentivirus transduction may be repeated 1-5 times using the same range of 1-1,000 viral genomes per target cell in culture.

Cellular Enrichment

[0201] In various embodiments, cells such as T cells are obtained from an HIV infected patient and cultured. Culturing can occur in multiwell plates in a culture medium comprising conditioned media ("CM"). The levels of supernatant p24^{gag} ("p24") and viral RNA levels may be assessed by standard means. Those patients whose CM-cultured cells have peak p24 supernatant levels of less than 1 ng/ml may be suitable patients for large-scale T-cell expansion in CM with or without the use of additional anti-viral agents. Additionally, different drugs or drug combinations of interest may be added to different wells and the impact on virus levels in the sample may be assessed by standard means. Those drug combinations providing adequate viral suppression are therapeutically useful combinations. It is within the capacity of a competent technician to determine what constitutes adequate viral suppression in relation to a particular subject. In order to test the effectiveness of drugs of interest in limiting viral expansion, additional factors such as anti-CD3 antibodies may be added to the culture to stimulate viral production. Unlike culture methods for HIV infected cell samples known in the art, CM allows the culture of T

cells for periods of over two months, thereby providing an effective system in which to assay long term drug effectiveness.

[0202] This approach allows the inhibition of gene expression driven by the HIV LTR promoter region in a cell population by the culture of cells in a medium comprising the CM. Culture in CM4 likely inhibits HIV LTR driven gene expression by altering one or more interactions between transcription mediating proteins and HIV gene expression regulatory elements. Transcription-mediating proteins of interest include host cell encoded proteins such as AP-1, NFkappaB, NF-AT, IRF, LEF-1 and Spl, and the HIV encoded protein Tat. HIV gene expression regulatory elements of interest include binding sites for AP-1, NFkappaB, NF-AT, IRF, LEF-1 and Spl, as well as the transacting responsive element ("TAR") which interacts with Tat.

[0203] In a preferred embodiment, the HIV infected cells are obtained from a subject with susceptible transcription mediating protein sequences and susceptible HIV regulatory element sequences. In a more preferred embodiment, the HIV infected cells are obtained from a subject having wild-type transcription-mediating protein sequences and wild-type HIV regulatory sequences.

[0204] Another method of enriching T Cells utilizes immunoaffinity-based selection. This method includes the simultaneous enrichment or selection of a first and second population of cells, such as a CD4+ and CD8+ cell population. Cells containing primary human T cells are contacted with a first immunoaffinity reagent that specifically binds to CD4 and a second immunoaffinity reagent that specifically binds to CD8 in an incubation composition, under conditions whereby the immunoaffinity reagents specifically bind to CD4 and CD8 molecules, respectively, on the surface of cells in the sample. Cells bound to the first and/or the second immunoaffinity reagent are recovered, thereby generating an enriched composition comprising CD4+ cells and CD8+ cells. This approach may include incubation of the composition with a concentration of the first and/or second immunoaffinity reagent that is at a sub-optimal yield concentration. Notably, in some embodiments, transduced cells are a mixed T cell population, and in other embodiments transduced cells are not a mixed T cell population.

[0205] In some embodiments, immunoaffinity-based selection is used where the solid support is a sphere, such as a bead, such as a microbead or nanobead. In other embodiments, the bead can be a magnetic bead. In another embodiment, the antibody contains one or more binding partners capable of forming a reversible bond with a binding reagent immobilized on the solid surface, such as a sphere or chromatography matrix, wherein the antibody is reversibly mobilized to the solid surface. In some embodiments, cells expressing a cell surface marker bound by the antibody on said solid surface are capable of being recovered from the matrix by disruption of the reversible binding between the binding reagent and binding partner. In some embodiments, the binding reagent is streptavidin or is a streptavidin analog or mutant. In some embodiments, immunoaffinity-based selection is used to capture cells specifically responding to HIV proteins, vaccines or peptides based on expression of cytokines in these cells. A bi-specific capture reagent binds to T cells and also captures cytokine released from that cell. The cytokine is preferably interferon gamma but may include tumor necrosis factor alpha or other cytokines known to be produced by T cells. The immobilized cytokine

is recognized by a second immune-affinity reagent that is modified by a magnetic bead. Cells with the first capture reagent-cytokine-second immune-affinity reagent and magnetic bead are retained on a magnetic column and are thus purified away from cells that did not express cytokine after HIV protein, vaccine or peptide stimulation and are maintained in a viable state. Removing the matrix from a magnetic field allows release of the labeled cells and capture as a highly enriched population. In some cases, the enriched cells may be cultured for 1-30 days to increase in number before being stimulated with a polyclonal mitogen such as anti-CD3/anti-CD28 microbeads or similar stimulation reagents that are compatible with lentivirus transduction.

[0206] Stable transduction of primary cells of the hematopoietic system and/or hematopoietic stem cells may be obtained by contacting, in vitro or ex vivo, the surface of the cells with both a lentiviral vector and at least one molecule which binds the cell surface. The cells may be cultured in a ventilated vessel comprising two or more layers under conditions conducive to growth and/or proliferation. In some embodiments, this approach may be used in conjunction with non-CD4+ T cell depletion and/or broad polyclonal expansion.

[0207] In another approach to T cell enrichment, PBMCs are stimulated with a peptide and enriched for cells secreting a cytokine, such as interferon-gamma. This approach generally involves stimulating a mixture of cells containing T cells with antigen, and effecting a separation of antigen-stimulated cells according to the degree to which they are labeled with the product. Antigen stimulation is achieved by exposing the cells to at least one antigen under conditions effective to elicit antigen-specific stimulation of at least one T cell. Labeling with the product is achieved by modifying the surface of the cells to contain at least one capture moiety, culturing the cells under conditions in which the product is secreted, released and specifically bound ("captured" or "entrapped") to said capture moiety; and labeling the captured product with a label moiety, where the labeled cells are not lysed as part of the labeling procedure or as part of the separation procedure. The capture moiety may incorporate detection of cell surface glycoproteins CD3 or CD4 to refine the enrichment step and increase the proportion of antigen-specific T cells in general, of CD4+ T cells in specific.

[0208] The following examples are given to illustrate aspects of the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples. All printed publications referenced herein are specifically incorporated by reference.

EXAMPLES

Example 1: Development of a Lentiviral Vector System

[0209] A lentiviral vector system was developed as summarized in FIG. 3 (linear form) and FIG. 4 (circularized form). Referring first to the top portion of FIG. 3, a representative therapeutic vector has been designed and produced with the following elements being from left to right: hybrid 5' long terminal repeat (RSV/5' LTR) (SEQ ID NOS: 34-35), Psi sequence (RNA packaging site) (SEQ ID NO: 36), RRE (Rev-response element) (SEQ ID NO: 37), cPPT (polypurine tract) (SEQ ID NO: 38), EF-1 α promoter (SEQ ID NO: 4), miR30CCR5 (SEQ ID NO: 1), miR21Vif (SEQ ID NO:

2), miR185Tat (SEQ ID NO: 3), Woodchuck Post-Transcriptional Regulatory Element (WPRE) (SEQ ID NOS: 32 or 80), and Δ U3 3' LTR (SEQ ID NO: 39). The therapeutic vector detailed in FIG. 3 is also referred to herein as AGT103.

[0210] Referring next to the middle portion of FIG. 3, a helper plasmid has been designed and produced with the following elements being from left to right: CAG promoter (SEQ ID NO: 41); HIV component gag (SEQ ID NO: 43); HIV component pol (SEQ ID NO: 44); HIV Int (SEQ ID NO: 45); HIV RRE (SEQ ID NO: 46); and HIV Rev (SEQ ID NO: 47).

[0211] Referring next to the lower portion of FIG. 3, an envelope plasmid has been designed and produced with the following elements being from left to right: RNA polymerase II promoter (CMV) (SEQ ID NO: 60) and vesicular stomatitis virus G glycoprotein (VSV-G) (SEQ ID NO: 62).

[0212] Lentiviral particles were produced in 293T/17 HEK cells (purchased from American Type Culture Collection, Manassas, Va.) following transfection with the therapeutic vector, the envelope plasmid, and the helper plasmid (as shown in FIG. 3). The transfection of 293T/17 HEK cells, which produced functional viral particles, employed the reagent Poly(ethylenimine) (PEI) to increase the efficiency of plasmid DNA uptake. The plasmids and DNA were initially added separately in culture medium without serum in a ratio of 3:1 (mass ratio of PEI to DNA). After 2-3 days, cell medium was collected and lentiviral particles were purified by high-speed centrifugation and/or filtration followed by anion-exchange chromatography. The concentration of lentiviral particles can be expressed in terms of transducing units/ml (TU/ml). The determination of TU was accomplished by measuring HIV p24 levels in culture fluids (p24 protein is incorporated into lentiviral particles), measuring the number of viral DNA copies per cell by quantitative PCR, or by infecting cells and using light (if the vectors encode luciferase or fluorescent protein markers).

[0213] As mentioned above, a 3-vector system (i.e., a 2-vector lentiviral packaging system) was designed for the production of lentiviral particles. A schematic of the 3-vector system is shown in FIG. 4. The schematic of FIG. 4 is a circularized version of the linear system previously described in FIG. 3. Briefly, and with reference to FIG. 4, the top-most vector is a helper plasmid, which, in this case, includes Rev. The vector appearing in the middle of FIG. 4 is the envelope plasmid. The bottom-most vector is the previously described therapeutic vector.

[0214] Referring more specifically to FIG. 4, the Helper plus Rev plasmid includes a CAG enhancer (SEQ ID NO: 40); a CAG promoter (SEQ ID NO: 41); a chicken beta actin intron (SEQ ID NO: 42); a HIV gag (SEQ ID NO: 43); a HIV Pol (SEQ ID NO: 44); a HIV Int (SEQ ID NO: 45); a HIV RRE (SEQ ID NO: 46); a HIV Rev (SEQ ID NO: 47); and a rabbit beta globin poly A (SEQ ID NO: 48).

[0215] The Envelope plasmid includes a CMV promoter (SEQ ID NO: 60); a beta globin intron (SEQ ID NO: 61); a VSV-G (SEQ ID NO: 62); and a rabbit beta globin poly A (SEQ ID NO: 63).

[0216] In an alternate vector system, and with respect to FIG. 6, the vector sequences are provided herein as SEQ ID NOS: 105-107.

[0217] Synthesis of a 2-Vector Lentiviral Packaging System Including Helper (Plus Rev) and Envelope plasmids.

[0218] Materials and Methods:

[0219] Construction of the Helper Plasmid:

[0220] The helper plasmid was constructed by initial PCR amplification of a DNA fragment from the pNL4-3 HIV plasmid (NIH Aids Reagent Program) containing Gag, Pol, and Integrase genes. Primers were designed to amplify the fragment with EcoRI and NotI restriction sites which could be used to insert at the same sites in the pCDNA3 plasmid (Invitrogen). The forward primer was (5'-TAAGCA-GAATTC ATGAATTTGCCAGGAAGAT-3') (SEQ ID NO: 81) and reverse primer was (5'-CCATACAATGAATGGA-CCTAGGCGGCCGACGAAT-3') (SEQ ID NO: 82). The sequence for the Gag, Pol, Integrase fragment was as follows:

(SEQ ID NO: 83)

GAATTCATGAATTTGCCAGGAAGATGGAACCAAAATGATAGGGGGAAT
TGGAGGTTTTATCAAAGTAAGACAGTATGATCAGATACTCATAGAAATCT
GCGGACATAAAGCTATAGGTACAGTATTAGTAGGACCTACACCTGTCAAC
ATAATTGGAAGAAATCTGTTGACTCAGATTGGCTGCACTTTAAATTTTCC
CATTAGTCCTATTGAGACTGTACCAAGTAAATTAAGCCAGGAATGGATG
GCCCCAAAGTTAAACAATGGCCATTGACAGAAGAAAAATAAAGCATT
GTAGAAATTTGTACAGAAATGGAAGGAAGAAAAATTTCAAAATTTGG
GCCTGAAATCCATACAATACTCCAGTATTTGCCATAAAGAAAAAGACA
GTACTAAATGGAGAAATTTAGTAGATTTAGAGAACTTAATAAGAGAAT
CAAGATTTCTGGGAAGTTCAATTAGGAATACCACATCCTGCAGGGTTAA
ACAGAAAAATCAGTAACAGTACTGGATGTGGCGATGCATATTTTTCAG
TTCCCTTAGATAAAGACTTCAGGAAGTATATGCATTTACCATACCTAGT
ATAAACAATGAGACACCAGGGATTAGATATCAGTACAATGTGCTTCCACA
GGGATGGAAGGATCACCAGCAATATTCAGTGTAGCATGACAAAAATCT
TAGAGCCTTTTAGAAAAACAAATCCAGACATAGTCATCTATCAATACATG
GATGATTTGTATGTAGGATCTGACTTAGAAATAGGCGAGCATAGAACAA
AATAGAGGAATGAGACAACTCTGTTGAGGTGGGATTACACACCCAG
ACAAAAACATCAGAAAGAACCTCCATTCCTTTGGATGGGTTATGAATCT
CATCTGTATAATGGACAGTACAGCCTATAGTGTGCGAGAAAGGACAG
CTGGACTGTCAATGACATACAGAAATAGTGGGAAATGAATTGGGCAA
GTCAGATTTATGCAGGATTAAGTAAGGCAATTATGTAACTTCTTAGG
GGAACCAAGCACTAACAGAAGTAGTACCATAACAGAAGAAGCAGAGCT
AGAACTGGCAGAAACAGGGAGATTCTAAAAGAACCGGTACATGGAGTGT
ATTATGACCCATCAAAGACTTAATAGCAGAAATACAGAAGCAGGGGCAA
GGCCAAATGGACATATCAAATTTATCAAGAGCCATTTAAAAATCTGAAAC
AGGAAAGTATGCAAGAAATGAAGGGTGCACACCTAATGATGTGAAACAAT
TAACAGAGGCAGTACAAAAATAGCCACAGAAAGCATAGTAATATGGGGA
AAGACTCCTAAATTTAAATTACCCATACAAAAGGAACATGGGAAGCATG

-continued

GTGGACAGAGTATTGGCAAGCCACCTGGATTCCCTGAGTGGGAGTTTGTC
ATACCCCTCCCTTAGTGAAGTTATGGTACCAGTTAGAGAAAGAACCCATA
ATAGGAGCAGAACTTTCTATGTAGATGGGCAGCCAATAGGGAACTAA
ATTAGGAAAAGCAGGATATGTAAGTACAGAGGAAGACAAAAGTTGTCC
CCCTAACGACACACAATCAGAAGACTGAGTTACAAGCAATTCATCTA
GCTTTGCAGGATTCGGGATTAGAAGTAAACATAGTGACAGACTCACAATA
TGCAATTGGGAATCATTCAGCACAACCAGATAAGAGTGAATCAGAGTTAG
TCAGTCAAATAATAGAGCAGTTAATAAAAAAGGAAAAAGTCTACCTGGCA
TGGGTACCAGCACACAAAGGAATGGAGGAAATGAACAGTAGATAAAT
GGTCAGTGTGGAATCAGGAAAGTACTATTTTTAGATGGAATAGATAAGG
CCCCAAGAAGACATGAGAAATATCACAGTAATTGGAGAGCAATGGCTAGT
GATTTTAACCTACCACCTGTAGTAGCAAAAGAAATAGTAGCCAGCTGTGA
TAAATGTCAGCTAAAAGGGGAAGCCATGCATGGACAAGTAGACTGTAGCC
CAGGAATATGGCAGCTAGATTGTACACATTTAGAAGGAAAAGTTATCTTG
GTAGCAGTTTATGTAGCCAGTGGATATATAGAAGCAGAAGTAATCCAGC
AGAGACAGGGCAAGAAACAGCATACTTCCCTCTTAAATTAGCAGGAAGAT
GGCCAGTAAAAACAGTACATACAGACAATGGCAGCAATTTACCACTACT
ACAGTTAAGGCCCGCTGTTGGTGGCGGGGATCAAGCAGGAATTTGGCAT
TCCCTACAATCCCCAAAGTCAAGGAGTAATAGAATCTATGAATAAAGAA
TAAAGAAATTTATAGGACAGGTAAGAGATCAGGCTGAACATCTTAAGACA
GCAGTACAAATGGCAGTATTCATCCACAATTTTAAAGAAAAGGGGGAT
TGGGGGTACAGTGCAGGGGAAAGAAATAGTAGACATAATAGCAACAGACA
TACAACTAAGAATTACAAAAACAAATTACAAAAATTCAAAAATTTTCGG
GTTTATTACAGGGACAGCAGAGATCCAGTTTGGAAAGGACAGCAAGCT
CCTCTGGAAGGTGAAGGGCAGTAGTAATACAAGATAATAGTGACATAA
AAGTAGTGCCAGAAGAAAGCAAAGATCATCAGGATTATGGAACACAG
ATGGCAGGTGATGATTGTGTGGCAAGTAGACAGGATGAGGATTAA

[0221] Next, a DNA fragment containing the Rev, RRE, and rabbit beta globin poly A sequence with XbaI and XmaI flanking restriction sites was synthesized by MWG Operon. The DNA fragment was then inserted into the plasmid at the XbaI and XmaI restriction sites. The DNA sequence was as follows:

(SEQ ID NO: 84)

TCTAGAATGGCAGGAAGAAGCGGAGACAGCGACGAAGAGCTCATCAGAAC
AGTCAGACTCATCAAGCTTCTCTATCAAGCAACCCACTCCCAATCCCG
AGGGGACCCGACAGGCCCGAAGGAATAGAAGAAGAGTGGAGAGAGAGA
CAGAGACAGATCCATTGATTAGTGAACGGATCCTTGGCACTTATCTGGG
ACGATCTGCGGAGCCTGTGCTCTTACAGTACCACCGCTTGAGAGACTTA
CTCTTGATTGTAACGAGGATTGTGGAACTTCTGGACGCAGGGGTGGGA
AGCCCTCAAATATTGGTGAATCTCCTACAATATTGGAGTACAGGAGCTAA

-continued

AGAATAGAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACT
 ATGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTC
 TGGTATAGTGCAGCAGCAGAACAAATTTGCTGAGGGCTATTGAGGCGCAAC
 AGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGA
 ATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTAGATCTTTT
 TCCCCTCTGCCAAAATTATGGGACATCATGAAGCCCCCTTGAGCATCTGA
 CTTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTGGAAAT
 TTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTAAA
 ACATCAGAATGAGTATTGTTTAGAGTTTGGCAACATATGCCATATGCT
 GGCTGCCATGAACAAAGGTGGCTATAAAGAGGTCAATCAGTATATGAAACA
 GCCCCCTGCTGTCCATTCTTATTCATAGAAAAGCCTTGACTTGAGGTT
 AGATTTTTTTTATATTTTGTGTTTATTTTTCTTTAACATCCCTA
 AAATTTCTCTTACATGTTTACTAGCCAGATTTTCTCTCTCTGACT
 ACTCCCAGTCATAGCTGTCCCTCTTCTCTTATGAAGATCCCTCGACCTGC
 AGCCCAAGCTTGCGTAATCATGGTCATAGCTGTTCTGTGTGAAATTG
 TTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
 AAGCCTGGGGTGCCATAATGAGTGAGCTAACTCACATTAATTGCGTTGCGC
 TCACTGCCCGCTTTCCAGTCGGGAAACCTGTCTGCCAGCGGATCCGCAT
 CTAATTAGTCAGCAACCATAGTCCCGCCCCTAACCTCGCCCATCCGCGC
 CCTAACTCCGCCAGTTCCGCCCATTTCTCCGCCCATGGCTGACTAATTT
 TTTTATTTATGACAGAGCCGAGGCGCCTCGGCTCTGAGCTATTCCAG
 AAGTAGTGAGGAGGCTTTTGGAGGCTAGGCTTTGCAAAAAGCTAAC
 TTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAA
 TTTCAAAATAAAGCATTTTTTCTACTGCATTCTAGTTGTGGTTGTCCA
 AACTCATCAATGTATCTTATCAGCGCCGCCCGGG

[0222] Finally, the CMV promoter of pCDNA3.1 was replaced with the CAG enhancer/promoter plus a chicken beta actin intron sequence. A DNA fragment containing the CAG enhancer/promoter/intron sequence with MluI and EcoRI flanking restriction sites was synthesized by MWG Operon. The DNA fragment was then inserted into the plasmid at the MluI and EcoRI restriction sites. The DNA sequence was as follows:

(SEQ ID NO: 85)
 ACGCGTTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCC
 CATATATGGAGTTCGCGTTACATAACTTACGGTAAATGGCCCGCTGGC
 TGACCGCCCAACGACCCCGCCCATTTGACGTCAATAATGACGTATGTTCC
 CATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGACTATT
 TACGGTAAACTGCCCACTTGGCAGTACATCAAGTGATCATATGCCAAGT
 ACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGC
 CCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTAT

-continued

TAGTCATCGCTATTACCATGGGTGCGAGGTGAGCCCCAGTTCTGCTTCAC
 TCTCCCCATCTCCCCCCTCCCCACCCCAATTTTGTATTTATTTATTT
 TTTAATTATTTTGTGTCAGCGATGGGGCGGGGGGGGGGGGGCGCGCGCC
 AGGCGGGGCGGGGCGGGGCGAGGGGCGGGGCGGGGCGAGGCGGAGAGGTG
 CGGCGGCAGCCAATCAGAGCGGCGCGCTCCGAAAGTTTCTTTTATGGCG
 AGGCGGCGGGGCGGGCGGCGCTATAAAAAGCGAAGCGCGCGGGCGGGG
 AGTCGCTGCGTTGCTTTCGCCCCGTGCCCGCTCCGCGCGGCTCGCGCC
 GCCCGCCCCGGCTCTGACTGACCGCGTTACTCCACAGGTGAGCGGGCGG
 GACGGCCCTTCTCTCCGGGCTGTAATTAGCGCTTGTTTAAATGACGGCT
 CGTTTCTTTTCTGTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCC
 CTTTGTGCGGGGGGAGCGGCTCGGGGGGTGCGTGCCTGTGTGTGTGCGT
 GGGGAGCGCCGCGTGCAGCGCGCGCTGCCCGCGGCTGTGAGCGCTGCGG
 GCGCGGCGCGGGGCTTTGTGCGCTCCGCGTGTGCGGAGGGGAGCGCGGC
 CGGGGGCGGTGCCCGCGGTGCGGGGGGCTGCGAGGGGAACAAAGGCTG
 CGTGCGGGGTGTGTGCGTGGGGGGGTGAGCAGGGGTTGTTGGCGCGCGG
 TCGGGCTGTAACCCCCCTGCACCCCTCCCGAGTTGCTGAGCACGG
 CCGGGCTTCGGGTGCGGGGCTCCGTGCGGGGCGTGGCGGGGGCTCGCCG
 TGCCGGGCGGGGGGTGCGCGCAGGTGGGGGTGCCGGGCGGGGCGGGGCGG
 CCTCGGGCGGGGAGGGGCTCGGGGAGGGGCGCGCGGCGCCCGAGCGCC
 GGCGGCTGTGAGGCGCGGCGAGCCGAGCCATTGCCTTTATGGTAATC
 GTGCGAGAGGGCGCAGGACTTCTTTTGTCCCAATCTGGCGGAGCCGAA
 ATCTGGGAGGCGCGCGCACCCCTCTAGCGGGCGCGGGCGAAGCGGTG
 CGGCGCCGGCAGGAAGGAAATGGGCGGGGAGGGCTTCGTGCGTCCCGC
 GCGCGCTCCCTTCTCCATCTCCAGCCTCGGGGCTGCCGAGGGGGAGC
 GCTGCTTCGGGGGGGACGGGGCAGGGCGGGGTTGCGCTTCTGGCGTGTG
 ACCGGCGGGAATTC

[0223] Construction of the VSV-G Envelope Plasmid:

[0224] The vesicular stomatitis Indiana virus glycoprotein (VSV-G) sequence was synthesized by MWG Operon with flanking EcoRI restriction sites. The DNA fragment was then inserted into the pCDNA3.1 plasmid (Invitrogen) at the EcoRI restriction site and the correct orientation was determined by sequencing using a CMV specific primer. The DNA sequence was as follows:

(SEQ ID NO: 86)
 GAATTCATGAAGTGCTTTTGTACTTAGCCTTTTATTGCGGGTGAA
 TTGCAAGTTCACCATAGTTTTTCCACACAACCAAAAGGAACTGGAAAA
 ATGTTCTTCTAATTACCATATTGCGCGTCAAGCTCAGATTTAAATTGG
 CATAATGACTTAATAGGCACAGCCTTACAAGTCAAAATGCCCAAGAGTCA
 CAAGGCTATTCAAGCAGACGGTTGGATGTGTATGCTTCCAAATGGGTCA
 CTACTTGTGATTTCCGCTGGTATGGACCGAAGTATATAACACATTCCATC

-continued

CGATCCTTCACTCCATCTGTAGAACAAATGCAAGGAAAGCATTGAACAAAC
GAAACAAGGAACCTGGCTGAATCCAGGCTTCCCTCCTCAAAGTTGTGGAT
ATGCAACTGTGACGGATGCCAAGCAGTGATTGTCCAGGTGACTCCTCAC
CATGTGCTGGTTGATGAATACACAGGAGAATGGGTGATTACAGTTTCAT
CAACGGAAAAATGCAGCAATTACATATGCCCCACTGTCCATAACTCTACAA
CCTGGCATTCTGACTATAAGGTCAAAGGGCTATGTGATTCTAACCTCATT
TCCATGGACATCACCTTCTTCTCAGAGGACGGAGAGCTATCATCCCTGGG
AAAGGAGGGCACAGGGTTGAGAAGTAAGTACTTTGCTTATGAACTGGAG
GCAAGGCCTGCAAAATGCAATACTGCAAGCATTGGGGAGTCAGACTCCCA
TCAGGTGTCTGGTTCGAGATGGCTGATAAGGATCTCTTTGCTGCGCCAG
ATTCCTTGAATGCCCAGAGGGTCAAGTATCTCTGCTCCATCTCAGACCT
CAGTGGATGTAAGTCTAATTCAGGACGTTGAGAGGATCTTGGATTATTC
CTCTGCCAAGAACTGGAGCAAAATCAGAGCGGTCTTCCAATCTCTCC
AGTGGATCTCAGCTATCTTGCTCTAAAAACCAGGAACCGGTCTTGCTT
TCACCATAATCAATGGTACCCTAAATACTTTGAGACCAGATACATCAGA
GTCGATATTGCTGCTCCAATCTCTCAAGAATGGTCGAATGATCAGTGG
AACTACCACAGAAAGGGAACGTGGGATGACTGGGCACCATATGAAGACG
TGGAAATTGGACCCAATGGAGTTCTGAGGACCAAGTTGAGATATAAGTTT
CCTTTATACATGATTGGACATGGTATGTTGGACTCCGATCTTCATCTTAG
CTCAAAGGCTCAGGTGTTGCAACATCTCACATTCAAGACGCTGCTTCGC
AACTTCTCTGATGATGAGAGTTTATTTTTTGGTGATACTGGGCTATCCAAA
AATCCAATCGAGCTTGTAGAAGGTTGGTTCAGTAGTTGGAAGCTCTAT
TGCTCTTTTTTTCTTTATCATAGGGTTAATCATTGGACTATTCTTGGTTC
TCCGAGTTGGTATCCATCTTTGCATTAAATTAAGCACACCAAGAAAAGA
CAGATTTATACAGACATAGAGATGAGAATTC

[0225] A 4-vector system (i.e., a 3-vector lentiviral packaging system) has also been designed and produced using the methods and materials described herein. A schematic of the 4-vector system is shown in FIG. 5. Briefly, and with reference to FIG. 5, the top-most vector is a helper plasmid, which, in this case, does not include Rev. The vector second from the top is a separate Rev plasmid. The vector second from the bottom is the envelope plasmid. The bottom-most vector is the previously described therapeutic vector.

[0226] Referring, in part, to FIG. 5, the Helper plasmid includes a CAG enhancer (SEQ ID NO: 49); a CAG promoter (SEQ ID NO: 50); a chicken beta actin intron (SEQ ID NO: 51); a HIV gag (SEQ ID NO: 52); a HIV Pol (SEQ ID NO: 53); a HIV Int (SEQ ID NO: 54); a HIV RRE (SEQ ID NO: 55); and a rabbit beta globin poly A (SEQ ID NO: 56).

[0227] The Rev plasmid includes a RSV promoter (SEQ ID NO: 57); a HIV Rev (SEQ ID NO: 58); and a rabbit beta globin poly A (SEQ ID NO: 59).

[0228] The Envelope plasmid includes a CMV promoter (SEQ ID NO: 60); a beta globin intron (SEQ ID NO: 61); a VSV-G (SEQ ID NO: 62); and a rabbit beta globin poly A (SEQ ID NO: 63).

[0229] Synthesis of a 3-Vector Lentiviral Packaging System Including Helper, Rev, and Envelope Plasmids.

[0230] Materials and Methods:

[0231] Construction of the Helper Plasmid without Rev:

[0232] The Helper plasmid without Rev was constructed by inserting a DNA fragment containing the RRE and rabbit beta globin poly A sequence. This sequence was synthesized by MWG Operon with flanking XbaI and XmaI restriction sites. The RRE/rabbit poly A beta globin sequence was then inserted into the Helper plasmid at the XbaI and XmaI restriction sites. The DNA sequence is as follows:

(SEQ ID NO: 87)
TCTAGAAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTA
TGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCT
GGTATAGTGCAGCAGCAGAACAATTGCTGAGGGCTATTGAGGCGCAACA
GCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGAA
TCTTGGCTGTGGAAGATACCTAAGGATCAACAGCTCCTAGATCTTTTT
CCCTCTGCCAAAAATTATGGGGACATCATGAAGCCCTTGAGCATCTGAC
TTCTGGCTAATAAAGGAAATTTATTTTTCATTGCAATAGTGTGTGGAATT
TTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAATCATTTAAAA
CATCAGAATGAGTATTTGGTTTAGAGTTTGGAACATATGCCATATGCTG
GCTGCCATGAACAAAGGTGGCTATAAAGAGGTCATCAGTATATGAACAG
CCCCCTGCTGCCATTCTCTTATCCATAGAAAAGCCTTGACTTGAGGTTA
GATTTTTTTTATATTTTGTGTTATTTTTTCTTTAACATCCCTAA
AATTTTCTTACATGTTTTACTAGCCAGATTTTTCTCTCTCTCTGACTA
CTCCAGTCATAGCTGTCCCTCTCTCTTATGAAGATCCCTCGACCTGCA
GCCCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAAATTGT
TATCCGCTCACAATTCACACAACATACGAGCCGGAAGCATAAAGTGTA
AGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCT
CACTGCCCGCTTTCCAGTCGGGAAACCTGTGTCGACGCGATCCGCATC
TCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCGCCCC
CTAACTCCGCCAGTTCGCCCATTTCTCGCCCATGAGTACTAATTTT
TTTTATTTATGCAGAGGCCGAGGCCCTCGGCCTCTGAGCTATTCCAGA
AGTAGTGAGGAGGCTTTTTTGGAGGCTAGGCTTTTGCAAAAGCTAACT
TGTTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAAT
TTCACAAATAAAGCATTTTTTCTACTGCATTCTAGTTGTGTTTGTCCAA
ACTCATCAATGTATCTTATCACCCTGGG

[0233] Construction of the Rev Plasmid:

[0234] The RSV promoter and HIV Rev sequence was synthesized as a single DNA fragment by MWG Operon with flanking MfeI and XbaI restriction sites. The DNA fragment was then inserted into the pCDNA3.1 plasmid (Invitrogen) at the MfeI and XbaI restriction sites in which the CMV promoter is replaced with the RSV promoter. The DNA sequence was as follows:

(SEQ ID NO: 88)

```

CAATTGCGATGTACGGGCCAGATATACGCGTATCTGAGGGGACTAGGGTG
TGTTTAGGCGAAAAGCGGGGCTTCGGTTGTACGCGGTTAGGAGTCCCCTC
AGGATATAGTAGTTTCGCTTTTGCATAGGGAGGGGAAATGTAGTCTTAT
GCAATACACTTGTAGTCTTGCAACATGGTAACGATGAGTTAGCAACATGC
CTTACAAGGAGAGAAAAAGCACCGTGCATGCCGATTGGTGAAGTAAGGT
GGTACGATCGTGCCTTATTAGGAAGGCAACAGACAGGTCTGACATGGATT
GGACGAACCACTGAATTCGCGATTGAGAGATAATTGTATTAAAGTGCCCT
AGCTCGATACAATAAACGCCATTTGACCATTACCCACATTGGTGTGCACC
TCCAAGCTCGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGAGCCCAT
CCACGCTGTTTTCGACTCCATAGAAGACACCGGGACCGATCCAGCCTCCC
CTCGAAGCTAGCGATTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAG
CGACGAAGAACTCCTCAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAA
GCAACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAAGGAATAGA
AGAAGAAGGTGGAGAGAGAGACAGAGACAGATCCATTGATTAGTGAACG
GATCCTTAGCACTTATCTGGGACGATCTGCGGAGCCTGTGCCTCTTCAGC
TACCACCGCTTGAGAGACTTACTCTTGATTGTAACGAGGATTGTGGAAC
TCTGGGACGCAGGGGGTGGGAAGCCCTCAAATATTGGTGAATCTCCTAC
AATATTGGAGTCAGGAGCTAAAGAATAGTCTAGA

```

[0235] The plasmids for the 2-vector and 3-vector packaging systems could be modified with similar elements and the intron sequences could potentially be removed without loss of vector function. For example, the following elements could replace similar elements in the 2-vector and 3-vector packaging system:

[0236] Promoters: Elongation Factor-1 (EF-1) (SEQ ID NO: 64), phosphoglycerate kinase (PGK) (SEQ ID NO: 65), and ubiquitin C (UbC) (SEQ ID NO: 66) can replace the CMV (SEQ ID NO: 60) or CAG promoter (SEQ ID NO: 100). These sequences can also be further varied by addition, substitution, deletion or mutation.

[0237] Poly A sequences: SV40 poly A (SEQ ID NO: 67) and bGH poly A (SEQ ID NO: 68) can replace the rabbit beta globin poly A (SEQ ID NO: 48). These sequences can also be further varied by addition, substitution, deletion or mutation.

[0238] HIV Gag, Pol, and Integrase sequences: The HIV sequences in the Helper plasmid can be constructed from different HIV strains or clades. For example, HIV Gag (SEQ ID NO: 69);

[0239] HIV Pol (SEQ ID NO: 70); and HIV Int (SEQ ID NO: 71) from the Bal strain can be interchanged with the gag, pol, and int sequences contained in the helper/helper plus Rev plasmids as outlined herein. These sequences can also be further varied by addition, substitution, deletion or mutation.

[0240] Envelope: The VSV-G glycoprotein can be substituted with membrane glycoproteins from feline endogenous virus (RD114) (SEQ ID NO: 72), gibbon ape leukemia virus (GALV) (SEQ ID NO: 73), Rabies (FUG) (SEQ ID NO: 74), lymphocytic choriomeningitis virus (LCMV) (SEQ ID NO: 75), influenza A fowl plague virus (FPV) (SEQ ID NO: 76), Ross River alphavirus (RRV) (SEQ ID NO: 77), murine

leukemia virus 10A1 (MLV) (SEQ ID NO: 78), or Ebola virus (EboV) (SEQ ID NO: 79). Sequences for these envelopes are identified in the sequence portion herein. Further, these sequences can also be further varied by addition, substitution, deletion or mutation.

[0241] In summary, the 3-vector versus 4-vector systems can be compared and contrasted, in part, as follows. The 3-vector lentiviral vector system contains: 1. Helper plasmid: HIV Gag, Pol, Integrase, and Rev/Tat; 2. Envelope plasmid: VSV-G/FUG envelope; and 3. Therapeutic vector: RSV 5'LTR, Psi Packaging Signal, Gag fragment, RRE, Env fragment, cPPT, WPRE, and 3'delta LTR. The 4-vector lentiviral vector system contains: 1. Helper plasmid: HIV Gag, Pol, and Integrase; 2. Rev plasmid: Rev; 3. Envelope plasmid: VSV-G/FUG envelope; and 4. Therapeutic vector: RSV 5'LTR, Psi Packaging Signal, Gag fragment, RRE, Env fragment, cPPT, WPRE, and 3'delta LTR. Sequences corresponding with the above elements are identified in the sequence listings portion herein.

Example 2: Development of an Anti-HIV Lentivirus Vector

[0242] The purpose of this example was to develop an anti-HIV lentivirus vector.

[0243] Inhibitory RNA Designs.

[0244] The sequence of *Homo sapiens* chemokine C-C motif receptor 5 (CCR5) (GC03P046377) mRNA was used to search for potential siRNA or shRNA candidates to knockdown CCR5 levels in human cells. Potential RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Broad Institute or the BLOCK-iT RNAi Designer from Thermo Scientific. Individual selected shRNA sequences were inserted into lentiviral vectors immediately 3' to a RNA polymerase III promoter such as H1, U6, or 7SK to regulate shRNA expression. These lentivirus-shRNA constructs were used to transduce cells and measure the change in specific mRNA levels. The shRNA most potent for reducing mRNA levels were embedded individually within a microRNA backbone to allow for expression by either the CMV or EF-1alpha RNA polymerase II promoters. The microRNA backbone was selected from mirbase.org. RNA sequences were also synthesized as synthetic siRNA oligonucleotides and introduced directly into cells without using a lentiviral vector.

[0245] The genomic sequence of Bal strain of human immunodeficiency virus type 1 (HIV-1 85US_BaL, accession number AY713409) was used to search for potential siRNA or shRNA candidates to knockdown HIV replication levels in human cells. Based on sequence homology and experience, the search focused on regions of the Tat and Vif genes of HIV although an individual of skill in the art will understand that use of these regions is non-limiting and other potential targets might be selected. Importantly, highly conserved regions of gag or pol genes could not be targeted by shRNA because these same sequences were present in the packaging system complementation plasmids needed for vector manufacturing. As with the CCR5 (NM 000579.3, NM 001100168.1-specific) RNAs, potential HIV-specific RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Gene-E Software Suite hosted by the Broad Institute (broadinstitute.org/mai/public) or the BLOCK-iT RNAi Designer from Thermo Scientific (madesigner.thermofisher.

com/maexpress/setOption.

do?designOption=shrna&pid=67126273 60706061801). Individual selected shRNA sequences were inserted into lentiviral vectors immediately 3' to a RNA polymerase III promoter such as H1, U6, or 7SK to regulate shRNA expression. These lentivirus-shRNA constructs were used to transduce cells and measure the change in specific mRNA levels. The shRNA most potent for reducing mRNA levels were embedded individually within a microRNA backbone to allow for expression by either the CMV or EF-1alpha RNA polymerase II promoters.

[0246] Vector Constructions.

[0247] For CCR5, Tat or Vif shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by Eurofins MWG Operon, LLC. Overlapping sense and antisense oligonucleotide sequences were mixed and annealed during cooling from 70 degrees Celsius to room temperature. The lentiviral vector was digested with the restriction enzymes BamHI and EcoRI for one hour at 37 degrees Celsius. The digested lentiviral vector was purified by agarose gel electrophoresis and extracted from the gel using a DNA gel extraction kit from Invitrogen. The DNA concentrations were determined and vector to oligo (3:1 ratio) were mixed, allowed to anneal, and ligated. The ligation reaction was performed with T4 DNA ligase for 30 minutes at room temperature. 2.5 microliters of the ligation mix were added to 25 microliters of STBL3 competent bacterial cells. Transformation was achieved after heat-shock at 42 degrees Celsius. Bacterial cells were spread on agar plates containing ampicillin and drug-resistant colonies (indicating the presence of ampicillin-resistance plasmids) were recovered, purified and expanded in LB broth. To check for insertion of the oligo sequences, plasmid DNA were extracted from harvested bacteria cultures with the Invitrogen DNA mini prep kit. Insertion of the shRNA sequence in the lentiviral vector was verified by DNA sequencing using a specific primer for the promoter used to regulate shRNA expression. Exemplary vector sequences that were determined to restrict HIV replication can be found in FIG. 7. For example, the shRNA sequences with the highest activity against CCR5, Tat or Vif gene expression were then assembled into a microRNA (miR) cluster under control of the EF-1 alpha promoter. The promoter and miR sequences are depicted in FIG. 7.

[0248] Further, and using standard molecular biology techniques (e.g., Sambrook; *Molecular Cloning: A Laboratory Manual*, 4th Ed.) as well as the techniques described herein, a series of lentiviral vectors have been developed as depicted in FIG. 8 herein.

[0249] Vector 1 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a H1 element (SEQ ID NO: 101); a shCCR5 (SEQ ID NOS: 16, 18, 20, 22, or 24-Y); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0250] Vector 2 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a H1 element (SEQ ID NO: 101); a shRev/Tat (SEQ ID NO: 10); a H1 element (SEQ ID NO: 101); a shCCR5 (SEQ ID NOS: 16, 18, 20, 22, or 24); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0251] Vector 3 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a H1 element (SEQ ID NO: 101); a shGag (SEQ ID NO: 12); a H1 element (SEQ ID NO: 101); a shCCR5 (SEQ ID NOS: 16, 18, 20, 22, or 24); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0252] Vector 4 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a 7SK element (SEQ ID NO: 103); a shRev/Tat (SEQ ID NO: 10); a H1 element (SEQ ID NO: 101); a shCCR5 (SEQ ID NOS: 16, 18, 20, 22, or 24); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0253] Vector 5 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a EF1 element (SEQ ID NO: 4); miR30CCR5 (SEQ ID NO: 1); MiR21Vif (SEQ ID NO: 2); miR185Tat (SEQ ID NO: 3); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0254] Vector 6 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a EF1 element (SEQ ID NO: 4); miR30CCR5 (SEQ ID NO: 1); MiR21Vif (SEQ ID NO: 2); miR155Tat (SEQ ID NO: 104); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0255] Vector 7 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a EF1 element (SEQ ID NO: 4); miR30CCR5 (SEQ ID NO: 1); MiR21Vif (SEQ ID NO: 2); miR185Tat (SEQ ID NO: 3); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

[0256] Vector 8 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a EF1 element (SEQ ID NO: 4); miR30CCR5 (SEQ ID NO: 1); MiR21Vif (SEQ ID NO: 2); miR185Tat (SEQ ID NO: 3); and a long terminal repeat portion (SEQ ID NO: 102).

[0257] Vector 9 was developed and contains, from left to right: a long terminal repeat (LTR) portion (SEQ ID NO: 35); a CD4 element (SEQ ID NO: 30); miR30CCR5 (SEQ ID NO: 1); miR21Vif (SEQ ID NO: 2); miR185Tat (SEQ ID NO: 3); a posttranscriptional regulatory element of woodchuck hepatitis virus (WPRE) (SEQ ID NOS: 32, 80); and a long terminal repeat portion (SEQ ID NO: 102).

Development of Vectors

[0258] It should be noted that not all vectors developed for these experiments necessarily worked as might be predicted. More specifically, a lentivirus vector against HIV might include three main components: 1) inhibitory RNA to reduce the level of HIV binding proteins (receptors) on the target cell surface to block initial virus attachment and penetration; 2) overexpression of the HIV TAR sequence that will sequester viral Tat protein and decrease its ability to transactivate viral gene expression; and 3) inhibitory RNA that attack important and conserved sequences within the HIV genome.

[0259] With respect to the first point above, a key cell surface HIV binding protein is the chemokine receptor CCR5. HIV particles attach to susceptible T cells by binding to the CD4 and CCR5 cell surface proteins. Because CD4 is an essential glycoprotein on the cell surface that is important for the immunological function of T cells, this was not chosen as a target to manipulate its expression levels. However, people born homozygous for null mutations in the CCR5 gene and completely lacking receptor expression, live normal lives save for enhanced susceptibility to a few infectious diseases and the possibility of developing rare autoimmunity. Thus, modulating CCR5 was determined to be a relatively safe approach and was a primary target in the development of anti-HIV lentivirus vectors.

[0260] With respect to the second point above, the viral TAR sequence is a highly structured region of HIV genomic RNA that binds tightly to viral Tat protein. The Tat:TAR complex is important for efficient generation of viral RNA. Over-expression of the TAR region was envisioned as a decoy molecule that would sequester Tat protein and decrease the levels of viral RNA. However, TAR proved toxic to most mammalian cells including cells used for manufacturing lentivirus particles. Further, TAR was inefficient for inhibiting viral gene expression in other laboratories and has been discarded as a viable component in HIV gene therapy.

[0261] In various embodiments, viral gene sequences have been identified that meet 3 criteria: i) Sequences that are reasonably conserved across a range of HIV isolates representative of the epidemic in a geographic region of interest; ii) reduction in RNA levels due to the activity of an inhibitory RNA in a viral vector will reduce the corresponding protein levels by an amount sufficient to meaningfully reduce HIV replication; and iii) the viral gene sequence(s) targeted by inhibitory RNA are not present in the genes required for packaging and assembling viral vector particles during manufacturing. In various embodiments, a sequence at the junction of HIV Tat and Rev genes and a second sequence within the HIV Vif gene have been targeted by inhibitory RNA. The Tat/Rev targeting has an additional benefit of reducing HIV envelope glycoprotein expression because this region overlaps with the envelope gene in the HIV genome.

[0262] Various methods for vector development and testing relies first on identifying suitable targets (as described herein) followed by constructing plasmid DNAs expressing individual or multiple inhibitory RNA species for testing in

cell models, and finally constructing lentivirus vectors containing inhibitory RNA with proven anti-HIV function. The lentivirus vectors are tested for toxicity, yield during in vitro production, and effectiveness against HIV in terms of reducing CCR5 expression levels or lowering viral gene products to inhibit virus replication.

[0263] Table 2 below demonstrates progression through multiple versions of inhibitory constructs until arriving at a clinical candidate. Initially, shRNA (short homology RNA) molecules were designed and expressed from plasmid DNA constructs.

[0264] Plasmids 1-4, as detailed in Table 2 below, tested shRNA sequences against Gag, Pol and RT genes of HIV. While each shRNA was active for suppressing viral protein expression in a cell model, there were two important problems that prevented further development. First, the sequences were targeted to a laboratory isolate of HIV that was not representative of Clade B HIV strains currently circulating in North America and Europe. Second, these shRNA targeted critical components in the lentivirus vector packaging system and would severely reduce vector yield during manufacturing. Plasmid 5, as detailed in Table 2, was selected to target CCR5 and provided a lead candidate sequence. Plasmids 6, 7, 8, 9, 10, and 11, as detailed in Table 2, incorporated the TAR sequence and it was found they produced unacceptable toxicity for mammalian cells including cells used for lentivirus vector manufacturing. Plasmid 2, as detailed in Table 2, identified a lead shRNA sequence capable of reducing Tat RNA expression. Plasmid 12, as detailed in Table 2, demonstrated the effectiveness of shCCR5 expressed as a microRNA (miR) in a lentiviral vector and confirmed it should be in the final product. Plasmid 13, as detailed in Table 2, demonstrated the effectiveness of a shVif expressed as a microRNA (miR) in a lentiviral vector and confirmed it should be in the final product. Plasmid 14, as detailed in Table 2, demonstrated the effectiveness of shTat expressed as a microRNA (miR) in a lentiviral vector and confirmed it should be in the final product. Plasmid 15, as detailed in Table 2, contained the miR CCR5, miR Tat and miR Vif in the form of a miR cluster expressed from a single promoter. These miR do not target critical components in the lentivirus vector packaging system and proved to have negligible toxicity for mammalian cells. The miRs within the cluster were equally effective to individual miR that were tested previously, and the overall impact was a substantial reduction in replication of a CCR5-tropic HIV BaL strain.

TABLE 2

Development of HIV Vectors					
Internal Code	Material	Description	Remarks	Decision	
1 SIH-H1-shRT-1,3	Lentiviral vector	shRNA construct for RT of LAI strain	Wrong target, lab virus, no virus test	Abandon	
2 SIH-H1-shRT43 (Tat/Rev NL4-3)	Lentiviral vector	H1 promoter shRNA Tat/Rev overlap	Tat protein knock-down >90%	Lead	

TABLE 2-continued

Development of HIV Vectors				
Internal Code	Material	Description	Remarks	Decision
<p>Vector Construction: For Rev/Tat (RT) shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by MWG Operon. Two different Rev/Tat target sequences were tested for their ability to decrease Tat mRNA expression. The RT1,3 target sequence is (5'-ATGGCAGGAAGAAGCGGAG-3') (SEQ ID NO: 89) and shRNA sequence is (5'-ATGGCAGGAAGAAGCGGAGTTCAAGAGACTCCGCTTCTCCTGCCATTTTT-3') (SEQ ID NO: 90). The RT43 sequence is (5'-GCGGAGACGCGACGAAGAGC-3') (SEQ ID NO: 9) and shRNA sequence is (5'-GCGGAGACGCGACGAAGAGCTTCAAGAGAGCTCTTCGTCGCTGTCTCCGCTTTTT-3') (SEQ ID NO: 10). Oligonucleotide sequences were inserted into the pSIH lentiviral vector (System Biosciences).</p> <p>Functional test for shRNA against Rev/Tat: The ability of the vector to reduce Tat expression was tested using a luciferase reporter plasmid which contained the Rev/Tat target sequences inserted into the 3'-UTR (untranslated region of the mRNA). Either the shRT1,3 or shRT43 plasmid was co-transfected with the plasmid containing luciferase and the Rev/Tat target sequence. There was a 90% reduction in light emission indicating strong function of the shRT43 shRNA sequence but less than 10% with the shRT1,3 plasmid.</p> <p>Conclusion: The SIH-H1-shRT43 was superior to SIH-H1-shRT-1,3 in terms of reducing mRNA levels in the Luciferase assay system. This indicates potent inhibitory activity of the shRT43 sequence and it was selected as a lead candidate for further development.</p>				
3 SIH-H1-shGag-1	Lentiviral vector	shRNA construct for LAI	Inhibits Gag expression but will inhibit packaging	Abandon
<p>Vector Construction: For Gag shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by MWG Operon. A Gag target sequence was tested for their ability to decrease Gag mRNA expression. The Gag target sequence is (5'-GAAGAAATGATGACAGCAT-3') (SEQ ID NO: 11) and shRNA sequence is (5'-GAAGAAATGATGACAGCATTCAAGAGAATGCTGTCATCATTTCTTCTTTT-3') (SEQ ID NO: 12). Oligonucleotide sequences were inserted into the pSIH lentiviral vector (System Biosciences).</p> <p>Functional test for shRNA against Gag: The ability of the vector to reduce Gag expression was tested using a luciferase reporter plasmid which contained the Gag target sequences inserted into the 3'-UTR (untranslated region of the mRNA). The Gag plasmid was co-transfected with the plasmid containing luciferase and the Gag target sequence. There was nearly a 90% reduction in light emission indicating a strong effect of the shGag shRNA sequence.</p> <p>Conclusion: This shRNA sequence is potent against HIV Gag expression but was abandoned. The lentivirus packaging system requires production of Gag from the helper plasmid and shRNA inhibition of Gag will reduce lentivirus vector yield. This shRNA sequence could be used as an oligonucleotide inhibitor of HIV or incorporated into an alternate viral vector packaging system that uses a different vector genome or is modified to resist inhibition by this shRNA.</p>				
4 SIH-H1-shPol-1	Lentiviral vector	shRNA construct for Pol	Inhibits Pol expression but will inhibit packaging	Abandon
<p>Vector Construction: A Pol shRNA was constructed with oligonucleotide sequences containing BamHI and EcoRI restriction sites that were synthesized by MWG Operon. A Pol target sequence was tested for its ability to decrease Pol mRNA expression. The Pol target sequence is (5'-CAGGAGCAGATGATACAG-3') (SEQ ID NO: 13) and shRNA sequence is (5'-CAGGAGATGATACAGTTCAAGAGACTGTATCATCTGCTCCTGTTTT-3') (SEQ ID NO: 14). Oligonucleotide sequences were inserted into the pSIH lentiviral vector (System Biosciences).</p> <p>Functional tests for shRNA against HIV Pol: The ability of the vector to reduce Pol expression was tested using a luciferase reporter plasmid which contained the Pol target sequences inserted into the 3'-UTR (untranslated region of the mRNA). The Pol plasmid was co-transfected with the plasmid containing luciferase and the Pol target sequence. There was a 60% reduction in light emission indicating a strong effect of the shPol shRNA sequence.</p> <p>Conclusion: This shRNA sequence is potent against HIV Pol expression but was abandoned. The lentivirus packaging system requires production of Pol from the helper plasmid and shRNA inhibition of Pol will reduce lentivirus vector yield. This shRNA sequence could be used as an oligonucleotide inhibitor of HIV or incorporated into an alternate viral vector packaging system that uses a different vector genome or is modified to resist inhibition by this shRNA.</p>				

TABLE 2-continued

Development of HIV Vectors				
Internal Code	Material	Description	Remarks	Decision
5 SIH-H1-shCCR5-1	Lentiviral vector	shRNA construct for CCR5	Best of 5 candidates, Extracellular CCR5 protein reduction >90%	Lead
<p>Vector Construction: A CCR5 shRNA was constructed with oligonucleotide sequences containing BamHI and EcoRI restriction sites that were synthesized by MWG Operon. Oligonucleotide sequences were inserted into the pSIH lentiviral vector (System Biosciences). The CCR5 target sequence #1, which focuses on CCR5 gene sequence 1 (SEQ ID NO: 25), is (5'-GTGTCAAGTCCAATCTATG-3') (SEQ ID NO: 15) and the shRNA sequence is (5'-GTGTCAAGTCCAATCTATGTTCAAGAGACATAGATTGGACTTGACACTTTT-3') (SEQ ID NO: 16). The CCR5 target sequence #2, which focuses on CCR5 gene sequence 2 (SEQ ID NO: 26), is (5'-GAGCATGACTGACATCTAC-3') (SEQ ID NO: 17) and the shRNA sequence is (5'-GAGCATGACTGACATCTACTTCAAGAGAGTAGATGTCAGTCATGCTCTTTT-3') (SEQ ID NO: 18). The CCR5 target sequence #3, which focuses on CCR5 gene sequence 3 (SEQ ID NO: 27), is (5'-GTAGCTCTAACAGGTTGGA-3') (SEQ ID NO: 19) and the shRNA sequence is (5'-GTAGCTCTAACAGGTTGATTCAAGAGATCCAACCTGTTAGAGCTACTTTT-3') (SEQ ID NO: 20). The CCR5 target sequence #4, which focuses on CCR5 gene sequence 4 (SEQ ID NO: 28), is (5'-GTTTCAAGAACTACCTCTTA-3') (SEQ ID NO: 21) and the shRNA sequence is (5'-GTTTCAAGAACTACCTCTTATTCAAGAGATAAGAGGTAGTTTCTGAACCTTTT-3') (SEQ ID NO: 22). The CCR5 target sequence #5, which focuses on CCR5 gene sequence 5 (SEQ ID NO: 29), is (5'-GAGCAAGCTCAGTTTACACC-3') (SEQ ID NO: 23) and the shRNA sequence is (5'-GAGCAAGCTCAGTTTACACCTTCAAGAGAGGTGTAAGTCTGCTCTTTT-3') (SEQ ID NO: 24).</p> <p>Functional test for shRNA against CCR5: The ability of a CCR5 shRNA sequence to knock-down CCR5 RNA expression was initially tested by co-transfecting each of the lentiviral plasmids, in separate experiments for each plasmid, containing one of the five CCR5 target sequences with a plasmid expressing the human CCR5 gene. CCR5 mRNA expression was then assessed by qPCR analysis using CCR5-specific primers.</p> <p>Conclusion: Based on the reduction in CCR5 mRNA levels the shRNCCR5-1 was most potent for reducing CCR5 gene expression. This shRNA was selected as a lead candidate.</p>				
6 SIH-U6-TAR	Lentiviral vector	U6 promoter-TAR	Toxic to cells	Abandon
7 SIH-U6-TAR-H1-shCCR5	Lentiviral vector	U6 promoter-TAR-H1-shCCR5	Toxic to cells	Abandon
8 U6-TAR-H1-shRT	Lentiviral vector	U6 promoter-TAR-H1-RT	Suppress HIV, toxic to cells, poor packaging	Abandon
9 U6-TAR-7SK-shRT	Lentiviral vector	Change shRNA promoter to 7SK	Toxic, poor packaging	Abandon
10 U6-TAR-H1-shRT-H1-shCCR5	Lentiviral vector	U6 promoter-TAR-H1-RT-H1-shCCR5	Toxic, poor packaging, H1 repeats	Abandon
11 U6-TAR-7SK-shRT-H1-CCR5	Lentiviral vector	Change shRNA promoter to 7SK	Toxic, poor packaging	Abandon

Vector Construction: A TAR decoy sequence containing flanking KpnI restriction sites was synthesized by MWG operon and inserted into the pSIH lentiviral vector (System Biosciences) at the KpnI site. In this vector, TAR expression is regulated by the U6 promoter. The TAR decoy sequence is (5'-CTTGCAATGATGTCGTAATTTGCGTCTTACCTCGTTCTCGACAGCGACCATCTGAGCCTGGGAGCTCTCTGGCTGTCTAGTAAGCTGGTACAGAAGTTGACGAAATTTTACTGAGCAAGAAA-3') (SEQ ID NO: 8). Expression of the TAR decoy sequence was determined by qPCR analysis using specific primers for the TAR sequence. Additional vectors were constructed also containing the TAR sequence. The H1 promoter and shRT sequence was inserted in this vector in the XhoI site. The H1 shRT sequence is (5'-

TABLE 2-continued

Development of HIV Vectors				
Internal Code	Material	Description	Remarks	Decision
<p>GAACGCTGACGTCAATCAACCCGCTCCAAGGAATCGCGGGCCAGTGTCACTAGGC GGGAACACCCAGCGCGCTGCGCCCTGGCAGGAAGATGGCTGTGAGGGACAGGG GAGTGGCGCCCTGCAATATTTGCATGTGCTATGTGTTCTGGGAATCACCATAAA CGTGAAATGTCTTTGGATTGGGAATCTTATAAGTTCTGTATGAGACCACTGGAT CCGCGGAGACAGCGACGAAGAGCTTCAAGAGAGCTCTTCGTCGCTGTCTCCGCTTT TT-3') (SEQ ID NO: 91). This vector could express TAR and knockdown RT. The 7SK promoter was also substituted for the H1 promoter to regulate shRT expression. Another vector was constructed containing U6 TAR, H1 shRT, and H1 shCCR5. The H1 shCCR5 sequence was inserted into the SpeI site of the plasmid containing U6 TAR and H1 shRT. The H1 CCR5 sequence is (5'- GAACGCTGACGTCAATCAACCCGCTCCAAGGAATCGCGGGCCAGTGTCACTAGGC GGGAACACCCAGCGCGCTGCGCCCTGGCAGGAAGATGGCTGTGAGGGACAGGG GAGTGGCGCCCTGCAATATTTGCATGTGCTATGTGTTCTGGGAATCACCATAAA CGTGAAATGTCTTTGGATTGGGAATCTTATAAGTTCTGTATGAGACCACTGGAT CCGTGTCAGTCAATCTATGTTCAGAGACATAGATTGGACTTGACACTTTT-3') (SEQ ID NO: 92). The 7SK promoter was also substituted for the H1 promoter to regulate shRT expression.</p>				
<p>Functional test for TAR decoy activity: We tested the effect of SIH-U6-TAR on packaging efficiency. When TAR sequence was included, the yield of vector in the SIH packaging system was reduced substantially.</p>				
<p>Conclusion: Lentivirus vectors expressing the TAR decoy sequence are unsuitable for commercial development due to low vector yields. These constructs were abandoned.</p>				
12 shCCR5	Lentiviral vector	microRNA sequence	Extracellular CCR5 protein reduction >90%	Lead

Vector Construction: A CCR5 microRNA was constructed with oligonucleotide sequences containing BsrGI and NotI restriction sites that were synthesized by MWG Operon. Oligonucleotide sequences were inserted into the pCDH lentiviral vector (System Biosciences). The EF-1 promoter was substituted for a CMV promoter that was used in the plasmid construct Test Material 5. The EF-1 promoter was synthesized by MWG Operon containing flanking ClaI and BsrGI restriction sites and inserted into the pCDH vector containing shCCR5-1. The EF-1 promoter sequence is (5'-
CCGGTGCCTAGAGAAGGTGGCGCGGGTAACTGGGAAAGTGATGTCGTGACTG
GCTCCGCTTTTCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCC
GTGAACGTTCTTTTCGCAACGGGTTTCCCGCCAGAACACAGGTAAAGTCCCGTGTG
TGTTTCCCGCGGCTGGCCCTTTACGGGTTATGGCCCTTGCCTGCTGCTTGAATTA
CTTCCACGCCCTGGCTGCAGTACGTGATTCTTGATCCCGAGCTTCGGGTTGGAAG
TGGGTGGGAGAGTTCGAGGCCTTGCCTTAAGGAGCCCTTCGCTCGTGTGTTGAG
TTGAGGCTGGCCTGGGCGCTGGGCGCGCCGCGTGCGAATCTGGTGGCACCTTCG
CGCTGTCTCGCTGCTTTTCGATAAGTCTCTAGCCATTTAAATTTTGTATGACCTGC
TGCGACGCTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAGATCTGCACA
CTGGTATTTCCGTTTGGGGCCGCGGCGCGCAGCGGGCCGCTGCGTCCAGCGC
ACATGTTCCGGGAGGCGGGCTGCGAGCGCGGCCACCGAGAATCGGACGGGG
TAGTCTCAAGCTGGCCGCTGCTCTGGTGCCTGGCCTCGCGCCGCTGTATCGC
CCCGCCCTGGGCGGCAAGGCTGGCCCGGTGCGCACCAAGTTGCGTGAGCGGAAAGA
TGGCGCTTCCCGCCCTGCTGCGAGGAGCTCAAAATGGAGGACGCGCGCTCGG
GAGAGCGGGCGGGTGAATCACCCACAAAGGAAAGGGCTTTCCGCTCTCAGC
CGTCGCTTCATGTGACTCCACGAGTACCGGCGCGCTCCAGGCACCTCGATTAGT
TCTCGAGCTTTGGAGTACGTCGCTTTTAGGTTGGGGGAGGGGTTTATGCGATG
GAGTTTCCCCACACTGAGTGGGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGA
TGTAATTCCTCTGGAATTTGCCCTTTTGGAGTTGGATCTTGGTTCACTCTCAAGC
CTCAGACAGTGGTTCAAAGTTTTTCTTCCATTACAGGTGTCGTGA-3') (SEQ ID
NO: 4).

Functional test for lentivirus CDH-shCCR5-1: The ability of the miR CCR5 sequences to knock-down CCR5 expression was determined by transducing CEM-CCR5 T cells and measuring cell surface CCR5 expression after staining with a fluorescently-labeled monoclonal antibody against CCR5 and measuring the intensity of staining, that is directly proportional to the number of cell surface CCR5 molecules, by analytical flow cytometry. The most effective shRNA sequence for targeting CCR5 was CCR5 shRNA sequence #1. However, the most effective CCR5 targeting sequence for constructing the synthetic microRNA sequence was overlapping with CCR5 sequence #5; this conclusion was based on sequence alignments and experience with miRNA construction. Finally, the miR30 hairpin sequence was used to construct the synthetic miR30 CCR5 sequence which is (5'-
AGGTATATGTGTTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAG
CCACAGATGGGTAGAGCAAGCACAGTTTACCGCTGCCCTACTGCCCTCGGACTTCAA
GGGGCTT-3') (SEQ ID NO: 1). The miR CCR5 target sequence is (5'-
GAGCAAGCTCAGTTTACA-3') (SEQ ID NO: 5). At multiplicity of infection equal to 5,

TABLE 2-continued

Development of HIV Vectors				
Internal Code	Material	Description	Remarks	Decision
<p>generating on average 1.25 genome copies of integrated lentivirus per cell, CCR5 expression levels were reduced by $\geq 90\%$ indicating potent inhibition of CCR5 mRNA by the miR30CCR5 micro RNA construct in a lentivirus vector.</p> <p>Conclusion: The miR30CCR5 construct is potent for reducing CCR5 cell surface expression and is a lead candidate for a therapeutic lentivirus for HIV.</p>				
13 shVif	Lentiviral vector	microRNA sequence	Vif protein reduction $>80\%$	Lead
<p>Vector Construction: A Vif microRNA was constructed with oligonucleotide sequences containing BsrGI and NotI restriction sites that were synthesized by MWG Operon. Oligonucleotide sequences were inserted into the pCDH lentiviral vector (System Biosciences) containing an EF-1 promoter. Based on sequence alignments and experience with constructing synthetic miRNA, the miR21 hairpin sequence was used to construct the synthetic miR21 Vif sequence which is (5'-CATCTCCATGGCTGTACCACCTTGTGCGGGGATGTGTACTTCTGAACCTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGGTATCTTTCATCTGACCA-3') (SEQ ID NO: 2). The miR Vif target sequence is (5'-GGGATGTGTACTTCTGAACCT-3') (SEQ ID NO: 6).</p> <p>Functional test for potency of miR21Vif: The ability of the miR Vif sequence to knock-down Vif expression was determined by measuring Vif protein expression by immunoblot analysis using an anti-Vif monoclonal antibody to identify the Vif protein.</p> <p>Conclusion: the miR21Vif reduced Vif protein expression by ≥ 10-fold as determined by quantitative image analysis of immunoblot data. This was sufficient to justify miR21Vif as a lead candidate for our therapeutic lentivirus.</p>				
14 shTat	Lentiviral vector	microRNA sequence	Tat RNA reduction $>80\%$	Lead
<p>Vector Construction: A Tat microRNA was constructed with oligonucleotide sequences containing BsrGI and NotI restriction sites that were synthesized by MWG Operon. The microRNA cluster was inserted into the pCDH lentiviral vector (System Biosciences) containing an EF-1 promoter. Based on sequence alignments and experience in the construction of synthetic miRNA, the miR185 hairpin sequence was selected for constructing a synthetic miR185 Tat sequence which is (5'-GGGCTCGGCTCGAGCAGGGGCGAGGGATTCCGCTTCTTCTGCCATAGCGTGGTCCCCCTCCCTATGGCAGGCAGAGCGGCACCTTCCCTCCCAATGACCGCGCTTTCGTCG-3') (SEQ ID NO: 7).</p> <p>Functional test for potency of miR185Tat: The ability of miR Tat to knock-down Tat expression was determined by measuring Tat mRNA expression by RT-PCR analysis using Tat specific primers. We compared the miR185Tat with a similar miR155Tat on the basis of reducing the relative levels of Tat mRNA.</p> <p>Conclusion: The miR185Tat was approximately twice as potent for reducing Tat mRNA compare to miR155Tat, and was selected as the lead candidate for our therapeutic lentivirus.</p>				
15 shCCR5-shVif-shTat	Lentiviral vector	microRNA cluster sequence	CCR5 reduction $>90\%$, Vif protein reduction $>80\%$, RNA reduction $>80\%$, $>95\%$ inhibition of HIV replication	Candidate Tat
<p>Vector Construction: A miR30CCR5 miR21Vif miR185Tat microRNA cluster sequence was constructed with a synthetic DNA fragment containing BsrGI and NotI restriction sites that was synthesized by MWG Operon. The DNA fragment was inserted into the pCDH lentiviral vector (System Biosciences) containing the EF-1 promoter. The miR cluster sequence is (5'-AGGTATATTGCTGTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAGCCACAGATGGGTAGAGCAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAAAGGGGCTTCCCGGCATCTCCATGGCTGTACCACCTTGTGCGGGGATGTGTACTTCTGAACCTGTGTTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGGTATCTTTCATCTGACCACTAGCGGCTGGCTCGAGCAGGGGCGAGGGATTCCGCTTCTTCTGCCATAGCGTGGTCCCTCCCTATGGCAGGCAGAGCGGCACCTTCCCTCCCAATGACCGCGCTTTCGTC-3') (SEQ ID NO: 31) and incorporates Test Material</p>				

TABLE 2-continued

Development of HIV Vectors				
Internal Code	Material	Description	Remarks	Decision
12, Test Material 13 and Test Material 14 into a single cluster that can be expressed under control of the EF-1 promoter.				
Functional test for potency of the Lentivirus Vector AGT103 containing the microRNA cluster of miR30CCR5, miR21Vif and miR185Tat: The AGT103 vector was tested for potency against CCR5 using the assay for reduction in cell surface CCR5 expression (Test Material 12). The AGT103 vector was tested for potency against Vif using the assay for reduction in cell surface Vif expression (Test Material 13). The AGT103 vector was tested for potency against Tat using the assay for reduction in cell surface Tat expression (Test Material 14).				
Conclusion: Potency for reducing CCR5 expression by the miRNA cluster was similar to potency observed for the miR30CCR5 alone. Potency for reducing Vif expression by the miRNA cluster was similar to potency observed for the miR21Vif alone. Potency for reducing Tat expression by the miRNA cluster was similar to potency observed for the miR185Tat alone. The miRNA cluster is potent for reducing cell surface CCR5 levels and for inhibiting two HIV genes. Thus, AGT103 containing this miRNA cluster was selected as the therapeutic vector construct for our HIV functional cure program.				

[0265] Functional Assays.

[0266] Individual lentivirus vectors containing CCR5, Tat or Vif shRNA sequences and, for experimental purposes, expressing green fluorescent protein (GFP) under control of the CMV Immediate Early Promoter, and designated AGT103/CMV-GFP were tested for their ability to knock-down CCR5, Tat or Vif expression. Mammalian cells were transduced with lentiviral particles either in the presence or absence of polybrene. Cells were collected after 2-4 days; protein and RNA were analyzed for CCR5, Tat or Vif expression. Protein levels were tested by Western blot assay or by labeling cells with specific fluorescent antibodies (CCR5 assay), followed by analytical flow cytometry comparing modified and unmodified cell fluorescence using either the CCR5-specific or isotype control antibodies.

[0267] Starting Testing of Lentivirus.

[0268] T cell culture medium was made using RPMI 1640 supplemented with 10% FBS and 1% penicillin—streptomycin. Cytokine stocks of IL-2 10,000 units/ml, IL-12 1 µg/ml, IL-7 1 µg/ml, IL-15 1 µg/ml were also prepared in advance.

[0269] Prior to transduction with the lentivirus, an infectious viral titer was determined and used to calculate the amount of virus to add for the proper multiplicity of infection (MOI).

[0270] Day 0-12: Antigen-Specific Enrichment.

[0271] On day 0, cryopreserved PBMC were thawed, washed with 10 ml 37° C. medium at 1200 rpm for 10 minutes and resuspended at a concentration of 2×10⁶/ml in 37° C. medium. The cells were cultured at 0.5 ml/well in a 24-well plate at 37° C in 5% CO₂. To define the optimal stimulation conditions, cells were stimulated with combinations of reagents as listed in Table 3 below:

[0272] Final concentrations: IL-2=20 units/ml, IL-12=10 ng/ml, IL-7=10 ng/ml, IL-15=10 ng/ml, peptides=5 µg/ml individual peptide, MVA MOI=1.

[0273] On days 4 and 8, 0.5 ml fresh medium and cytokine at listed concentrations (all concentrations indicate the final concentration in the culture) were added to the stimulated cells.

[0274] Day 12-24: Non-Specific Expansion and Lentivirus Transduction.

[0275] On day 12, the stimulated cells were removed from the plate by pipetting and resuspended in fresh T cell culture medium at a concentration of 1×10⁶/ml. The resuspended cells were transferred to T25 culture flasks and stimulated with DYNABEADS® Human T-Activator CD3/CD28 following the manufacturer's instruction plus cytokine as listed above; flasks were incubated in the vertical position.

[0276] On day 14, AGT103/CMV-GFP was added at MOI 20 and cultures were returned to the incubator for 2 days. At this time, cells were recovered by pipetting, collected by centrifugation at 1300 rpm for 10 minutes, resuspended in the same volume of fresh medium, and centrifuged again to form a loose cell pellet. That cell pellet was resuspended in fresh medium with the same cytokines used in previous steps, with cells at 0.5×10⁶ viable cells per ml.

[0277] From days 14 to 23, the number of the cells was evaluated every 2 days and the cells were diluted to 0.5×10⁶/ml with fresh media. Cytokines were added every time.

[0278] On day 24, the cells were collected and the beads were removed from the cells. To remove the beads, cells were transferred to a suitable tube that was placed in the sorting magnet for 2 minutes. Supernatant containing the cells was transferred to a new tube. Cells were then cultured for 1 day in fresh medium at 1×10⁶/ml. Assays were

TABLE 3

1	2	3	4	5	6
IL-2 + IL-12	IL-7 + IL-15	Peptides + IL-2 + IL-12	Peptides + IL-7 + IL-15	MVA + IL- 2 + IL-12	MVA + IL- 7 + IL-15

performed to determine the frequencies of antigen-specific T cells and lentivirus transduced cells.

[0279] To prevent possible viral outgrowth, amprenavir (0.5 ng/ml) was added to the cultures on the first day of stimulation and every other day during the culture.

[0280] Examine Antigen-Specific T Cells by Intracellular Cytokine Staining for IFN-Gamma.

[0281] Cultured cells after peptide stimulation or after lentivirus transduction at 1×10^6 cells/ml were stimulated with medium alone (negative control), Gag peptides (5 μ g/ml individual peptide), or PHA (5 μ g/ml, positive control). After 4 hours, BD GolgiPlug™ (1:1000, BD Biosciences) was added to block Golgi transport. After 8 hours, cells were washed and stained with extracellular (CD3, CD4 or CD8; BD Biosciences) and intracellular (IFN-gamma; BD Biosciences) antibodies with BD Cytotfix/Cytoperm™ kit following the manufacturer's instruction. Samples were analyzed on a BD FACSCalibur™ Flow Cytometer. Control samples labeled with appropriate isotype-matched antibodies were included in each experiment. Data were analyzed using Flowjo software.

[0282] Lentivirus transduction rate was determined by the frequency of GFP+ cells. The transduced antigen-specific T cells are determined by the frequency of CD3+CD4+GFP+ IFN gamma+ cells; tests for CD3+CD8+GFP+IFN gamma+ cells are included as a control.

[0283] These results indicate that CD4 T cells, the target T cell population, can be transduced with lentiviruses that are designed to specifically knock down the expression of HIV-specific proteins, thus producing an expandable population of T cells that are immune to the virus. This example serves as a proof of concept indicating that the disclosed lentiviral constructs can be used in combination with vaccination to produce a functional cure in HIV patients.

Example 4: CCR5 Knockdown with Experimental Vectors

[0284] AGTc120 is a HeLa cell line that stably expresses large amounts of CD4 and CCR5. AGTc120 was transduced with or without LV-CMV-mCherry (the red fluorescent protein mCherry expressed under control of the CMV Immediate Early Promoter) or AGT103/CMV-mCherry. Gene expression of the mCherry fluorescent protein was controlled by a CMV (cytomegalovirus immediate early promoter) expression cassette. The LV-CMV-mCherry vector lacked a microRNA cluster, while AGT103/CMV-mCherry expressed therapeutic miRNA against CCR5, Vif, and Tat.

[0285] As shown in FIG. 9A, transduction efficiency was >90%. After 7 days, cells were collected and stained with fluorescent monoclonal antibody against CCR5 and subjected to analytical flow cytometry. Isotype controls are shown in gray on these histograms plotting Mean Fluorescence Intensity of CCR5 APC (x axis) versus cell number normalized to mode (y axis). After staining for cell surface CCR5, cells treated with no lentivirus or control lentivirus (expressing only the mCherry marker) showed no changes in CCR5 density while AGT103 (right section) reduced CCR5 staining intensity to nearly the levels of isotype control. After 7 days, cells were infected with or without R5-tropic HIV reporter virus Bal-GFP. 3 days later, cells were collected and analyzed by flow cytometry. More than 90% of cells were transduced. AGT103-CMV/CMVmCherry reduced CCR5 expression in transduced AGTc120 cells and blocked R5-tropic HIV infection compared with cells treated with the Control vector.

[0286] FIG. 9B shows the relative insensitivity of transduced AGTc120 cells to infection with HIV. As above, the lentivirus vectors express mCherry protein and a transduced cell that was also infected with HIV (expressing GFP) would appear as a double positive cell in the upper right quadrant of the false color flow cytometry dot plots. In the absence of HIV (upper panels), there were no GFP+ cells under any condition. After HIV infection (lower panels), 56% of cells were infected in the absence of lentivirus transduction and 53.6% of cells became infected in AGTc120 cells transduced with the LV-CMV-mCherry. When cells were transduced with the therapeutic AGT103/CMV-mCherry vector, only 0.83% of cells appeared in the double positive quadrant indicating they were transduced and infected.

[0287] Dividing 53.62 (proportion of double positive cells with control vector) by 0.83 (the proportion of double positive cells with the therapeutic vector) shows that AGT103 provided greater than 65-fold protection against HIV in this experimental system.

Example 5: Regulation of CCR5 Expression by shRNA Inhibitor Sequences in a Lentiviral Vector

[0288] Inhibitory RNA Design.

[0289] The sequence of *Homo sapiens* chemokine receptor CCR5 (CCR5, NC 000003.12) was used to search for potential siRNA or shRNA candidates to knockdown CCR5 levels in human cells. Potential RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Broad Institute or the BLOCK-IT RNA iDesigner from Thermo Scientific. A shRNA sequence may be inserted into a plasmid immediately after a RNA polymerase III promoter such as H1, U6, or 7SK to regulate shRNA expression. The shRNA sequence may also be inserted into a lentiviral vector using similar promoters or embedded within a microRNA backbone to allow for expression by an RNA polymerase II promoter such as CMV or EF-1 alpha. The RNA sequence may also be synthesized as a siRNA oligonucleotide and utilized independently of a plasmid or lentiviral vector.

[0290] Plasmid Construction.

[0291] For CCR5 shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by MWG Operon. Oligonucleotide sequences were annealed by incubating at 70° C. then cooled to room temperature. Annealed oligonucleotides were digested with the restriction enzymes BamHI and EcoRI for one hour at 37° C., then the enzymes were inactivated at 70° C. for 20 minutes. In parallel, plasmid DNA was digested with the restriction enzymes BamHI and EcoRI for one hour at 37° C. The digested plasmid DNA was purified by agarose gel electrophoresis and extracted from the gel using a DNA gel extraction kit from Invitrogen. The DNA concentration was determined and the plasmid to oligonucleotide sequence was ligated in the ratio 3:1 insert to vector. The ligation reaction was done with T4 DNA ligase for 30 minutes at room temperature. 2.5 μ L of the ligation mix were added to 25 μ L of STBL3 competent bacterial cells. Transformation required heat shock at 42° C. Bacterial cells were spread on agar plates containing ampicillin and colonies were expanded in L broth. To check for insertion of the oligo sequences, plasmid DNA was extracted from harvested bacterial cultures using the Invitrogen DNA Miniprep kit and tested by restriction enzyme digestion. Insertion of the

shRNA sequence into the plasmid was verified by DNA sequencing using a primer specific for the promoter used to regulate shRNA expression.

[0292] Functional Assay for CCR5 mRNA Reduction:
[0293] The assay for inhibition of CCR5 expression required co-transfection of two plasmids. The first plasmid contains one of five different shRNA sequences directed against CCR5 mRNA. The second plasmid contains the cDNA sequence for human CCR5 gene. Plasmids were co-transfected into 293T cells. After 48 hours, cells were lysed and RNA was extracted using the RNeasy kit from Qiagen. cDNA was synthesized from RNA using a Super Script Kit from Invitrogen. The samples were then analyzed by quantitative RT-PCR using an Applied Biosystems Step One PCR machine. CCR5 expression was detected with SYBR Green from Invitrogen using the forward primer (5'-AGGAATTGATGGCGAGAAGG-3') (SEQ ID NO: 93) and reverse primer (5'-CCCCAAAGAAGGTCAAGG-TAATCA-3') (SEQ ID NO: 94) with standard conditions for polymerase chain reaction analysis. The samples were normalized to the mRNA for beta actin gene expression using the forward primer (5'-AGCGCGGCTACAGCTTCA-3') (SEQ ID NO: 95) and reverse primer (5'-GGCGACGTAG-CACAGCTTCT-3') (SEQ ID NO: 96) with standard conditions for polymerase chain reaction analysis. The relative expression of CCR5 mRNA was determined by its Ct value normalized to the level of actin messenger RNA for each sample. The results are shown in FIG. 10.

[0294] As shown in FIG. 10A, CCR5 knock-down was tested in 293T cells by co-transfection of the CCR5 shRNA construct and a CCR5-expressing plasmid. Control samples were transfected with a scrambled shRNA sequence that did not target any human gene and the CCR5-expressing plasmid. After 60 hours post-transfection, samples were harvested and CCR5 mRNA levels were measured by quantitative PCR. Further, as shown in FIG. 10B, CCR5 knock-down after transduction with lentivirus expressing CCR5 shRNA-1 (SEQ ID NO: 16).

Example 6: Regulation of HIV Components by shRNA Inhibitor Sequences in a Lentiviral Vector

[0295] Inhibitory RNA Design.
The sequences of HIV type 1 Rev/Tat (5'-GCGGAGACA-GCGACGAAGAGC-3') (SEQ ID NO: 9) and Gag (5'-GAAGAAATGATGACAGCAT-3') (SEQ ID NO: 11) were used to design:

Rev/Tat:

(5'GCGGAGACAGCGACGAAGAGCTTCAAGA-GAGCTCTTCGTCGCTGTCTCCGCTTT TT-3') (SEQ ID NO: 10) and

Gag:

[0296] (5'GAAGAAATGATGACAGCATTTCAAGA-GAATGCTGTCATCATTTCTTCTTTT-3') (SEQ ID NO: 12) shRNA that were synthesized and cloned into plasmids as described above.

[0297] Plasmid Construction.

[0298] The Rev/Tat or Gag target sequences were inserted into the 3'UTR (untranslated region) of the firefly luciferase gene used commonly as a reporter of gene expression in cells or tissues. Additionally, one plasmid was constructed to express the Rev/Tat shRNA and a second plasmid was

constructed to express the Gag shRNA. Plasmid constructions were as described above.

[0299] Functional Assay for shRNA Targeting of Rev/Tat or Gag mRNA:

[0300] Using plasmid co-transfection we tested whether a shRNA plasmid was capable of degrading luciferase messenger RNA and decreasing the intensity of light emission in co-transfected cells. A shRNA control (scrambled sequence) was used to establish the maximum yield of light from luciferase transfected cells. When the luciferase construct containing a Rev/Tat target sequence inserted into the 3'-UTR (untranslated region of the mRNA) was co-transfected with the Rev/Tat shRNA sequence there was nearly a 90% reduction in light emission indicating strong function of the shRNA sequence. A similar result was obtained when a luciferase construct containing a Gag target sequence in the 3'-UTR was co-transfected with the Gag shRNA sequence. These results indicate potent activity of the shRNA sequences.

[0301] As shown in FIG. 11A, knock-down of the Rev/Tat target gene was measured by a reduction of luciferase activity, which was fused with the target mRNA sequence in the 3'UTR, by transient transfection in 293T cells. As shown in FIG. 11B, knock-down of the Gag target gene sequence fused with the luciferase gene. The results are displayed as the mean±SD of three independent transfection experiments, each in triplicate.

Example 7: AGT103 Decreases Expression of Tat and Vif

[0302] Cells were transfected with exemplary vector AGT103/CMV-GFP. AGT103 and other exemplary vectors are defined in Table 3 below.

TABLE 3

Vector Designation	Composition
AGT103	EF1-miR30CCR5-miR21Vif-miR185-Tat-WPRE
Control-mCherry	CMV-mCherry
AGT103/CMV-mCherry	CMV-mCherry-EF1-miR30CCR5-miR21Vif-miR185-Tat-WPRE-
Control-GFP	CMV-mCherry
AGT103/CMV-GFP	CMV-GFP-EF1-miR30CCR5-miR21Vif-miR185-Tat-WPRE-

Abbreviations:
EF-1: elongation factor 1 transcriptional promoter
miR30CCR5 - synthetic microRNA capable of reducing CCR5 protein on cell surfaces
miR21Vif - synthetic microRNA capable of reducing levels of HIV RNA and Vif protein expression
miR185Tat - synthetic micro RNA capable of reducing levels of HIV RNA and Tat protein expression
CMV - Immediate early transcriptional promoter from human cytomegalovirus
mCherry - coding region for the mCherry red fluorescent protein
GFP - coding region for the green fluorescent protein
WPRE - Woodchuck hepatitis virus post transcriptional regulatory element

[0303] A T lymphoblastoid cell line (CEM; CCRF-CEM; American Type Culture Collection Catalogue number CCL119) was transduced with AGT103/CMV-GFP. 48 hours later the cells were transfected with an HIV expression plasmid encoding the entire viral sequence. After 24 hours, RNA was extracted from cells and tested for levels of intact Tat sequences using reverse transcriptase polymerase chain reaction. Relative expression levels for intact Tat RNA were reduced from approximately 850 in the presence of control

lentivirus vector, to approximately 200 in the presence of AGT103/CMV-GFP for a total reduction of >4 fold, as shown in FIG. 12.

Example 8: Regulation of HIV Components by Synthetic MicroRNA Sequences in a Lentiviral Vector

[0304] Inhibitory RNA Design.

[0305] The sequence of HIV-1 Tat and Vif genes were used to search for potential siRNA or shRNA candidates to knockdown Tat or Vif levels in human cells. Potential RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from the Broad Institute or the BLOCK-IT RNA iDesigner from Thermo Scientific. The selected shRNA sequences most potent for Tat or Vif knockdown were embedded within a microRNA backbone to allow for expression by an RNA polymerase II promoter such as CMV or EF-1 alpha. The RNA sequence may also be synthesized as a siRNA oligonucleotide and used independently of a plasmid or lentiviral vector.

[0306] Plasmid Construction.

[0307] The Tat target sequence (5'-TCCGCTTCTTCCTGCCATAG-3') (SEQ ID NO: 7) was incorporated into the miR185 backbone to create a Tat miRNA (5'-GGGCTG-GCTCGAGCAGGGGGCGAGGGATTCCGCTTCTTCCTGCCATAGCGTGGT CCCCTCCCCTATGGCAGGCA-GAAGCGGCACCTTCCCTCCCAATGACCGCGTCTTC G TCG-3') (SEQ ID NO: 3) that was inserted into a lentivirus vector and expressed under control of the EF-1 alpha promoter. Similarly, the Vif target sequence (5'-GGGATGTGTACTTCTGAACCT-3') (SEQ ID NO: 6) was incorporated into the miR21 backbone to create a Vif miRNA (5'-CATCTCCATGGCTGTACCACCTT-GTCGGGGGATGTGTACTTCTGAACCTTGTGTTGAATCTCATGGAGTTCAGAAGAACAACATCCGCACT-GACATTTTGGTATCTTTTCATCTG ACCA-3') (SEQ ID NO: 2) that was inserted into a lentivirus vector and expressed under control of the EF-1 alpha promoter. The resulting Vif/Tat miRNA-expressing lentivirus vectors were produced in 293T cells using a lentiviral vector packaging system. The Vif and Tat miRNA were embedded into a microRNA cluster consisting of miR CCR5, miR Vif, and miR Tat all expressed under control of the EF-1 promoter.

[0308] Functional Assay for miR185/Tat Inhibition of Tat mRNA Accumulation.

[0309] A lentivirus vector expressing miR185 Tat (LV-EF1-miR—CCR5-Vif-Tat) was used at a multiplicity of infection equal to 5 for transducing 293T cells. 24 hours after transduction the cells were transfected with a plasmid expressing HIV strain NL4-3 (pNL4-3) using Lipofectamine2000 under standard conditions. 24 hours later RNA was extracted and levels of Tat messenger RNA were tested by RT-PCR using Tat-specific primers and compared to actin mRNA levels for a control.

[0310] Functional Assay for miR21 Vif Inhibition of Vif Protein Accumulation.

[0311] A lentivirus vector expressing miR21 Vif (LV-EF1-miR—CCR5-Vif-Tat) was used at a multiplicity of infection equal to 5 for transducing 293T cells. 24 hours after transduction, the cells were transfected with a plasmid expressing HIV strain NL4-3 (pNL4-3) using Lipofectamine2000. 24 hours later cells were lysed and total soluble protein was tested to measure the content of Vif protein. Cell lysates

were separated by SDS-PAGE according to established techniques. The separated proteins were transferred to nylon membranes and probed with a Vif-specific monoclonal antibody or actin control antibody.

[0312] As shown in FIG. 13A, Tat knock-down was tested in 293T cells transduced with either a control lentiviral vector or a lentiviral vector expressing either synthetic miR185 Tat or miR155 Tat microRNA. After 24 hours, the HIV vector pNL4-3 was transfected with Lipofectamine2000 for 24 hours and then RNA was extracted for qPCR analysis with primers for Tat. As shown in FIG. 13B, Vif knock-down was tested in 293T cells transduced with either a control lentiviral vector or a lentiviral vector expressing a synthetic miR21 Vif microRNA. After 24 hours, the HIV vector pNL4-3 was transfected with Lipofectamine2000 for 24 hours and then protein was extracted for immunoblot analysis with an antibody for HIV Vif.

Example 9: Regulation of CCR5 Expression by Synthetic microRNA Sequences in a Lentiviral Vector

[0313] CEM-CCR5 cells were transduced with a lentiviral vector containing a synthetic miR30 sequence for CCR5 (AGT103: TGTAAGCTGAGCTTGCTCTA (SEQ ID NO: 97), AGT103-R5-1: TGTAAGCTGAGCTTGCTCGC (SEQ ID NO: 98), or AGT103-R5-2: CATAGATTGGACTTGACAC (SEQ ID NO: 99). After 6 days, CCR5 expression was determined by FACS analysis with an APC-conjugated CCR5 antibody and quantified by mean fluorescence intensity (MFI). CCR5 levels were expressed as % CCR5 with LV-Control set at 100%. The target sequence of AGT103 and AGT103-R5-1 is in the same region as CCR5 target sequence #5. The target sequence of AGT103-R5-2 is the same as CCR5 target sequence #1. AGT103 (2% of total CCR5) is most effective at reducing CCR5 levels as compared with AGT103-R5-1 (39% of total CCR5) and AGT103-R5-2 which does not reduce CCR5 levels. The data is demonstrated in FIG. 14 herein.

Example 10: Regulation of CCR5 Expression by Synthetic microRNA Sequences in a Lentiviral Vector Containing Either a Long or Short WPRE Sequence

[0314] Vector Construction.

[0315] Lentivirus vectors often require an RNA regulatory element for optimal expression of therapeutic genes or genetic constructs. A common choice is to use the Woodchuck hepatitis virus post transcriptional regulatory element (WPRE). We compared AGT103 that contains a full-length WPRE:

(SEQ ID NO: 32)
(5' AATCAACCTCTGATTACAAAATTTGTGAAGATTGACTGGTATTCTT
AACTATGTTGCTCCTTTTACGCTATGTGGATACGCTGCTTTAATGCCTTT
GTATCATGCTATTGCTTCCCGTATGGCTTTTCTCTCTCTCTGTATA
AATCCTGGTGTGCTGTCTTTATGAGGAGTTGTGGCCCGTTGTGAGGCAA
CGTGGCGTGGTGTGCACTGTGTTTGTGACGCAACCCCACTGGTTGGGG
CATTGCCACCACCTGTGCACTCCTTTCCGGGACTTTCGCTTCCCCCTCC

-continued

CTATTGCCACGGCGAACTCATCGCCGCTGCCTGCCCGCTGCTGGACA
GGGGCTCGGCTGTTGGGCACTGACAATTCCGTGGTGTGTCGGGAAATC
ATCGTCCTTTCTTGGCTGCTCGCCTGTGTTGCCACCTGGATTCTGCGCG
GGACGTCCTTCTGCTACGTCCTTCGGCCCTCAATCCAGCGGACCTTCCT
TCCCGCGGCTGCTGCCGGCTCTGCGGCTCTTCCGCGTCTTCGCTTCG
CCCTCAGACGAGTCGGATCTCCCTTTGGGCCGCTCCCCGCT-3')

with a modified AGT103 vector containing a shortened
WPRE element

(SEQ ID NO: 80)
(5'AATCAACCTCTGGATTACAAAATTTGTGAAAGATTGACTGATATTC
TAACTATGTTGCTCCTTTTACGCTGTGTGGATATGCTGCTTAATGCCTC
TGTATCATGCTATTGCTTCCGTACGGCTTTCGTTTTCTCCTCCTGTAT
AAATCCTGGTGTGCTGCTCTTTATGAGGAGTTGTGGCCGTTGTCCGTCA
ACGTGGCGTGGTGTGCTCTGTGTTGCTGACGCAACCCCCACTGGCTGGG
GCATTGCCACCACCTGTCAACTCCTTTCTGGGACTTTCGCTTTCCCCCTC
CCGATCGCCACGGCAGAACTCATCGCCGCTGCCTTGCCGCTGCTGGAC
AGGGGCTAGGTTGCTGGGCACTGATAATCCGTGGTGTGTC-3').

[0316] Functional Assay for Modulating Cell Surface
CCR5 Expression as a Function of Long Versus Short
WPRE Element in the Vector Sequence.

[0317] AGT103 containing long or short WPRE elements
were used for transducing CEM-CCR5 T cells a multiplicity
of infection equal to 5. Six days after transduction cells were
collected and stained with a monoclonal antibody capable of
detecting cell surface CCR5 protein. The antibody was
conjugated to a fluorescent marker and the intensity of
staining is directly proportional to the level of CCR5 on the
cell surface. A control lentivirus had no effect on cell surface
CCR5 levels resulting in a single population with a mean
fluorescence intensity of 73.6 units. The conventional
AGT103 with a long WPRE element reduced CCR5 expres-
sion to a mean fluorescence intensity level of 11 units.
AGT103 modified to incorporate a short WPRE element
resulted in a single population of cells with mean fluores-
cence intensity of 13 units. Accordingly, substituting a short
WPRE element had little or no effect on the capacity for
AGT103 to reduce cell surface CCR5 expression.

[0318] As shown in FIG. 14, CEM-CCR5 cells were
transduced with AGT103 containing either a long or short
WPRE sequence. After 6 days, CCR5 expression was deter-
mined by FACS analysis with an APC-conjugated CCR5
antibody and quantified as mean fluorescence intensity
(MFI). CCR5 levels were expressed as % CCR5 with
LV-Control set at 100%. The reduction in CCR5 levels was
similar for AGT103 with either the short (5.5% of total
CCR5) or long (2.3% of total CCR5) WPRE sequence.

Example 11: Regulation of CCR5 Expression by
Synthetic microRNA Sequences in a Lentiviral
Vector with or without a WPRE Sequence

[0319] Vector Construction.

[0320] In order to test whether WPRE was required for
AGT103 down regulation of CCR5 expression we con-
structed a modified vector without WPRE element
sequences.

[0321] Functional assay for modulating cell surface CCR5
expression as a function of including or not including a long
WPRE element in the AGT103 vector. In order to test
whether WPRE was required for AGT103 modulation of
CCR5 expression levels we transduced CEM-CCR5 T cells
with AGT103 or a modified vector lacking WPRE using a
multiplicity of infection equal to 5. Six days after transduc-
tion cells were collected and stained with a monoclonal
antibody capable of recognizing cell surface CCR5 protein.
The monoclonal antibody was directly conjugated to a
fluorescent marker and the intensity of staining is directly
proportional to the number of CCR5 molecules per cell
surface. A lentivirus control vector had no effect on cell
surface CCR5 levels resulting in a uniform population with
mean fluorescence intensity of 164. The lentivirus vector
(AGT103 with a long WPRE and also expressing GFP
marker protein), AGT103 lacking GFP but containing a long
WPRE element, or AGT103 lacking both GFP and WPRE
all were similarly effective for modulating cell surface
CCR5 expression. After removing GFP, AGT103 with or
without WPRE elements were indistinguishable in terms of
their capacity for modulating cell surface CCR5 expression.

[0322] CEM-CCR5 cells were transduced with AGT103
with or without GFP and WPRE. After 6 days, CCR5
expression was determined by FACS analysis with an APC-
conjugated CCR5 antibody and quantified as mean fluores-
cence intensity (MFI). CCR5 levels were expressed as %
CCR5 with LV-Control set at 100%. The reduction in CCR5
levels was similar for AGT103 with (0% of total CCR5) or
without (0% of total CCR5) the WPRE sequence. This data
is demonstrated in FIG. 16.

Example 12: Regulation of CCR5 Expression by a
CD4 Promoter Regulating Synthetic microRNA
Sequences in a Lentiviral Vector

[0323] Vector Construction.

[0324] A modified version of AGT103 was constructed to
test the effect of substituting alternate promoters for express-
ing the microRNA cluster that suppresses CCR5, Vif and Tat
gene expression. In place of the normal EF-1 promoter we
substituted the T cell-specific promoter for CD4 glycopro-
tein expression using the sequence:

(SEQ ID NO: 30)
(5'TGTTGGGGTTCAAATTTGAGCCCCAGCTGTAGCCCTCTGCAAGAA
AAAAAAAAAAAAAAAAAGAACAAAGGGCCTAGATTTCCCTTCTGAGCCCCA
CCCTAAGATGAAGCCTCTTCTTTCAAGGGAGTGGGGTGGGGTGGAGCG
GATCCTGTGACGCTTTGCTCTCTGTGGCTGGCAGTTTCTCCAAGGGTA
ACAGGTGTGACGCTGGCTGAGCCTAGGCTGAACCTGAGACATGCTACCTC
TGTCTTCTCATGGCTGGAGGCAGCCTTTGTAAGTCACAGAAAGTAGCTGA

-continued

GGGGCTCTGGAAAAAGACAGCCAGGGTGGAGGTAGATTGGTCTTTGACT
CCTGATTAAAGCCTGATTCTGCTTAACCTTTTCCCTTGACTTTGGCATT
TCACTTTGACATGTTCCCTGAGAGCCTGGGGGTGGGAACCCAGCTCCA
GCTGGTGACGTTTGGGGCCGGCCAGGCCTAGGGTGTGGAGGAGCCTTGC
CATCGGGCTTCTGTCTCTCTTCATTTAAGCAGACTCTGCAGA-3') .

[0325] Functional Assay Comparing EF-1 and CD4 Gene Promoters in Terms of Potency for Reducing Cell Surface CCR5 Protein Expression.

[0326] AGT103 modified by substituting the CD4 gene promoter for the normal EF-1 promoter was used for transducing CEM-CCR5 T cells. Six days after transduction cells were collected and stained with a monoclonal antibody capable of recognizing cell surface CCR5 protein. The monoclonal antibody was conjugated to a fluorescent marker and staining intensity is directly proportional to the level of cell surface CCR5 protein. A control lentivirus transduction resulted in a population of CEM-CCR5 T cells that were stained with a CCR5-specific monoclonal antibody and produced a mean fluorescence intensity of 81.7 units. The modified AGT103 using a CD4 gene promoter in place of the EF-1 promoter for expressing microRNA showed a broad distribution of staining with a mean fluorescence intensity roughly equal to 17.3 units. Based on this result, the EF-1 promoter is at least similar and likely superior to the CD4 gene promoter for microRNA expression. Depending on the desired target cell population, the EF-1 promoter is universally active in all cell types and the CD4 promoter is only active in T-lymphocytes.

[0327] CEM-CCR5 cells were transduced with a lentiviral vector containing a CD4 promoter regulating a synthetic microRNA sequence for CCR5, Vif, and Tat (AGT103). After 6 days, CCR5 expression was determined by FACS analysis with an APC-conjugated CCR5 antibody and quantified as mean fluorescence intensity (MFI). CCR5 levels were expressed as % CCR5 with LV-Control set at 100%. In cells transduced with LV-CD4-AGT103, CCR5 levels were 11% of total CCR5. This is comparable to that observed for LV-AGT103 which contains the EF1 promoter. This data is demonstrated in FIG. 17.

Example 13: Detecting HIV Gag-Specific CD4 T Cells

[0328] Cells and Reagents.

[0329] Viable frozen peripheral blood mononuclear cells (PBMC) were obtained from a vaccine company. Data were obtained with a representative specimen from an HIV+ individual who was enrolled into an early stage clinical trial (TRIAL REGISTRATION: clinicaltrials.gov NCT01378156) testing a candidate HIV therapeutic vaccine. Two specimens were obtained for the "Before vaccination" and "After vaccination" studies. Cell culture products, supplements and cytokines were from commercial suppliers. Cells were tested for responses to recombinant Modified Vaccinia Ankara 62B from Geovax Corporation as described in Thompson et al. (2016). "DNA/MVA Vaccination of HIV-1 Infected Participants with Viral Suppression on Antiretroviral Therapy, followed by Treatment Interruption: Elicitation of Immune Responses without Control of Re-Emergent Virus." *PLoS One* 11(10): e0163164. Synthetic peptides representing the entire HIV-1 Gag polyprotein were

obtained from GeoVax the HIV (GAG) Ultra peptide sets were obtained from JPT Peptide Technologies GmbH (www.jpt.com), Berlin, Germany. HIV (GAG) Ultra contains 150 peptides each being 15 amino acids in length and overlapping by 11 amino acids. They were chemically synthesized then purified and analyzed by liquid chromatography—mass spectrometry. Collectively these peptides represent major immunogenic regions of the HIV Gag polyprotein and are designed for average coverage of 57.8% among known HIV strains. Peptide sequences are based on the HIV sequence database from the Los Alamos National Laboratory (<http://www.hiv.lanl.gov/content/sequence/NEALIGN/align.html>). Peptides are provided as dried trifluoroacetate salts, 25 micrograms per peptide, and are dissolved in approximately 40 microliters of DMSO then diluted with PBS to final concentration. Monoclonal antibodies for detecting CD4 and cytoplasmic IFN-gamma were obtained from commercial sources and intracellular staining was done with the BD Pharmingen Intracellular Staining Kit for interferon-gamma. Peptides were resuspended in DMSO and we include a DMSO only control condition.

[0330] Functional Assay for Detecting HIV-Specific CD4+ T Cells.

[0331] Frozen PBMC were thawed, washed and resuspended in RPMI medium containing 10% fetal bovine serum, supplements and cytokines. Cultured PBMC collected before or after vaccination were treated with DMSO control, MVA GeoVax (multiplicity of infection equal to 1 plaque forming unit per cell), Peptides GeoVax (1 microgram/ml) or HIV (GAG) Ultra peptide mixture (1 microgram/ml) for 20 hours in the presence of Golgi Stop reagent. Cells were collected, washed, fixed, permeabilized and stained with monoclonal antibodies specific for cell surface CD4 or intracellular interferon-gamma. Stained cells were analyzed with a FACSCalibur analytical flow cytometer and data were gated on the CD4+ T cell subset. Cells highlighted within boxed regions are double-positive and designated HIV-specific CD4 T cells on the basis of interferon-gamma expression after MVA or peptide stimulation. Numbers within the boxed regions show the percentage of total CD4 that were identified as HIV-specific. We did not detect strong responses to DMSO or MVA. Peptides from GeoVax elicited fewer responding cells compared to HIV (GAG) Ultra peptide mixture from JPT but differences were small and not significant.

[0332] As shown in FIG. 18, PBMCs from a HIV-positive patient before or after vaccination were stimulated with DMSO (control), recombinant MVA expressing HIV Gag from GeoVax (MVA GeoVax), Gag peptide from GeoVax (Pep GeoVax, also referred to herein as Gag peptide pool 1) or Gag peptides from JPT (HIV (GAG) Ultra, also referred to herein as Gag peptide pool 2) for 20 hours. IFN γ production was detected by intracellular staining and flow cytometry using standard protocols. Flow cytometry data were gated on CD4 T cells. Numbers captured in boxes are the percentage of total CD4 T cells designated "HIV-specific" on the basis of cytokine response to antigen-specific stimulation.

Example 14: HIV-Specific CD4 T Cell Expansion and Lentivirus Transduction

[0333] Designing and testing methods for enriching PBMC to increase the proportion of HIV-specific CD4 T

cells and transducing these cells with AGT103 to produce the cellular product AGT103T.

[0334] The protocol was designed for ex vivo culture of PBMC (peripheral blood mononuclear cells) from HIV-positive patients who had received a therapeutic HIV vaccine. In this example, the therapeutic vaccine consisted of three doses of plasmid DNA expressing HIV Gag, Pol and Env genes followed by two doses of MVA 62-B (modified vaccinia Ankara number 62-B) expressing the same HIV Gag, Pol, and Env genes. The protocol is not specific for a vaccine product and only requires a sufficient level of HIV-specific CD4⁺ T cells after immunization. Venous blood was collected and PBMC were purified by Ficoll-Paque density gradient centrifugation. Alternately, PBMC or defined cellular fractions can be prepared by positive or negative selection methods using antibody cocktails and fluorescence activated or magnetic bead sorting. The purified PBMC are washed and cultured in standard medium containing supplements, antibiotics and fetal bovine serum. To these cultures, a pool of synthetic peptides was added representing possible T cell epitopes within the HIV Gag polyprotein. Cultures are supplemented by adding cytokines interleukin-2 and interleukin-12 that were selected after testing combinations of interleukin-2 and interleukin-12, interleukin 2 and interleukin-7, interleukin 2 and interleukin-15. Peptide stimulation is followed by a culture interval of approximately 12 days. During the 12 days culture, fresh medium and fresh cytokine supplements were added approximately once every four days.

[0335] The peptide stimulation interval is designed to increase the frequency of HIV-specific CD4 T cells in the PBMC culture. These HIV-specific CD4 T cells were activated by prior therapeutic immunization and can be re-stimulated and caused to proliferate by synthetic peptide exposure. Our goal is to achieve greater than or equal to 1% of total CD4 T cells being HIV-specific by end of the peptide stimulation culture period.

[0336] On approximately day 12 of culture cells are washed to remove residual materials then stimulated with synthetic beads decorated with antibodies against CD4 T cell surface proteins CD3 and CD28. This well-established method for polyclonal stimulation of T cells will reactivate the cells and make them more susceptible for AGT103 lentivirus transduction. The lentivirus transduction is performed on approximately day 13 of culture and uses a multiplicity of infection between 1 and 5. After transduction cells are washed to remove residual lentivirus vector and cultured in media containing interleukin-2 and interleukin-12 with fresh medium and cytokines added approximately once every four days until approximately day 24 of culture.

[0337] Throughout the culture interval the antiretroviral drug Saquinavir is added at a concentration of approximately 100 nM to suppress any possible outgrowth of HIV.

[0338] On approximately day 24 of culture cells are harvested, washed, a sample is set aside for potency and release assay, then the remaining cells are suspended in cryopreservation medium before freezing in single aliquots of approximately 1×10^{10} cells per dose that will contain approximately 1×10^8 HIV-specific CD4 T cells that are transduced with AGT103.

[0339] Potency of the cell product (AGT103T) is tested in one of two alternate potency assays. Potency assay 1 tests for the average number of genome copies (integrated AGT103 vector sequences) per CD4 T cell. The minimum

potency is approximately 0.5 genome copies per CD4 T cell in order to release the product. The assay is performed by positive selection of CD3 positive/CD4 positive T cells using magnetic bead labeled monoclonal antibodies, extracting total cellular DNA and using a quantitative PCR reaction to detect sequences unique to the AGT103 vector. Potency assay 2 tests for the average number of genome copies of integrated AGT103 within the subpopulation of HIV-specific CD4 T cells. This assay is accomplished by first stimulating the PBMC with the pool of synthetic peptides representing HIV Gag protein. Cells are then stained with a specific antibody reagent capable of binding to the CD4 T cell and also capturing secreted interferon-gamma cytokine. The CD4 positive/interferon-gamma positive cells are captured by magnetic bead selection, total cellular DNA is prepared, and the number of genome copies of AGT103 per cell is determined with a quantitative PCR reaction. Release criterion based on potency using Assay 2 require that greater than or equal to 0.5 genome copies per HIV-specific CD4 T-cell are present in the AGT103 cell product.

[0340] Functional Test for Enriching and Transducing HIV-Specific CD4 T Cells from PBMC of HIV-Positive Patients that Received a Therapeutic HIV Vaccine.

[0341] The impact of therapeutic vaccination on the frequency of HIV-specific CD4 T cells was tested by a peptide stimulation assay (FIG. 19, Panel B). Before vaccination the frequency of HIV-specific CD4 T cells was 0.036% in this representative individual. After vaccination, the frequency of HIV-specific CD4 T cells was increased approximately 2-fold to the value of 0.076%. Responding cells (HIV-specific) identified by accumulation of cytoplasmic interferon-gamma, were only detected after specific peptide stimulation.

[0342] We also tested whether peptide stimulation to enrich for HIV-specific CD4 T cells followed by AGT103 transduction would reach our goal of generating approximately 1% of total CD4 T cells in culture that were both HIV-specific and transduced by AGT103. In this case, we used an experimental version of AGT103 that expresses green fluorescence protein (see GFP). In FIG. 19, Panel C the post-vaccination culture after peptide stimulation (HIV (GAG) Ultra) and AGT103 transduction demonstrated that 1.11% of total CD4 T cells were both HIV-specific (based on expressing interferon-gamma in response to peptide stimulation) and AGT103 transduced (based on expression of GFP).

[0343] Several patients from a therapeutic HIV vaccine study were tested to assess the range of responses to peptide stimulation and to begin defining eligibility criteria for entering a gene therapy arm in a future human clinical trial. FIG. 19 Panel D show the frequency of HIV-specific CD4 T cells in 4 vaccine trial participants comparing their pre- and post-vaccination specimens. In three cases the post-vaccination specimens show a value of HIV-specific CD4 T cells that was greater than or equal to 0.076% of total CD4 T cells. The ability to reach this value was not predicted by the pre-vaccination specimens as patient 001-004 and patient 001-006 both started with pre-vaccination values of 0.02% HIV-specific CD4 T cells but one reached an eventual post-vaccination value of 0.12% HIV-specific CD4 T cells while the other individual fail to increase this value after vaccination. The same three patients that responded well to vaccine, in terms of increasing the frequency of HIV-specific CD4 T cells, also showed substantial enrichment of HIV-

specific CD4 T cells after peptide stimulation and culture. In the three cases shown in FIG. 19 Panel E, peptide stimulation and subsequent culture generated samples where 2.07%, 0.72% or 1.54% respectively of total CD4 T cells were HIV-specific. These values indicate that a majority of individuals responding to a therapeutic HIV vaccine will have a sufficiently large ex vivo response to peptide stimulation in order to enable our goal of achieving approximately 1% of total CD4 T cells that are HIV-specific and transduced with AGT103 in the final cell product.

[0344] As shown in FIG. 19, Panel A describes the schedule of treatment. Panel B demonstrates that PBMCs were stimulated with Gag peptide or DMSO control for 20 hours. IFN gamma production was detected by intracellular staining by FACS. CD4⁺ T cells were gated for analysis. Panel C demonstrates CD4⁺ T cells were expanded and transduced with AGT103-GFP using the method as shown in Panel A. Expanded CD4⁺ T cells were rested in fresh medium without any cytokine for 2 days and re-stimulated with Gag peptide or DMSO control for 20 hours. IFN gamma production and GFP expression was detected by FACS. CD4⁺ T cells were gated for analysis. Panel D demonstrates frequency of HIV-specific CD4⁺ T cells (IFN gamma positive, pre- and post-vaccination) were detected from 4 patients. Panel E demonstrates Post-vaccination PBMCs from 4 patients were expanded and HIV-specific CD4⁺ T cells were examined.

Example 15: Dose Response

[0345] Vector Construction.

[0346] A modified version of AGT103 was constructed to test the dose response for increasing AGT103 and its effects on cell surface CCR5 levels. The AGT103 was modified to include a green fluorescent protein (GFP) expression cassette under control of the CMV promoter. Transduced cells expression the miR30CCR5 miR21Vif miR185Tat micro RNA cluster and emit green light due to expressing GFP.

[0347] Functional Assay for Dose Response of Increasing AGT103-GFP and Inhibition of CCR5 Expression.

[0348] CEM-CCR5 T cells were transduced with AGT103-GFP using multiplicity of infection per cell from 0 to 5. Transduced cells were stained with a fluorescently conjugated (APC) monoclonal antibody specific for cell surface CCR5. The intensity of staining is proportional to the number of CCR5 molecules per cell surface. The intensity of green fluorescence is proportional to the number of integrated AGT103-GFP copies per cell.

[0349] As shown in FIG. 20, Panel A demonstrates the dose response for increasing AGT103-GFP and its effects on cell surface CCR5 expression. At multiplicity of infection equal to 0.4 only 1.04% of cells are both green (indicating transduction) and showing significantly reduced CCR5 expression. At multiplicity of infection equal to 1 the number of CCR5low, GFP+ cells increases to 68.1%. At multiplicity of infection equal to 5 the number of CCR5low, GFP+ cells increased to 95.7%. These data are presented in histogram form in FIG. 20, Panel B that shows a normally distribution population in terms of CCR5 staining, moving toward lower mean fluorescence intensity with increasing doses of AGT103-GFP. The potency of AGT103-GFP is presented in graphical form in FIG. 20, Panel C showing the percentage inhibition of CCR5 expression with increasing doses of AGT103-GFP. At multiplicity of infection equal to 5, there was greater than 99% reduction in CCR5 expression levels.

Example 16: AGT103 Efficiently Transduces Primary Human CD4⁺ T Cells

[0350] Transducing Primary CD4 T Cells with AGT103 Lentivirus Vector.

[0351] A modified AGT103 vector containing the green fluorescence protein marker (GFP) was used at multiplicities of infection between 0.2 and 5 for transducing purified, primary human CD4 T cells.

[0352] Functional Assay for Transduction Efficiency of AGT103 in Primary Human CD4 T Cells.

[0353] CD4 T cells were isolated from human PBMC (HIV-negative donor) using magnetic bead labeled antibodies and standard procedures. The purified CD4 T cells were stimulated ex vivo with CD3/CD28 beads and cultured in media containing interleukin-2 for 1 day before AGT103 transduction. The relationship between lentivirus vector dose (the multiplicity of infection) and transduction efficiency is demonstrated in FIG. 21, Panel A showing that multiplicity of infection equal to 0.2 resulted in 9.27% of CD4 positive T cells being transduced by AGT103 and that value was increased to 63.1% of CD4 positive T cells being transduced by AGT103 with a multiplicity of infection equal to 5. In addition to achieving efficient transduction of primary CD4 positive T cells it is also necessary to quantify the number of genome copies per cell. In FIG. 21, Panel B total cellular DNA from primary human CD4 T cells transduced at several multiplicities of infection were tested by quantitative PCR to determine the number of genome copies per cell. In a multiplicity of infection equal to 0.2 we measured 0.096 genome copies per cell that was in good agreement with 9.27% GFP positive CD4 T cells in panel A. Multiplicity of infection equal to 1 generated 0.691 genome copies per cell and multiplicity of infection equal to 5 generated 1.245 genome copies per cell.

[0354] As shown in FIG. 21, CD4⁺ T cells isolated from PBMC were stimulated with CD3/CD28 beads plus IL-2 for 1 day and transduced with AGT103 at various concentrations. After 2 days, beads were removed and CD4⁺ T cells were collected. As shown in Panel A, frequency of transduced cells (GFP positive) were detected by FACS. As shown in Panel B, the number of vector copies per cell was determined by qPCR. At a multiplicity of infection (MOI) of 5, 63% of CD4⁺ T cells were transduced with an average of 1 vector copy per cell.

Example 17: AGT103 Inhibits HIV Replication in Primary CD4⁺ T Cells

[0355] Protecting Primary Human CD4 Positive T Cells from HIV Infection by Transducing Cells with AGT103.

[0356] Therapeutic lentivirus AGT103 was used for transducing primary human CD4 positive T cells at multiplicities of infection between 0.2 and 5 per cell. The transduced cells were then challenged with a CXCR4-tropic HIV strain NL4.3 that does not require cell surface CCR5 for penetration. This assay tests the potency of microRNA against Vif and Tat genes of HIV in terms of preventing productive infection in primary CD4 positive T cells, but uses an indirect method to detect the amount of HIV released from infected, primary human CD4 T cells.

[0357] Functional Assay for AGT103 Protection Against CXCR4-Tropic HIV Infection of Primary Human CD4 Positive T Cells.

[0358] CD4 T cells were isolated from human PBMC (HIV-negative donor) using magnetic bead labeled antibodies and standard procedures. The purified CD4 T cells were stimulated ex vivo with CD3/CD28 beads and cultured in media containing interleukin-2 for 1 day before AGT103 transduction using multiplicities of infection between 0.2 and 5. Two days after transduction the CD4 positive T cell cultures were challenged with HIV strain NL4.3 that was engineered to express the green fluorescent protein (GFP). The transduced and HIV-exposed primary CD4 T cell cultures were maintained for 7 days before collecting cell-free culture fluids containing HIV. The cell-free culture fluids were used to infect a highly permissive T cell line C8166 for 2 days. The proportion of HIV-infected C8166 cells was determined by flow cytometry detecting GFP fluorescence. With a mock lentivirus infection, the dose of 0.1 multiplicity of infection for NL4.3 HIV resulted in an amount of HIV being released into culture fluids that was capable of establishing productive infection in 15.4% of C8166 T cells. With the dose 0.2 multiplicity of infection for AGT103, this value for HIV infection of C8166 cells is reduced to 5.3% and multiplicity of infection equal to 1 for AGT103 resulted in only 3.19% of C8166 T cells being infected by HIV. C8166 infection was reduced further to 0.62% after AGT103 transduction using a multiplicity of infection equal to 5. There is a clear dose response relationship between the amount of AGT103 used for transduction and the amount of HIV released into the culture medium.

[0359] As shown in FIG. 22, CD4⁺ T cells isolated from PBMC were stimulated with CD3/CD28 beads plus IL-2 for 1 day and transduced with AGT103 at various concentrations (MOI). After 2 days, beads were removed and CD4⁺ T cells were infected with 0.1 MOI of HIV NL4.3-GFP. 24 hours later, cells were washed 3 times with PBS and cultured with IL-2 (30U/ml) for 7 days. At the end of the culture, supernatant was collected to infect the HIV permissive cell line C8166 for 2 days. HIV-infected C8166 cells (GFP positive) were detected by FACS. There was a reduction in viable HIV with an increase in the multiplicity of infection of AGT103 as observed by less infection of C8166 cells (MOI 0.2=65.6%, MOI 1=79.3%, and MOI 5=96%).

Example 18: AGT103 Protects Primary Human CD4⁺ T Cells from HIV-Induced Depletion

[0360] AGT103 Transduction of Primary Human CD4 T Cells to Protect Against HIV-Mediated Cytopathology and Cell Depletion.

[0361] PBMC were obtained from healthy, HIV-negative donors and stimulated with CD3/CD28 beads then cultured for 1 day in medium containing interleukin-2 before AGT103 transduction using multiplicities of infection between 0.2 and 5.

[0362] Functional Assay for AGT103 Protection of Primary Human CD4 T Cells Against HIV-Mediated Cytopathology.

[0363] AGT103-transduced primary human CD4 T cells were infected with HIV NL 4.3 strain (CXCR4-tropic) that does not require CCR5 for cellular entry. When using the CXCR4-tropic NL 4.3, only the effect of Vif and Tat microRNA on HIV replication is being tested. The dose of HIV NL 4.3 was 0.1 multiplicity of infection. One day after

HIV infection, cells were washed to remove residual virus and cultured in medium plus interleukin-2. Cells were collected every three days during a 14-day culture then stained with a monoclonal antibody that was specific for CD4 and directly conjugated to a fluorescent marker to allow measurement of the proportion of CD4 positive T cells in PBMC. Untreated CD4 T cells or CD4 T cells transduced with the control lentivirus vector were highly susceptible to HIV challenge and the proportion of CD4 positive T cells in PBMC fell below 10% by day 14 culture. In contrast, there was a dose-dependent effect of AGT103 on preventing cell depletion by HIV challenge. With a AGT103 dose of 0.2 multiplicity of infection more than 20% of PBMC were CD4 T cells by day 14 of culture and this value increased to more than 50% of PBMC being CD4 positive T cells by day 14 of culture with a AGT103 dose of multiplicity of infection equal to 5. Again, there is a clear dose response effect of AGT103 on HIV cytopathogenicity in human PBMC.

[0364] As shown in FIG. 23, PBMCs were stimulated with CD3/CD28 beads plus IL-2 for 1 day and transduced with AGT103 at various concentrations (MOI). After 2 days, beads were removed and cells were infected with 0.1 MOI of HIV NL4.3. 24 hours later, cells were washed 3 times with PBS and cultured with IL-2 (30U/ml). Cells were collected every 3 days and the frequency of CD4⁺ T cells were analyzed by FACS. After 14 days of exposure to HIV, there was an 87% reduction in CD4⁺ T cells transduced with LV-Control, a 60% reduction with AGT103 MOI 0.2, a 37% reduction with AGT103 MOI 1, and a 17% reduction with AGT103 MOI 5.

Example 19: Generating a Population of CD4⁺ T Cells Enriched for HIV-Specificity and Transduced with AGT103/CMV-GFP

[0365] Therapeutic vaccination against HIV had minimal effect on the distribution of CD4⁺, CD8⁺ and CD4⁺/CD8⁺ T cells. As shown in FIG. 24A, the CD4 T cell population is shown in the upper left quadrant of the analytical flow cytometry dot plots, and changes from 52% to 57% of total T cells after the vaccination series. These are representative data.

[0366] Peripheral blood mononuclear cells from a participant in an HIV therapeutic vaccine trial were cultured for 12 days in medium+/-interleukin-2/interleukin-12 or +/-interleukin-7/interleukin-15. Some cultures were stimulated with overlapping peptides representing the entire p55 Gag protein of HIV-1 (HIV (GAG) Ultra peptide mixture) as a source of epitope peptides for T cell stimulation. These peptides are 10-20 amino acids in length and overlap by 20-50% of their length to represent the entire Gag precursor protein (p55) from HIV-1 BaL strain. The composition and sequence of individual peptides can be adjusted to compensate for regional variations in the predominant circulating HIV sequences or when detailed sequence information is available for an individual patient receiving this therapy. At culture end, cells were recovered and stained with anti-CD4 or anti-CD8 monoclonal antibodies and the CD3⁺ population was gated and displayed here. The HIV (GAG) Ultra peptide mixture stimulation for either pre- or post-vaccination samples was similar to the medium control indicating that HIV (GAG) Ultra peptide mixture was not toxic to cells and was not acting as a polyclonal mitogen. The results of this analysis can be found in FIG. 24B.

[0367] HIV (GAG) Ultra peptide mixture and interleukin-2/interleukin-12 provided for optimal expansion of antigen-specific CD4 T cells. As shown in the upper panels of FIG. 24C, there was an increase in cytokine (interferon-gamma) secreting cells in post-vaccination specimens exposed to HIV (GAG) Ultra peptide mixture. In the pre-vaccination sample, cytokine secreting cells increased from 0.43 to 0.69% as a result of exposure to antigenic peptides. In contrast, the post-vaccination samples showed an increase of cytokine secreting cells from 0.62 to 1.76% of total CD4 T cells as a result of peptide stimulation. These data demonstrate the strong impact of vaccination on the CD4 T cell responses to HIV antigen.

[0368] Finally, AGT103/CMV-GFP transduction of antigen-expanded CD4 T cells produced HIV-specific and HIV-resistant helper CD4 T cells that are needed for infusion into patients as part of a functional cure for HIV (in accordance with other various aspects and embodiments, AGT103 alone is used; for example, clinical embodiments may not include the CMV-GFP segment). The upper panels of FIG. 24D show the results of analyzing the CD4+ T cell population in culture. The x axis of FIG. 24D shows Green Fluorescent Protein (GFP) emission indicating that individual cells were transduced with the AGT103/CMV-GFP. As shown in FIG. 24D, in the post-vaccination samples 1.11% of total CD4 T cells that were both cytokine secreting was recovered, indicating that the cells are responding specifically to HIV antigen, and transduced with AGT103/CMV-GFP. This is the target cell population and the clinical product intended for infusion and functional cure of HIV. With the efficiency of cell expansion during the antigen stimulation and subsequent polyclonal expansion phases of ex vivo culture, 4×10^8 antigen-specific, lentivirus transduced CD4 T cells can be produced. This exceeds the target for cell production by 4-fold and will allow achievement of a count of antigen-specific and HIV-resistant CD4 T cells of approximately 40 cells/microliter of blood or around 5.7% of total circulating CD4 T cells.

[0369] Table 4 below shows the results of the ex vivo production of HIV-specific and HIV-resistant CD4 T cells using the disclosed vectors and methods.

TABLE 4

Material/manipulation	Total CD4 T cells	Percentage HIV-specific	Percentage HIV-specific and HIV-resistant
Leukapheresis pack from HIV + patient	$\sim 7 \times 10^8$	~ 0.12	N/A
Peptide expansion ex vivo	$\sim 8 \times 10^8$	~ 2.4	N/A
Mitogen expansion	$\sim 1.5 \times 10^{10}$	~ 2.4	N/A
Lentivirus transduction	$\sim 1.5 \times 10^{10}$	~ 2.4	~ 1.6

Example 20: Clinical Study for Treatment of HIV

[0370] AGT103T is a genetically modified autologous PBMC containing $>5 \times 10^7$ HIV-specific CD4 T cells that are also transduced with AGT103 lentivirus vector.

[0371] A Phase I clinical trial will test the safety and feasibility of infusing ex vivo modified autologous CD4 T cells (AGT103T) in adult research participants with confirmed HIV infection, CD4+ T-cell counts >600 cells per mm^3 of blood and stable virus suppression below 200 copies per ml of plasma while on cART. All study participants will

continue receiving their standard antiretroviral medications through the Phase I clinical trial. Up to 40 study participants receive two doses by intramuscular injection 8 weeks apart, of recombinant modified vaccinia Ankara (rMVA) expressing HIV Gag, Pol and Env proteins. Seven to 10 days after the second immunization a blood sample is collected for in vitro testing to measure the frequency of CD4+ T-cells that respond to stimulation with a pool of overlapping, synthetic peptides representing the HIV-1 Gag polypeptide. Subjects in the upper half of vaccine responders, based on measuring the frequency of Gag-specific CD4 T cells are enrolled in the gene therapy arm and subjects in the lower half of responders do not continue in the study. We anticipate that the cut-off for higher responders is a HIV-specific CD4+ T cell frequency $\geq 0.065\%$ of total CD4 T cells. Subjects enrolled into the gene therapy arm of our trial undergo leukapheresis followed by purification of PBMC (using Ficoll density gradient centrifugation or negative selection with antibodies) that are cultured ex vivo and stimulated with HIV Gag peptides plus interleukin-2 and interleukin-12 for 12 days, then stimulated again with beads decorated with CD3/CD28 bispecific antibody. The antiretroviral drug Saquinavir is included at 100 nM to prevent emergence of autologous HIV during ex vivo culture. One day after CD3/CD28 stimulation cells are transduced with AGT103 at multiplicity of infection between 1 and 10. The transduced cells are cultured for an additional 7-14 days during which time they expand by polyclonal proliferation. The culture period is ended by harvesting and washing cells, setting aside aliquots for potency and safety release assays, and resuspending the remaining cells in cryopreservation medium. A single dose is $\leq 1 \times 10^{10}$ autologous PBMC. The potency assay measures the frequency of CD4 T cells that respond to peptide stimulation by expressing interferon-gamma. Other release criteria include the product must include $\geq 0.5 \times 10^7$ HIV-specific CD4 T cells that are also transduced with AGT103. Another release criterion is that the number of AGT103 genome copies per cell must not exceed 3. Five days before infusion with AGT103T subjects receive one dose of busulfan (or Cytosan) conditioning regimen followed by infusion of $\leq 1 \times 10^{10}$ PBMC containing genetically modified CD4 T cells.

[0372] A Phase II study will evaluate efficacy of AGT103T cell therapy. Phase II study participants include individuals enrolled previously in our Phase I study who were judged to have successful and stable engraftment of genetically modified, autologous, HIV-specific CD4 T cells and clinical responses defined as positive changes in parameters monitored as described in efficacy assessments (1.3.). Study participants will be asked to add Maraviroc to their existing regimen of antiretroviral medication. Maraviroc is a CCR5 antagonist that will enhance the effectiveness of genetic therapy directed at reducing CCR5 levels. Once the Maraviroc regimen is in place subjects will be asked to discontinue the previous antiretroviral drug regimen and only maintain Maraviroc monotherapy for 28 days or until plasma viral RNA levels exceed 10,000 per ml on 2 sequential weekly blood draws. Persistently high viremia requires participants to return to their original antiretroviral drug regimen with or without Maraviroc according to recommendations of their HIV care physician.

[0373] If participants remain HIV suppressed (below 2,000 vRNA copies per ml of plasma) for >28 days on Maraviroc monotherapy, they will be asked to gradually

reduce Maraviroc dosing over a period of 4 weeks followed by intensive monitoring for an additional 28 days. Subjects who maintained HIV suppression with Maraviroc monotherapy are considered to have a functional cure. Subjects who maintain HIV suppression even after Maraviroc withdrawal also have a functional cure. Monthly monitoring for 6 months followed by less intensive monitoring will establish the durability of functional cure.

[0374] Patient Selection

[0375] Inclusion Criteria:

- [0376]** Aged between 18 and 60 years.
- [0377]** Documented HIV infection prior to study entry.
- [0378]** Must be willing to comply with study-mandated evaluations; including not changing their antiretroviral regimen (unless medically indicated) during the study period.
- [0379]** CD4+ T-cell count >500 cell per millimeter cubed (cells/mm³)
- [0380]** CD4+ T-cell nadir of >400 cells/mm³
- [0381]** HIV viral load <1,000 copies per milliliter (mL)
- [0382] Exclusion Criteria:**
 - [0383]** Any viral hepatitis
 - [0384]** Acute HIV infection
 - [0385]** HIV viral load >1,000 copies/mL
 - [0386]** Active or recent (prior 6 months) AIDS defining complication
 - [0387]** Any change in HIV medications within 12 weeks of entering the study
 - [0388]** Cancer or malignancy that has not been in remission for at least 5 years with the exception of successfully treated basal cell carcinoma of the skin
 - [0389]** Current diagnosis of NYHA grade 3 or 4 congestive heart failure or uncontrolled angina or arrhythmias
 - [0390]** History of bleeding problems
 - [0391]** Use of chronic steroids in past 30 days
 - [0392]** Pregnant or breast feeding
 - [0393]** Active drug or alcohol abuse
 - [0394]** Serious illness in past 30 days
 - [0395]** Currently participating in another clinical trial or any prior gene therapy

Safety assessments

- [0396]** Acute infusion reaction
- [0397]** Post-infusion safety follow-up

Efficacy Assessments—Phase I

- [0398]** Number and frequency of modified CD4 T cells.
- [0399]** Durability of modified CD4 T cells.
- [0400]** In vitro response to Gag peptide restimulation (ICS assay) as a measure of memory T cell function.
- [0401]** Polyfunctional anti-HIV CD8 T cell responses compare to pre- and post-vaccination time points.
- [0402]** Frequency of CD4 T cells making doubly spliced HIV mRNA after in vitro stimulation.

Efficacy Assessments—Phase II

- [0403]** Number and frequency of genetically modified CD4 T cells.
- [0404]** Maintenance of viral suppression (<2,000 vRNA copies per ml but 2 consecutive weekly draws not exceeding 5×10^4 vRNA copies per ml are permitted) with Maraviroc monotherapy.

[0405] Continued virus suppression during and after Maraviroc withdrawal.

[0406] Stable CD4 T cell count.

[0407] AGT103T Consists of Up to 1×10^{10} Genetically Modified, Autologous CD4+ T Cells Containing $\geq 5 \times 10^7$ HIV-Specific CD4 T Cells that are Also Transduced with AGT103 Lentivirus Vector.

[0408] A Phase I clinical trial will test the safety and feasibility of infusing ex vivo modified autologous CD4 T cells (AGT103T) in adult research participants with confirmed HIV infection, CD4+ T-cell counts >600 cells per mm³ of blood and stable virus suppression below 200 copies per ml of plasma while on cART. Up to 40 study participants receive two doses by intramuscular injection 8 weeks apart, of recombinant modified vaccinia Ankara (rMVA) expressing HIV Gag, Pol and Env proteins. Seven to 10 days after the second immunization a blood sample is collected for in vitro testing to measure the frequency of CD4+ T-cells that respond to stimulation with a pool of overlapping, synthetic peptides representing the HIV-1 Gag polypeptide. Subjects in the upper half of vaccine responders, based on measuring the frequency of Gag-specific CD4 T cells are enrolled in the gene therapy arm and subjects in the lower half of responders do not continue in the study. We anticipate that the cut-off for higher responders is a HIV-specific CD4+ T cell frequency $\geq 0.065\%$ of total CD4 T cells. Subjects enrolled into the gene therapy arm of our trial undergo leukapheresis and the CD4+ T cells are enriched by negative selection. The enriched CD4 subset is admixed with 10% the number of cells from the CD4-negative subset to provide a source and antigen-presenting cells. The enriched CD4 T cells are stimulated with HIV Gag peptides plus interleukin-2 and interleukin-12 for 12 days, then stimulated again with beads decorated with CD3/CD28 bispecific antibody. The antiretroviral drug Saquinavir is included at 100 nM to prevent emergence of autologous HIV during ex vivo culture. One day after CD3/CD28 stimulation cells are transduced with AGT103 at multiplicity of infection between 1 and 10. The transduced cells are cultured for an additional 7-14 days during which time they expand by polyclonal proliferation. The culture period is ended by harvesting and washing cells, setting aside aliquots for potency and safety release assays, and resuspending the remaining cells in cryopreservation medium. A single dose is $\leq 1 \times 10^{10}$ autologous cells enriched for the CD4+ T cell subset. The potency assay measures the frequency of CD4 T cells that respond to peptide stimulation by expressing interferon-gamma. Other release criteria include that the product must include $\geq 0.5 \times 10^7$ HIV-specific CD4 T cells that are also transduced with AGT103. Another release criterion is that the number of AGT103 genome copies per cell must not exceed 3. Five days before infusion with AGT103T subjects receive one dose of busulfuram (or Cytosan) conditioning regimen followed by infusion of $\leq 1 \times 10^{10}$ enriched and genetically modified CD4 T cell.

[0409] A Phase II study will evaluate efficacy of AGT103T cell therapy. Phase II study participants include individuals enrolled previously in our Phase I study who were judged to have successful and stable engraftment of genetically modified, autologous, HIV-specific CD4 T cells and clinical responses defined as positive changes in parameters monitored as described in efficacy assessments (1.3.). Study participants will be asked to add Maraviroc to their existing regimen of antiretroviral medication. Maraviroc is a CCR5 antagonist that will enhance the effectiveness of

genetic therapy directed at reducing CCR5 levels. Once the Maraviroc regimen is in place subjects will be asked to discontinue the previous antiretroviral drug regimen and only maintain Maraviroc monotherapy for 28 days or until plasma viral RNA levels exceed 10,000 per ml on 2 sequential weekly blood draws. Persistently high viremia requires participants to return to their original antiretroviral drug regimen with or without Maraviroc according to recommendations of their HIV care physician.

[0410] If participants remain HIV suppressed (below 2,000 vRNA copies per ml of plasma) for >28 days on Maraviroc monotherapy, they will be asked to gradually

reduce Maraviroc dosing over a period of 4 weeks followed by intensive monitoring for an additional 28 days. Subjects who maintained HIV suppression with Maraviroc monotherapy are considered to have a functional cure. Subjects who maintain HIV suppression even after Maraviroc withdrawal also have a functional cure. Monthly monitoring for 6 months followed by less intensive monitoring will establish the durability of functional cure.

Sequences

[0411] The following sequences are referred to herein:

SEQ ID NO:	Description	Sequence
1	miR30 CCR5	AGGTATATTGCTGTTGACAGTGAGCGACTGTAACTGAGCTTGCTCTACTGTGAAGCCACAGATGGGTAGAGCAAGCACAGTTTACCGCTGCCTACTGCCTCGGACTTCAAGGGGCTT
2	miR21 Vif	CATCTCCATGGCTGTACCACCTTGTCTGGGGATGTGTACTTCTGAACCTGTGTGAATCTCATGGAGTTCAGAAGAACACATCCGCACTGACATTTTGGTATCTTTCATCTGACCA
3	miR185 Tat	GGGCTTGGCTCGAGCAGGGGGCAGGGATTCCGCTTCTTCCTGCCATAGCGTGGTCCCTTCCCTATGGCAGGCAGAGCGGCACCTTCCCTCCC AATGACCGCTCTTCGTCG
4	Elongation Factor-1 alpha (EF1-alpha) promoter	CCGGTGCCTAGAGAAGGTGGCGCGGGGTAACTGGGAAAGTGATGTCGTGACTGGCTCCGCCCTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGACAGTAGTCGCCGTGAACGTTCTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAGTGCCGTGTGTGGTTCGCCGGGCTGGCTCTTTACGGGTATGGCCCTTGCGTGCCTTGAATTACTTCCACGCCCTGGCTGCAGTACGTGATCTTGATCCCGAGCTTCGGGTGGAAAGTGGGTGGGAGAGTTCGAGGCCCTTGCCTTAAGGAGCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGCCGCTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAATTTTGTATGACCTGCTGCGACGCTTTTCTGGCAAGATAGTCTGTAAATGCGGGCC AAGATCTGCACACTGGTATTTTCGGTTTTTGGGGCCGCGGGCGGCGACGGGGCCGTGCGTCCAGCGCACATGTTCCGGCGAGGCGGGCCGTGCGAGCGCGCCACCGAGAATCGGACGGGGGTAGTCTCAAGCTGGCGCGCTGTCTGGTGCCTGGCCTCGCGCCCGGTATCGCCCCGCCCTGGGCGCAAGGCTGGCCCGGTGCGCACCAAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCGCTCGGAGAGCGGGCGGGTGAAGTCAACCCACACAAGGAA AAGGGCCTTTCGCTCCTCAGCCGTCGCTTCATGTGACTCCA CGGAGTACCGGGCGCGCTCCAGGCACCTCGATTAGTTCTCGAGCTTTGGAGTACGTCGTTTAGGTTGGGGGAGGGGTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTAATTTCTCCTTGGAA TTTGCCCTTTTGAAGTTGGATCTTGGTTCAATTCACAGCCTCAGACAGTGGTTCAAAGTTTTTCTTCCATTCAGGTGTCTG TGA
5	CCR5 target sequence	GAGCAAGCTCAGTTTACA
6	Vif target sequence	GGGATGTGTAATCTGAACTT
7	Tat target sequence	TCCGCTTCTTCCTGCCATAG
8	TAR decoy sequence	CTTGCAATGATGTCGTAATTTGCGTCTTACCTCGTTCTCGACAGCGACCAGATCTGAGCCTGGGAGCTCTCTGGCTGTCAGTAAGCTGGTACAGAAGGTTGACGAAAATTTCTACTGAGCAAGAAA
9	Rev/Tat target sequence	GCGGAGACAGCGACGAAGAGC

-continued

SEQ ID NO:	Description	Sequence
10	Rev/Tat shRNA sequence	GCGGAGACAGCGACGAAGAGCTTCAAGAGAGCTCTTCGTC GCTGTCTCCGCTTTTT
11	Gag target sequence	GAAGAAATGATGACAGCAT
12	Gag shRNA sequence	GAAGAAATGATGACAGCATTTCAGAGAATGCTGTCATCA TTTCTTCTTTTT
13	Pol target sequence	CAGGAGCAGATGATACAG
14	Pol shRNA sequence	CAGGAGATGATACAGTTCAAGAGACTGTATCATCTGCTCCT GTTTTT
15	CCR5 target sequence #1	GTGTCAAGTCCAATCTATG
16	CCR5 shRNA sequence #1	GTGTCAAGTCCAATCTATGTTCAAGAGACATAGATTGGACT TGACACTTTTT
17	CCR5 target sequence #2	GAGCATGACTGACATCTAC
18	CCR5 shRNA sequence #2	GAGCATGACTGACATCTACTTCAAGAGAGTAGATGTCAGT CATGCTCTTTTT
19	CCR5 target sequence #3	GTAGCTCTAACAGGTTGGA
20	CCR5 shRNA sequence #3	GTAGCTCTAACAGGTTGGATTCAAGAGATCCAACCTGTTAG AGCTACTTTTT
21	CCR5 target sequence #4	GTTCAGAAACTACCTCTTA
22	CCR5 shRNA sequence #4	GTTCAGAAACTACCTCTTATTCAAGAGATAAGAGGTAGTTT CTGAACTTTTT
23	CCR5 target sequence #5	GAGCAAGCTCAGTTTACACC
24	CCR5 shRNA sequence #5	GAGCAAGCTCAGTTTACACCTTCAAGAGAGGTGTAACTG AGCTTGCTCTTTTT
25	<i>Homo sapiens</i> CCR5 gene, sequence 1	ATGGATTATCAAGTGTCAAGTCCAATCTATGACATCAATTA TTATACATCGGAGCCCTGCCAAAAATCAATGTGAAGCAA ATCGCAGCCCGCTCCTGCCTCCGCTCTACTCACTGGTGT CATCTTTGGTTTGTGGGC
26	<i>Homo sapiens</i> CCR5 gene, sequence 2	AACATGCTGGTCATCCTCATCCTGATAAACTGCAAAAGGCT GAAGAGCATGACTGACATCTACCTGCTCAACCTGGCCATCT CTGACCTGTTTTTCCTTCTTACTGTCCCTTCTGGGCTCACT ATGCTGCCGCCAGTGGGACTTTGGAAATACAATGTGTCAA CTCTTGACAGGGCTCTATTTTATAGGCTTCTTCTCTGGAATC TTCTTCATCATCCTCCTGACAATCGATAGGTACCTGGCTGT CGTCCATGCTGTGTTTGCTTTAAAAGCCAGGACGGTCACCT TTGGGGTGGTGACAAGTGTGATCACTTGGGTGGTGGCTGTG TTTGCCTCTCTCCAGGAATCATCTTTACAGATCTCAAAA AGAAGGTCTTCATTACACCTGCAGCTCTCATTTTCCATACA GTCAGTATCAATTCTGGAAGAATTTCCAGACATTAAAGATA GTCATCTTGGGGCTGGTCCTGCCGCTGCTTGTGTCATGGTCAT CTGCTACTCGGGAATCCTAAAACTCTGCTTCGGTGTGCGAA ATGAGAAGAAGAGGCACAGGGCTGTGAGGCTTATCTTCAC CATCATGATTGTTTATTTCTCTTCTGGGCTCCCTACAACAT TGTCTTCTCCTGAAC
27	<i>Homo sapiens</i> CCR5 gene, sequence 3	ACCTTCCAGGAATTCTTTGGCCTGAATAATTGCAGTAGCTC TAACAGGTTGGACCAAGCTATGCAGGTGA

-continued

SEQ ID	Description	Sequence
28	<i>Homo sapiens</i> CCR5 gene, sequence 4	CAGAGACTCTTGGGATGACGCACTGCTGCATCAACCCCATC ATCTATGCCTTTGTGCGGGAGAGTTTCAGAACTACCTCTT AGTCTTCTTCCAAAAGCACATTGCCAACGCTTCTGCAAAT GCTGTCTATTTTCCAG
29	<i>Homo sapiens</i> CCR5 gene, sequence 5	CAAGAGGCTCCCGAGCGAGCAAGCTCAGTTTACACCCGAT CCACTGGGAGCAGGAAATATCTGTGGGCTTGTGA
30	CD4 promoter sequence	TGTTGGGGTTCAAATTGAGCCCCAGCTGTTAGCCCTCTGC AAAGAAAAAAAAAAAAAAAAAAGAACAAAGGGCCTAGAT TTCCCTTCTGAGCCCCACCTAAGATGAAGCCTCTTCTTTCA AGGGAGTGGGGTTGGGGTGGAGGCGGATCCTGTCACTTT GCTCTCTCTGTGGCTGGCAGTTTCTCCAAAGGTAACAGGT GTCAGCTGGCTGAGCCTAGGCTGAACCTGAGACATGCTA CCTCTGTCTTCTCATGGCTGGAGGCGAGCCTTTGTAAGTCAC AGAAAGTAGCTGAGGGGCTCTGGAAAAAGACAGCCAGG GTGGAGGTAGATTGGTCTTTGACTCCTGATTTAAGCCTGAT TCTGCTTAACCTTTTCCCTTGACTTTGGCATTTCACCTTGA CATGTTCCCTGAGAGCCTGGGGGTGGGGAACCCAGCTCC AGCTGGTGACGTTTGGGGCCGCCAGGCCTAGGGTGTGG AGGAGCCTTGCCATCGGGCTTCTGTCTCTTCAATTAAG CACGACTCTGCAGA
31	miR30- CCR5/miR21- Vif/miR185 Tat microRNA cluster sequence	AGGTATATTGCTGTTGACAGTGAGCGACTGTAACTGAGCT TGCTCTACTGTGAAGCCACAGATGGGTAGAGCAAGCACAG TTTACCGCTGCCTACTGCCTCGGACTTCAAGGGGCTTCCCG GGCATCTCCATGGCTGTACCACCTTGTGCGGGGATGTGTAC TTCTGAACTTGTGTGAATCTCATGGAGTTTCAAGAAACAC ATCCGCACTGACATTTTGGTATCTTTCATCTGACCAGCTAG CGGGCTTGGCTCGAGCAGGGGCGAGGGATTCCGCTTCTT CCTGCCATAGCGTGGTCCCTCCCTATGGCAGGCAGAAGC GGCACCTTCCCTCCCAATGACCGCTCTTCGTC
32	Long WPRE sequence	AATCAACCTCTGATTACAAAATTTGTGAAAGATTGACTGGT ATTCTTAACATGTTGCTCCTTTTACGCTATGTGGATACGCT GCTTTAATGCCTTTGTATCATGCTATTGCTTCCCGTATGGCT TTCATTTTCTCCTCTTGTATAAATCCTGGTTGCTGTCTCTTT ATGAGGAGTTGTGGCCCGTTGTGAGGCAACGTGGCGTGGT GTGCACTGTGTTTGTGACGCAACCCCACTGGTTGGGGCA TTGCCACCACTGTGAGCTCCTTTCGGGACTTTCGCTTTCC CCCTCCCTATTGCCACGGCGGAACCTCATCGCCGCTGCCTT GCCCGCTGCTGACAGGGGCTCGGCTGTTGGGCACTGACA ATTCCGTGGTGTGTGCGGGAAATCATCGTCTTTCCTTGG CTGCTCGCCTGTGTTGCCACCTGGATTCTGCGCGGGACGTC CTTCTGCTACGTCCTTTCGGCCCTCAATCAGCGGACCTTC CTTCCGCGGGCTGCTGCCGCTCTGCGGCTCTTCCGCT CTTCGCTTCCGCTCAGACGAGTGGATCTCCCTTTGGGC CGCTCCCCGCT
33	Elongation Factor-1 alpha (EF1-alpha) promoter; miR30CCR5; miR21Vif; miR185Tat	CCGGTGCCTAGAGAAGGTGGCGCGGGTAACTGGGAAAG TGATGTCGTGTAAGCTCCGCTTTTCCCGAGGGTGGGG GAGAACCGTATATAAGTGCAAGTTCGCGTGAACGTTCTT TTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAGTGCCGT GTGTGGTTCCCGCGGGCTGGCCTCTTACGGGTTATGGCC CTTGGTGCCTTGAATTACTTCCACGCCCTGGCTGCAGTA CGTGATTCTTGATCCCGAGCTTCCGGTTGGAAGTGGGTGGG AGAGTTCGAGGCTTGCCTTAAGGAGCCCTTCGCTCGT GCTTGAGTTGAGGCTGGCTGGCGCTGGGCGCCCGCG TGCGAATCTGGTGGCACCTTCGCGCTGTCTCGCTGCTTTC GATAAGTCTCTAGCCATTTAAATTTTGTGATGACCTGTGTC GACGCTTTTTTCTGGCAAGATAGTCTGTAAATGCGGGCC AAGATCTGCACACTGGTATTTGGTTTTGGGGCCCGGGC GGCGACGGGGCCGTGCGTCCAGCGCACATGTTGCGCGA GGCGGGGCTGCGAGCGCGGCCACCGAGAATCGGACGGG GTAGTCTCAAGCTGGCGGCTGCTCTGGTGCCTGGCTCG CGCCGCGTGTATCGCCCGCCCTGGGCGGCAAGGCTGGC CCGGTCGGCACAGTTGCGTGAGCGGAAGATGGCCGCTT CCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACCGGGC GCTCGGAGAGCGGGCGGTGAGTCAACACACAAAGGAA AAGGGCTTTCCGCTCTCAGCCGCTGCTTCATGTGACTCCA

-continued

SEQ ID NO: Description	Sequence
	<p>CGGAGTACCGGGCGCCGTCCAGGCACCTCGATTAGTTCTCG AGCTTTTGGAGTACGTCGTCCTTAGGTTGGGGGAGGGGTT TTATGCGATGGAGTTTCCCACACTGAGTGGGTGGAGACTG AAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGA TTTGCCCTTTTGGAGTTTGGATCTTGGTTCATTCTCAAGCCT CAGACAGTGGTTCAAAGTTTTTCTTCCATTTCAGGTGTCG TGATGTACA AGGTATATTGCTGTTGACAGTGAGCGACTGTAAACTGAGCT TGCTCTACTGTGAAGCCACAGATGGGTAGAGCAAGCACAG TTTACCGCTGCCTACTGCCTCGGACTTCAAGGGGCTTCCCG GGCATCTCCATGGCTGTACCACCTTGTGGGGGATGTGTAC TTCTGAAGTGTGTTGAATCTCATGGAGTTCAGAAGAACAC ATCCGCACTGACATTTTGGTATCTTTCATCTGACCAGCTAG CGGGCTGGCTCGAGCAGGGGCGAGGGATTCCGCTTCTT CCTGCCATAGCGTGGTCCCTCCCTATGGCAGGCAGAAGC GGCACCTTCCCTCCCAATGACCGCGTCTTCGTC</p>
34 Rous Sarcoma virus (RSV) promoter	<p>GTAGTCTTATGCAACTACTCTTGTAGTCTTGCAACATGGTAA CGATGAGTTAGCAACATGCCTTACAAGGAGAGAAAAGCA CCGTGCATGCCGATTGGTGGAAAGTAAGGTGGTACGATCGT GCCTTATTAGGAAGGCAACAGACGGGTCTGACATGGATTG GACGAACCACTGAATTGCCGATTGCAGAGATATTGTATTT AAGTGCTTAGCTCGATACAATAAACG</p>
35 5' Long terminal repeat (LTR)	<p>GGTCTCTCTGGTTAGACCAGATCTGAGCCTGGGAGCTCTCT GGCTAACTAGGGAACCCACTGCTTAAGCCTCAATAAGCTT GCCTTGAGTGCTTCAAGTAGTGTGTGCCCGTCTGTTGTGTG ACTCTGGTAACTAGAGATCCCTCAGACCTTTTAGTCAGTG TGGAATCTCTAGCA</p>
36 Psi Packaging signal	<p>TACGCCAAAAATTTGACTAGCGAGGCTAGAAGGAGAGA G</p>
37 Rev response element (RRE)	<p>AGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCA CTATGGGCGCAGCCTCAATGACGCTGACGGTACAGGCCAG ACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAATTG TGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCAC AGTCTGGGCAATCAAGCAGCTCCAGGCAGAAATCCTGGCT GTGGAAGATACCTAAAGGATCAACAGCTCC</p>
38 Central polypurine tract (cPPT)	<p>TTTTAAAGAAAAGGGGGATTGGGGGGTACAGTGCCAGGG GAAAGAATAGTAGACATAATAGCAACAGACATACAACTA AAGAATTACAAAAACAAATTACAAAATTCAAATTTTA</p>
39 3' delta LTR	<p>TGGAAGGGCTAATTCACCTCCCAACGAAGATAAGATCTGCTT TTTGCTTGACTGGGTCTCTCTGGTTAGACCAGATCTGAGC CTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCTTAAGC CTCAATAAAGCTTGCTTGGAGTGCTTCAAGTAGTGTGTGCC CGTCTGTTGTGTGACTCTGGTAACTAGAGATCCCTCAGACC CTTTTAGTCAGTGTGGAATCTCTAGCAGTAGTAGTTTCAT GTCA</p>
40 Helper/Rev; CMV early (CAG) enhancer; Enhance Transcription	<p>TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATA GCCCCATATATGGAGTTCCGCGTTACATAACTTACGGTAAAT GGCCCCGCTGGCTGACCGCCCAACGACCCCCGCCATTGAC GTCAATAATGACGTATGTTCCATAGTAACGCCAATAGGG ACTTTCCATTGACGTCAATGGGTGGACTATTACGGTAAAC TGCCCACTTGGCAGTACATCAAGGTATCATATGCCAAGTA CGCCCCCTATTGACGTCAATGACGTAATGGCCCGGCTGG CATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTG GCAGTACATCTACGTATTAGTCATC</p>
41 Helper/Rev; Chicken beta actin (CAG) promoter; Transcription	<p>GCTATTACCATGGGTCGAGGTGAGCCCCACGTTCTGCTTCA CTCTCCCCATCTCCCCCCTCCCCACCCCAATTTGTATT TATTTATTTTAAATATTTTGTGCGAGGATGGGGCGGGG GGGGGGGGGCGCGCGCCAGGCGGGGCGGGGCGGGGCGA GGGGCGGGGCGGGGCGAGGCGGAGAGGTGCGGCGGCAGC CAATCAGAGCGGCGCGCTCCGAAAGTTTCTTTATGGCGA GGCGGCGGGCGGGCGGCCCTATAAAAAGCGAAGCGCGCG GCGGGCG</p>

-continued

SEQ ID	NO: Description	Sequence
42	Helper/Rev; Chicken beta actin intron; Enhance gene expression	GGAGTCGCTGCGTTGCCTTCGCCCCGTGCCCCGCTCCGCGC CGCCTCGCGCCGCCCGCCCGGCTCTGACTGACCGCGTTAC TCCACAGGTGAGCGGGCGGACGCGCCCTTCTCTCCGGG CTGTAATTAGCGCTTGGTTTAATGACGGCTCGTTTCTTTTCT GTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCCCT TTGTGCGGGGGGAGCGGCTCGGGGGTGCCTGCGTGTGT GTGTGCGTGGGAGCGCCGCTGCGGCCCGCGCTGCCCGG CGGCTGTGAGCGCTGCGGGCGCGGCGCGGGGCTTTGTGCG CTCCGCGTGTGCGGAGGGGAGCGCGGCCGGGGCGGTGC CCCCGCGTGGGGGGGCTGCGAGGGGAACAAAGGCTGCG TGCGGGGTGTGTGCTGGGGGGTGTGAGAGGGGTGTGGG CGCGCGGTGCGGCTGTAAACCCCCCTGCACCCCCCTCC CGAGTTGCTGAGCAGGCCCGGCTTCGGGTGCGGGGCTCC GTGCGGGGCTGGCGCGGGGCTCGCCGTGCCGGCGGGG GTGCGGCGAGGTGGGGTGC CGGGCGGGCGGGCGCGCT CGGGCGGGGAGGGCTCGGGGAGGGGCGCGCGGCCCGG GGAGCGCGGGCGGCTGTGAGGCGCGCGAGCCGAGCCA TTGCCCTTTATGGTAATCGTGCAGAGGGCGCAGGACTTC CTTTGTCCCAATCTGGCGGAGCCGAAATCTGGGAGCGC CGCCGCGCCCCCTCTAGCGGGCGCGGGCGAAGCGGTGCGG CGCCGCGAGGAAGGAATGGCGGGGAGGGCTTCGTGCG TCGCCGCGCGCGCTCCCTTCTCCATCTCCAGCCTCGGG CTGCCGAGGGGACGGCTGCCTTCGGGGGACGGGGCA GGCGGGGTTCGGCTTCTGCGTGTGACCGGCGG
43	Helper/Rev; HIV Gag; Viral capsid	ATGGGTGCGAGAGCGTCAGTATTAAGCGGGGAGAATTAG ATCGATGGGAAAAAATTCGGTTAAGCGCAGGGGAAAGAA AAAATATAAATTAAAAATATAGTATGGGCAAGCAGGGAG CTAGAACGATTGCGAGTTAATCCTGGCCTGTGTAGAAACATC AGAAGGCTGTAGACAAATACTGGGACAGCTACAACCATCC CTTCAGACAGGATCAGAAGAACTTAGATCATTATATAATAC AGTAGCAACCCCTCTATTGTGTGCATCAAGGATAGAGATA AAAGACACCAAGGAAGCTTTAGACAAGATAGAGGAAGAG CAAAACAAAAGTAAGAAAAAGCACAGCAAGCAGCAGCT GACACAGGACACAGCAATCAGGTGAGCAAAATTACCCTA TAGTGCAGAACATCCAGGGGCAATGGTACATCAGGCCAT ATCACCTAGAACTTTAAATGCATGGGTAAAAGTAGTAGAA GAGAAGGCTTTAGCCCAAGGTGATACCATGTTTTCAGC ATTATCAGAGGAGCCACCCACAAGATTAAACACCATG CTAAACACAGTGGGGGACATCAAGCAGCCATGCAATGT TAAAGAGAGCCATCAATAGGAAGCTGCAGAAATGGGATAG AGTGCATCCAGTGCATGCAAGGCTATTGCACCAAGCCAG ATGAGAGAACCAAGGGAGTGACATAGCAGGAACACTA GTACCTCTCAGGAACAAATAGGATGGATGACACATAATCC ACCTATCCAGTAGGAGAAATCTATAAAGATGGATAATC CTGGGATTAAATAAAATAGTAAGAAATGTATAGCCCTACCA GCATTCTGGACATAAGACAAGGACCAAGGAACCCCTTAG AGACTATGTAGACCGATTCTATAAACTCTAAGAGCCGAG CAAGCTTCACAAGAGGTAAAAAATGGATGACAGAAACCT TGTGGTCCAAAATGCGAACCCAGATTGTAAGACTATTTA AAAGCATTGGGACCAGGAGCGACACTAGAAGAAATGATGA CAGCATGTGAGGAGTGGGGGACCCGGCCATAAAGCAAG AGTTTTGGCTGAAGCAATGAGCCAAGTAACAAATCCAGCT ACCATAATGATACAGAAAGCAATTTAGGAACCAAGAA AGACTGTTAAGTGTTCATTTGTGGCAAGAAGGGCACAT AGCCAAAAATTCAGGGGCCCCAGGAAAAAGGGCTGTTGG AAATGTGGAAGGAAGGACACCAATGAAAGATTGTACTG AGAGACAGGCTAATTTTTAGGGAAGATCTGGCCTTCCAC AAGGGAAGGCCAGGAATTTCTTCAGAGCAGACCAAGC CAACAGCCCCACCAGAAGAGCTTCAGTTTGGGGAAGA GACAACAACCTCCCTCTCAGAAGCAGGAGCCGATAGACAAG GAAGTGTATCTTTAGCTTCCCTCAGATCACTCTTTGGCAG CGACCCCTCGTCACAATAA
44	Helper/Rev; HIV Pol; Protease and reverse transcriptase	ATGAATTTGCCAGGAAGATGGAAACCAAAATGATAGGGG GAATTGGAGGTTTATCAAAGTAGGACAGTATGATCAGAT ACTCATAGAAATCTGCGGACATAAAGCTATAGGTACAGTA TTAGTAGGACCTACACCTGTCAACATAATTGGAAGAAATCT GTTGACTCAGATTGGCTGCACTTTAAATTTTCCATTAGTCC TATTGAGACTGTACCAGTAAATTTAAAGCCAGGAATGGAT GGCCCAAAAGTTAAACAAATGGCCATTGACAGAAAGAAAA TAAAGCATTAGTAGAAATTTGTACAGAAATGGAAGGA AGGAAAAATTTCAAAATTTGGCCTGAAATCCATACAAT

-continued

SEQ ID	NO: Description	Sequence
		ACTCCAGTATTTGCCATAAAGAAAAAGACAGTACTAAAT GGAGAAAAATAGTAGATTTCAGAGAACTTAATAAGAGAAC TCAAGATTTCTGGGAAGTTCAATTAGGAATACCACATCCTG CAGGGTTAAAAACAGAAAAATCAGTAACAGTACTGGATGT GGGCGATGCATATTTTTCAGTTCCTTAGATAAAGACTTCA GGAAGTATACTGCATTTACCATACCTAGTATAACAATGAG ACACCAGGGATTAGATATCAGTACAATGTGCTTCCACAGG GATGGAAGGATCACCAGCAATATTCCAGTGTAGCATGAC AAAAACTTAGAGCCTTTTAGAAAACAAATCCAGACATA GTCATCTATCAATACATGGATGATTGTATGTAGGATCTGA CTTAGAAATAGGGCAGCATAGAACAAAAATAGAGGAACGT AGACAACTCTGTTGAGGTGGGATTACCACACCAGACA AAAAAATCAGAAAGAACCTCCATTCTTTGGATGGGTTAT GAACTCCATCCTGATAAATGGACAGTACAGCCTATAGTGCT GCCAGAAAAGGACAGCTGGACTGTCAATGACATACAGAAA TTAGTGGGAAAATTGAATTGGGCAAGTCAGATTTATGCAG GGATTAAGTAAGGCAATTATGTAACTTCTTAGGGGAAC CAAAGCACTAACAGAAGTAGTACCCTAACAGAAAGCA GAGCTAGAAGTGGCAGAAAACAGGGAGATTCTAAAAGAAC CGGTACATGGAGTGTATTATGACCCATCAAAGACTTAATA GCAGAAATACAGAAGCAGGGGCAAGGCCAATGGACATATC AAATTTATCAAGAGCCATTTAAAAATCTGAAAACAGGAAA ATATGCAAGAAATGAAGGGTGCCACACTAATGATGTGAAA CAATTAACAGAGGCAGTACAAAAATAGCCACAGAAAGCA TAGTAATATGGGAAAGACTCCTAAATTTAAATTACCCATA CAAAAGGAAACATGGGAAGCATGGTGGACAGAGTATTGGC AAGCCACCTGGATTCTTGAGTGGGAGTTGTCAATACCCCT CCCTTAGTGAAGTTATGGTACCAGTTAGAGAAAGAACCCA TAATAGGAGCAGAACTTCTATGTAGATGGGGCAGCCAA TAGGGAACTAAATTAGGAAAAGCAGGATATGTAATGAC AGAGGAAGACAAAAAGTTGTCCCCCTAACGGACACAACAA ATCAGAAGACTGAGTTACAAGCAATTCATCTAGCTTTGCAG GATTCGGGATTAGAAGTAAACATAGTGACAGACTCACAT ATGCATTGGGAATCATTCAAGCACAAACAGATAAGAGTGA ATCAGAGTTAGTCAGTCAAAATATAGAGCAGTTAATAAAA AAGGAAAAAGTCTACTTGGCATGGGTACCAGCACACAAG GAATTGGAGGAAATGAACAAGTAGATGGGTTGGTCAGTGC TGAATCAGGAAAGTACTA
45	Helper Rev; HIV Integrase; Integration of viral RNA	TTTTATAGATGGAATAGATAAGGCCCAAGAAGAACATGAGA AATATCACAGTAATTGGAGAGCAATGGCTAGTGATTTTAACT CTACCACCTGTAGTAGCAAAAGAAATAGTAGCCAGCTGTG ATAAATGTCAGCTAAAAGGGGAAGCCATGCATGGACAAGT AGACTGTAGCCAGGAATATGGCAGCTAGATTGTACACAT TTAGAAGGAAAAGTTATCTTGGTAGCAGTTTCATGTAGCCAG TGGATATATAGAAGCAGAAGTAATCCAGCAGAGACAGGG CAAGAAACAGCATACTTCTCTTAAATTAGCAGGAAGAT GGCCAGTAAAAACAGTACATACAGACAATGGCAGCAATTT CACCAGTACTACAGTTAAGGCCGCCCTGTTGGTGGGCGGGG ATCAAGCAGGAATTTGGCATTCCTTACAATCCCCAAAGTCA AGGAGTAATAGAATCTATGAATAAGAATTAAAGAAAAAT ATAGGACAGGTAAGAGATCAGGCTGAACATCTTAAGACAG CAGTACAAATGGCAGTATTCATCCCAATTTTAAAGAAAA AGGGGGGATTGGGGGTACAGTGCAGGGGAAAGAAATAGT AGACATAATAGCAACAGACATACAACTAAAGAAATTACAA AAACAAATTACAAAATTCAAAATTTTCGGGTTTATTACAG GGACAGCAGAGATCCAGTTTGGAAAGGACAGCAAGGCTC CTCTGGAAAGGTGAAGGGGAGTAGTAATACAAGATAATA GTGACATAAAAGTAGTGCCAAAGAAAGCAAGATCAT CAGGGATTATGGAACAGATGGCAGGTGATGTTGTGTG GCAAGTAGACAGGATGAGGATTAA
46	Helper/Rev; HIV RRE; Binds Rev element	AGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCA CTATGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAG ACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAATTGTC TGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCAC AGTCTGGGGCATCAAGCAGCTCCAGGCAAGAATCCTGGCT GTGGAAGATACCTAAGGATCAACAGCTCCT
47	Helper/Rev; HIV Rev; Nuclear	ATGGCAGGAAGAAGCGGAGACAGCGACGAAGAACTCCTC AAGGCAGTCAGACTCATCAAGTTTCTCTATCAAGCAACCC ACCTCCCAATCCCGAGGGGACCCGACAGGCCGGAAGGAAT

-continued

SEQ ID	NO: Description	Sequence
	export and stabilize viral mRNA	AGAAGAAGAAGGTGGAGAGAGAGACAGAGACAGATCCAT TCGATTAGTGAAACGGATCCTTAGCACTTATCTGGGACGATC TGCGGAGCCTGTGCCTCTTACGTACCACCGCTTGAGAGAC TTACTCTTGATTGTAACGAGGATTGTGGAACCTCTGGGACG CAGGGGGTGGGAAGCCCTCAAATATTGGTGAATCTCCTA CAATATTGGAGTCAGGAGCTAAGAATAG
48	Helper/Rev; Rabbit beta globin poly A; RNA stability	AGATCTTTTCCCTCTGCCAAAAATTATGGGGACATCATGA AGCCCTTGAGCATCTGACTTCTGGCTAATAAGGAAATTT ATTTTCATTGCAATAGTGTGTTGGAATTTTTGTGTCTCTCA CTCGGAAGGACATATGGGAGGGCAATCATTTAAACATC AGAATGAGTATTGGTTTAGAGTTTGGCAACATATGCCATA TGCTGGCTGCCATGAACAAAGGTGGCTATAAGAGGTCAT CAGTATATGAAACAGCCCCCTGCTGTCCATTCTTATTCCA TAGAAAAGCCTTGACTTGAGGTAGATTTTTTTTATATTTG TTTTGTGTATTTTTTCTTTAACATCCCTAAAATTTTCTTAA CATGTTTACTAGCCAGATTTTCTCTCTCTCTGACTACTC CCAGTCATAGCTGTCCCTCTTCTTATGAAGATC
49	Helper; CMV early (CAG) enhancer; Enhance transcription	TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATA GCCCCATATGGAGTTCGCGTTACATAACTTACGGTAAAT GGCCCCGCTGGCTGACCGCCCAACGACCCCGCCCATGAC GTCAATAATGACGTATGTTCCATAGTAACGCCAATAGGG ACTTTCCATTGACGTCAATGGGTGAGCTATTTACGTTAAAC TGCCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTA CGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCTGG CATTATGCCAGTACATGACCTTATGGGACTTCTTACTTGG GCAGTACATCTACGTATTAGTCATC
50	Helper; Chicken beta actin (CAG) promoter; Transcription	GCTATTACCATGGTCGAGGTGAGCCCCACGTTCTGCTTCA CTCTCCCCATCTCCCCCCCCCTCCCCACCCCAATTTTGATT TATTTATTTTTTAATATTTTGTGACGATGGGGCGGGG GGGGGGGGGCGCGCGCCAGGCGGGCGGGCGGGGCGA GGGGCGGGCGGGCGAGGCGGAGAGGTGCGGCGGCAGC CAATCAGAGCGGCGGCTCCGAAAGTTTCTTTTATGGCGA GGCGGCGGCGGCGGCGCCCTATAAAAGCGAAGCGCGCG GCGGGCG
51	Helper; Chicken beta actin intron; Enhance gene expression	GGAGTCGCTGCGTTGCCTTCGCCCGTGCCCGCTCCGCGC CGCCTCGCGCCCGCCCGCCCGCTCTGACTGACCGCGTTAC TCCCACAGGTGAGCGGGCGGACGGCCCTTCTCCTCCGGG CTGTAATTAGCGCTTGGTTTAATGACGGCTCGTTTCTTTCT GTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCCCT TTGTGCGGGGGGAGCGGCTCGGGGGTGCCTGCTGTGT GTGTGCGTGGGAGCGCCGCTGCGGCCCGCGCTGCCCG CGGCTGTGAGCGCTGCGGGCGCGGCGGGGCTTTGTGCG CTCCGCGTGTGCGGAGGGGAGCGCGCGGGGGCGGTGC CCCGCGTGTGCGGGGGGCTGCGAGGGGAACAAGGCTGCG TGCGGGGTGTGTGCTGGGGGGTGAGCAGGGGTGTGGG CGCGCGGTGCGGGCTGTAAACCCCCCTGCACCCCCCTCC CGAGTTGCTGAGCACGGCCCGGCTTCGGGTGCGGGGCTCC GTGCGGGGCTGGCGCGGGGCTCGCCGTGCCGGGCGGGG GTGGCGGCAAGTGGGGGTGCGGGCGGGGCGGGGCGCCT CGGGCGGGGAGGGCTCGGGGAGGGGCGCGGCGGCCCC GGAGCGCCGGCGGCTGTGAGGCGCGGCGAGCCGAGCCA TTGCCCTTTATGGTAATCGTGCAGAGGGCGCAGGGACTTC CTTTGTCCAAATCTGGCGAGCCGAAATCTGGGAGCGC CGCCGCACCCCTCTAGCGGGCGCGGCGAAGCGGTGCGG CGCGGCGAGGAAGGAAATGGGCGGGGAGGGCTTCGTGCG TCGCGCGCGCGCGTCCCCTTCTCCATCTCCAGCCTCGGG CTGCCGCGAGGGGACGGCTGCCTTCGGGGGGACGGGGCA GGGCGGGGTTGCGCTTCTGGCGTGTGACCGGCGG
52	Helper; HIV Gag; Viral capsid	ATGGGTGCGAGAGCGTCAGTATTAAGCGGGGAGAATTAG ATCGATGGGAAAAATTCGGTTAAGGCCAGGGGGAAGAA AAAAATATAAATAAAACATATAGTATGGGCAAGCAGGGAG CTAGAACGATTGCGAGTTAATCTGGCTGTTAGAAACATC AGAAGGCTGTAGACAAATACTGGGACAGCTACAACCATCC CTTCAGACAGGATCAGAAGAACTTAGATCATTATATAATAC AGTAGCAACCCCTCTATTGTGTGCATCAAAGGATAGAGATA AAAGACACCAAGGAAGCTTAGACAAGATAGAGGAAGAG CAAAACAAAAGTAAGAAAAAGCACAGCAAGCAGCAGCT

-continued

SEQ ID NO: Description	Sequence
	GACACAGGACACAGCAATCAGGTCAGCCAAAATTACCCTA TAGTGACAGAACATCCAGGGGCAAATGGTACATCAGGCCAT ATCACCTAGAACTTTAAATGCATGGGTAAAAGTAGTAGAA GAGAAGGCTTTAGCCAGAGTGATACCCATGTTTTCAGC ATTATCAGAAGGAGCCACCCACAAGATTAAACACCATG CTAAACACAGTGGGGGACATCAAGCAGCCATGCAATGT TAAAAGAGACCATCAATGAGGAAGCTGCAGAATGGGATAG AGTGCATCCAGTGCATGCAGGGCCTATTGCACCAGGCCAG ATGAGAGAACCAGGGGAAGTGACATAGCAGGAACACTA GTACCCCTTCAGGAACAAATAGGATGGATGACACATAATCC ACCTATCCAGTAGGAGAAATCTATAAAGATGGATAATC CTGGGATTAAATAAAATAGTAAGAATGTATAGCCCTACCA GCATTCTGGACATAAGACAAGGACCAAGGAACCCCTTAG AGACTATGTAGACCGATTCTATAAACTCTAAGAGCCGAG CAAGCTTCACAAGAGGTAAAAAATTGGATGACAGAAACCT TGTTGGTCCAAAATGCGAACCCAGATTGTAAGACTATTTTA AAAGCATTGGGACCGAGGACGACATAGAAGAAATGATGA CAGCATGTCAGGGAGTGGGGGACCCGGCCATAAAGCAAG AGTTTTGGCTGAAGCAATGAGCCAAGTAACAAATCCAGCT ACCATAAATGATACAGAAAGGCAATTTAGGAACCAAGAA AGACTGTTAAGTGTTCATTTGTGCCAAGAAGGGCAGAT AGCCAAAAATTGCAGGGCCCCAGGAAAAGGGCTGTTGG AAATGTGGAAGGAAGGACACCAATGAAGATTGTACTG AGAGACAGGCTAATTTTTAGGGAAGATCTGGCCTTCCAC AAGGAAGGCCAGGGAATTTCTTCAGAGCAGACCAGAGC CAACAGCCCCACAGAGAGAGCTTCAGGTTTGGGGAAGA GACAACTACTCCCTCTCAGAGCAGGAGCCGATAGACAAG GAACTGTATCCTTTAGCTTCCCTCAGATCACTCTTTGGCAG CGACCCCTCGTCACAATAA
53 Helper; HIV Pol; Protease and reverse transcriptase	ATGAATTTGCCAGGAAGATGGAAACCAAAATGATAGGGG GAATTTGGAGGTTTATCAAAGTAGGACAGTATGATCAGAT ACTCATAGAAATCTGCGGACATAAAGCTATAGGTACAGTA TTAGTAGGACCTACACCTGTCAACATAATGGAAGAAATCT GTTGACTCAGATGGCTGCACCTTAAATTTCCATTAGTCC TATTGAGACTGTACCAGTAAAATTAAAGCCAGGAATGGAT GGCCCAAAAGTTAAACAATGGCCATTGACAGAAGAAAAAA TAAAGCATTAGTAGAAATTTGTACAGAAATGGAAGGA AGGAAAAATTTCAAAATTTGGCCTGAAAATCCATACAT ACTCCAGTATTTGCCATAAAGAAAAAGACAGTACTAAAT GGAGAAAAATTAGTAGATTTCAGAGAACTTAATAAGAGAAC TCAAGATTTCTGGGAAGTTCAATTAGGAATACCACATCCTG CAGGGTTAAACAGAAAAATCAGTAACAGTACTGGATGT GGGCGATGCATATTTTCAGTTCCCTTAGATAAAGACTTCA GGAAGTATACTGCATTTACCATACCTAGTATAAACAATGAG ACACCAGGATTAGATATCAGTACAAATGTGCTTCCACAGG GATGGAAGGATCACCAGCAATTTCCAGTGTAGCATGAC AAAAATCTTAGAGCCCTTTAGAAAACAAAATCCAGACATA GTCATCTATCAATACATGGATGATTGTATGTAGGATCTGA CTTAGAAATAGGGCAGCATAGAACAAAAATAGAGGAACGT AGACAACATCTGTTGAGGTGGGGATTACCACACCAGACA AAAAACATCAGAAAGAACCTCCATTCCCTTTGGATGGGTTAT GAACTCCATCCTGATAAATGGACAGTACAGCCTATAGTGCT GCCAGAAAAGGACAGCTGGACTGTCAATGACATACAGAAA TTAGTGGGAAAATTGAATTGGGCAAGTCAGATTTATGCAG GGATTAAAGTAAGGCAATTATGTAACCTCTTAGGGGAAC CAAAGCACTAACAGAAAGTAGTACCCTAACAGAAAGCA GAGCTAGAACTGGCAGAAAAAGGGAGATTCTAAAAGAAC CGGTACATGGAGTGTATTATGACCCATCAAAAGACTTAATA GCAGAAATACAGAAAGGGGCAAGGCCAATGGACATATC AAATTTATCAAGAGCCATTTAAAAATCTGAAAAAGGAAA ATATGCAAGAATGAAGGGTGCCACACTAATGATGTGAAA CAATTAACAGAGGCAGTACAAAAAATAGCCACAGAAAGCA TAGTAATATGGGGAAGACTCCTAAATTTAAATTACCCATA CAAAAGGAACATGGGAAGCATGGTGGACAGAGTATTGGC AAGCCACCTGGATTCTGAGTGGGAGTTGTCAATACCCCT CCCTTAGTGAAGTTATGGTACCAGTTAGAGAAAGAACCA TAATAGGAGCAGAACTTTCTATGTAGATGGGGCAGCCAA TAGGGAAACTAAATTAGGAAAAGCAGGATATGTAAGTAC AGAGGAAGACAAAAAGTTGTCCCCCTAACGGACACAACAA ATCAGAAGACTGAGTTACAAGCAATTCATCTAGCTTGCAG GATTTCGGGATTAGAAGTAAACATAGTGACAGACTCACAAT

-continued

SEQ ID NO: Description	Sequence
	ATGCATTGGGAATCATTCAAGCACACCAGATAAGAGTGA ATCAGAGTTAGTCAGTCAAATAATAGAGCAGTTAATAAAA AAGGAAAAAGTCTACCTGGCATGGGTACCAGCACACAAAG GAATTGGAGGAAATGAACAAGTAGATGGGTTGGTCAGTGC TGAATCAGGAAAGTACTA
54 Helper; HIV Integrase; Integration of viral RNA	TTTTTAGATGGAATAGATAAGGCCCAAGAAGAACATGAGA AATATCACAGTAATTGGAGAGCAATGGCTAGTGATTTTAAC CTACCACCTGTAGTAGCAAAAGAAATAGTAGCCAGCTGTG ATAAATGTCAGCTAAAAGGGGAAGCCATGCATGGACAAGT AGACTGTAGCCAGGAATATGGCAGCTAGATTGTACACAT TTAGAAGGAAAAGTTATCTTGGTAGCAGTTTCATGTAGCCAG TGGATATATAGAAGCAGAAGTAATCCAGCAGAGACAGGG CAAGAAACAGCATACTTCTCTTAAATTAGCAGGAAGAT GGCCAGTAAAAACAGTACATACAGACAATGGCAGCAATTT CACCAGTACTACAGTTAAGGCCGCCTGTTGGTGGGCGGG ATCAAGCAGGAATTTGGCATTCCCTACAATCCCAGAGTCA AGGAGTAATAGAATCTATGAATAAGAATTAAAGAAAATT ATAGGACAGGTAAGAGATCAGGCTGAACATCTTAAGACAG CAGTACAAATGGCAGTATTCATCCCAATTTTAAAGAAA AGGGGGATTGGGGGTAAGTGCAGGGGAAAGAAATAGT AGACATAATAGCAACAGACATACAACTAAAGAATTACAA AAACAAATTACAAAATTCAAATTTTCGGGTTTATTACAG GGACAGCAGAGATCCAGTTTGGAAAGGACCAGCAAGCTC CTCTGGAAGGTGAAGGGGCAGTAGTAATACAAGATAATA GTGACATAAAAGTAGTGCCAAAGAAAGCAAGATCAT CAGGGATTATGGAAACAGATGGCAGGTGATGTTGTGTG GCAAGTAGCAGGATGAGGATTAA
55 Helper; HIV RRE; Binds Rev element	AGGAGCTTGTTCCTTGGGTCTTGGGAGCAGCAGGAAGCA CTATGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAG ACAATTATTTGTCTGGTATAGTGACAGCAGACAATTTGCT TGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCAC AGTCTGGGCATCAAGCAGCTCCAGGCAAGAACTCTGGCT GTGGAAGATACCTAAAGGATCAACAGCTCCT
56 Helper; Rabbit beta globin poly A; RNA stability	AGATCTTTTCCCTCTGCCAAAAATTATGGGGACATCATGA AGCCCCCTTGAGCATCTGACTTCTGGCTAATAAGGAAATTT ATTTTCATTGCAATAGTGTGTTGGAATTTTTGTGTCTCTCA CTCGGAAGGACATATGGGAGGGCAAATCATTTAAACATC AGAATGAGTATTTGGTTTAGAGTTTGGCAACATATGCCATA TGCTGGCTGCCATGAACAAAGGTGGCTATAAAGAGGTGAT CAGTATATGAAACAGCCCCCTGCTGTCCATTCTTATTCCA TAGAAAAGCCTTGACTTGAGGTTAGATTTTTTTTATATTTG TTTTGTGTTATTTTTTTCTTTAACATCCCTAAAATTTCTCTTA CATGTTTTTACTAGCCAGATTTTCTCTCTCTCTGACTACTC CCAGTCATAGCTGTCCCTCTTCTCTTATGAAGATC
57 Rev; RSV promoter; Transcription	ATGGCAGGAAGAAGCGGAGACAGCGACGAAGAACTCCTC AAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC ACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAAGGAAT AGAAGAAGAAGGTGGAGAGAGACAGAGACAGATCCAT TCGATTAGTGAACGGATCCTTAGCACTTATCTGGGACGATC TGCGGAGCCTGTGCCTCTTCAGCTACCACCGCTTGAGAGAC TTACTCTTGATTGTAAACGAGGATTGTGGAACCTCTGGGACG CAGGGGGTGGGAAGCCCTCAAATATTGGTGGAATCTCCTA CAATATTGGAGTCAGGAGCTAAAGAATAG
58 Rev; HIV Rev; Nuclear export and stabilize viral mRNA	ATGGCAGGAAGAAGCGGAGACAGCGACGAAGAACTCCTC AAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC ACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAAGGAAT AGAAGAAGAAGGTGGAGAGAGAGACAGAGACAGATCCAT TCGATTAGTGAACGGATCCTTAGCACTTATCTGGGACGATC TGCGGAGCCTGTGCCTCTTCAGCTACCACCGCTTGAGAGAC TTACTCTTGATTGTAAACGAGGATTGTGGAACCTCTGGGACG CAGGGGGTGGGAAGCCCTCAAATATTGGTGGAATCTCCTA CAATATTGGAGTCAGGAGCTAAAGAATAG
59 Rev; Rabbit beta globin poly A; RNA stability	AGATCTTTTCCCTCTGCCAAAAATTATGGGGACATCATGA AGCCCCCTTGAGCATCTGACTTCTGGCTAATAAGGAAATTT ATTTTCATTGCAATAGTGTGTTGGAATTTTTGTGTCTCTCA CTCGGAAGGACATATGGGAGGGCAAATCATTTAAACATC

-continued

SEQ ID NO: Description	Sequence
	AGAATGAGTATTTGGTTTAGAGTTTGGCAACATATGCCCAT ATGCTGGCTGCCATGAACAAAGGTTGGCTATAAAGAGGTC ATCAGTATATGAAACAGCCCCCTGCTGCCATTCTTATTC CATAGAAAAGCCTTGACTTGAGGTTAGATTTTTTTTATATTT TGTTTTGTGTTATTTTTTCTTTAACATCCCTAAATTTTCCT TACATGTTTACTAGCCAGATTTTCTCTCTCTCTGACTAC TCCCAGTCATAGCTGTCCCTCTTCTCTTATGGAGATC
60 Envelope; CMV promoter; Transcription	ACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGG GGTCATTAGTTCATAGCCCATATATGGAGTTCGCGGTTACA TAACTTACGGTAAATGGCCCGCTGGCTGACGCCCAACG ACCCCGCCCATTGACGTCAATAATGACGTATGTTCCCAT GTAACGCCAATAGGGACTTTCATTGACGTCAATGGGTGG AGTATTTACGGTAACTGCCCACTTGGCAGTACATCAAGTG TATCATATGCCAAGTACGCCCTTATTGACGTCAATGACGG TAAATGGCCCGCTGGCATTATGCCCAGTACATGACCTTAT GGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATC GCTATTACCATGGTGATGCGGTTTGGCAGTACATCAATGG GCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCC ACCCCATTGACGTCAATGGGAGTTTGTGTTGGCACCAAAAT CAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCAT GACGCAAAATGGGCGGTAGCGGTGTACGGTGGGAGGTCTAT ATAAGC
61 Envelope; Beta globin intron; Enhance gene expression	GTGAGTTTGGGGACCCCTTGATTGTTCTTTCTTTTCGCTATT GTAAATTCATGTTATATGGAGGGGGCAAAGTTTTCAGGGT GTTGTTAGAATGGGAAGATGTCCTTGTATCACCATTGGAC CCTCATGATAATTTTGTTCTTCTACTTCTACTCTGTGAC AACCATTGTCTCCTCTTATTTCTTTTCAATTTCTGTAACTTT TTCGTTAACTTTAGCTTGCAATTTGTAACGAATTTTAAATT CACTTTTGTTTATTTGTCAGATTGTAAGTACTTCTCTAATC ACTTTTTTTTCAAGGCAATCAGGGTATATTATTTGACTTTC AGCACAGTTTATAGAGAACAATTGTTATAATTAAATGATAAG GTAGAAATATTTCTGCATATAAATCTGGCTGGCGTGAAAT ATTCTTATTTGGTAGAAACAATACACCCCTGGTCATCATCT GCCTTCTCTTTATGTTTACAATGATATACACTGTTTGAGAT GAGGATAAAATACTCTGAGTCCAAACCGGCCCTCTGCT AACCATGTTTCATGCTTCTTCTCTTCTCTACAG
62 Envelope; VSV-G; Glycoprotein envelope-cell entry	ATGAAGTGCCTTTTGTACTTAGCCTTTTATTCATTGGGGTG AATTGCAAGTTCACCATAGTTTTCACACAACCAAAAGG AAACTGGAAAAATGTTCTTCTAATTACCATTATTGCCCGT CAAGCTCAGATTTAAATTGGCATAATGACTTAATAGGCACA GCCTTACAAGTCAAATGCCAAGAGTCAAGAGCTATTTC AAGCAGACGGTTGGATGTGTATGCTTCCAAATGGGTCACT ACTTGTGATTTCCGCTGGTATGGACCGAAGTATATAACACA TTCCATCCGATCCTTCACTCCATCTGTAGAACAATGCAAGG AAAGCATTGAACAAACGAAACAAAGAACTTGGCTGAATCC AGGCTTCCCTCCTCAAAGTTGTGGATATGCAACTGTGACGG ATGCCGAAGCAGTGATTGTCCAGGTGACTCCTCACCATGTG CTGGTTGATGAATACACAGGAGAATGGGTTGATTACAGTT CATCAACGGAATGCAGCAATTACATATGCCCCACTGTCC ATAACTCTACAACCTGGCATTTGACTATAAGGTCAAAGGG CTATGTGATTCTAACCTCATTTCCATGGACATCACCTTCTTC TCAGAGGACGGAGAGCTATCATCCCTGGGAAAGGAGGGCA CAGGGTTGAGAAGTAACACTTTGCTTATGAAACTGGAGGC AAGGCCTGCAAAATGCAATACTGCAAGCATTGGGGAGTCA GACTCCCATCAGGTGTCTGGTTCGAGATGGCTGATAAGGAT CTCTTTGCTGCAGCCAGATTCCCTGAATGCCAGAGGGTC AAGTATCTCTGCTCCATCTCAGACCTCAGTGGATGTAAGTC TAATTCAGGACGTTGAGAGGATCTTGGATTATTCCTCTGC CAAGAAACCTGGAGCAAAATCAGAGCGGGTCTTCCAATCT CTCCAGTGGATCTCAGCTATCTTGCTCCTAAAAACCCAGGA ACCGGTCCTGCTTTCACCATAAATCAATGGTACCCTAAATA CTTTGAGACCAGATACATCAGAGTCGATATTGCTGCTCCAA TCCTCTCAAGATGGTCGGAATGATCAGTGGAACTACCAC AGAAAGGGAACGTGGGATGACTGGGCACCATGAAGAC GTGGAAATGGACCAATGGAGTTCTGAGGACCAGTTTCAG GATATAAGTTTCTTTATACATGATTGGACATGGTATGTTG GACTCCGATCTTCATCTTAGCTCAAAGGCTCAGGTGTTCTGA ACATCCTCACATTCAAGACGCTGCTTCGCAACTTCTGATG ATGAGAGTTTATTTTTTGGTGATCTGGGCTATCCAAAAAT

-continued

SEQ ID	NO: Description	Sequence
		CCAATCGAGCTTGTAGAAGGTTGGTTCAGTAGTTGGAAAA GCTCTATTGCCTCTTTTCTTTATCATAGGGTAAATCATTG GACTATTCTTGGTTCTCCGAGTTGGTATCCATCTTGCATTA AATTAAGCACACCAAGAAAAGACAGATTATACAGACAT AGAGATGA
63	Envelope; Rabbit beta globin poly A; RNA stability	AGATCTTTTCCCTCTGCCAAAAATTATGGGGACATCATGA AGCCCCCTTGAGCATCTGACTTCTGGCTAATAAAGGAAATTT ATTTTCATTGCAATAGTGTGTTGGAATTTTTGTGTCTCTCA CTCGGAAGGACATATGGGAGGGCAAATCATTAAAAACATC AGAATGAGTATTGGTTTAGAGTTTGGCAACATATGCCCAT ATGCTGGCTGCCATGAACAAGGTTGGCTATAAAGAGGTC ATCAGTATATGAAACAGCCCCCTGCTGTCCATTCTTTATTC CATAGAAAAGCCTTGACTTGAGGTTAGATTTTTTTTATATTT TGTTTTGTGTATTTTTTCTTTAACATCCCTAAATTTTCCT TACATGTTTTACTAGCCAGATTTTCCCTCTCTCTGACTAC TCCCAGTCATAGCTGTCCCTCTCTCTTATGGAGATC
64	Promoter; EF- 1	CCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAG TGATGTCGTGTAAGTCCGCTTTTCCCGAGGGTGGGG GAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTCTT TTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAGTGCCGT GTGTGGTTCGCCGCGGCTGGCCTCTTACGGGTTATGGCC CTTGCCTGCCTGAATTACTTCCACGCCCTGGCTGCAGTA CGTGATTCTTGATCCCGAGCTTCGGGTTGGAAGTGGTGGG AGAGTTCGAGGCTTGCGCTTAAGGAGCCCCCTCGCCTCGT GCTTGAGTTGAGGCTGGCCTGGCGCTGGGGCCGCCGCG TGCGAATCTGGTGGCACCTTCGCGCTGTCTCGCTGCTTTC GATAAGTCTCTAGCCATTAAAAATTTTGATGACCTGCTGC GACGCTTTTTTTCTGGCAAGATAGTCTTGTAATGCGGGCC AAGATCTGCACACTGGTATTTTCGGTTTTTGGGGCCGCCGGC GGCGACGGGGCCGCTGCGTCCAGCGCACATGTTCCGGCA GGCGGGGCTGCGAGCGCGGCCACCGAGAATCGGACGGGG GTAGTCTCAAGCTGGCGGCTGCTCTGGTGCCTGGCCTCG CGCCGCCGTGTATCGCCCCGCGCTGGCGGCAAGGCTGGC CCGGTCGGCACCACTTGCGTGAAGCGGAAGATGGCCGCTT CCCCGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGC GCTCGGAGAGCGGGCGGGTGAAGTCAACACACAAAGGAA AAGGGCTTTCCGTCTCAGCCGTGCTTCATGTGACTCCA CGGAGTACCGGGCGCGTCCAGGCACCTCGATTAGTCTCTG AGCTTTTGGAGTACGTCGCTTTAGGTTGGGGGAGGGGTT TTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTG AAGTTAGGCCAGCTTGGCACTTGATGTAATTTCTCTTGAA TTTGCCCTTTTGTAGTTTGGATCTTGGTTCAATCTCAAGCCT CAGACAGTGGTTCAAAGTTTTTTCTTCCATTTCAAGTGTCTG TGA
65	Promoter; PGK	GGGGTTGGGGTTGCGCTTTTCCAAGGCAGCCCTGGGTTTG CGCAGGGACGCGGCTGCTCTGGGCGTGGTTCCGGGAAACG CAGCGCGCGACCCCTGGGTCTCGCACATTTCTACGTCGG TTCGACGCTACCCGGAATCTTCGCGCTACCCCTTGTTGGC CCCCCGGCGACGCTTCTGCTCCGCCCTAAGTCGGGAAGG TTCCTTGCGGTTTCGCGGCTGCGGACGTGACAAACGGAA GCCGCACGTCTCACTAGTACCTTCGCAGACGGACAGCGCC AGGGAGCAATGGCAGCGCGCCGACCGGATGGGCTGTGGC CAATAGCGGCTGCTCAGCAGGCGCGCGAGAGCAGCGGC CGGGAAGGGCGGTGCGGAGGCGGGGTGTGGGCGGTA GTGTGGGCCCTGTTCTGCCCCGCGGGTTCCTCGATTCTG CAAGCCTCCGGAGCGCACGTGCGCAGTCGGCTCCCTCGTTG ACCGAATCACCGACCTCTCTCCCCAG
66	Promoter; UbC	GCGCCGGGTTTTGGCGCTTCCCGCGGGCGCCCCCTCCTCA CGGCGAGCGCTGCCACGTGACGAGAGGGCGCAGGAGCGT TCCTGATCCTTCCGCCGAGCGCTCAGGACAGCGGCCCGCT GCTCATAAGACTCGGCCTTAGAACCCAGTATCAGCAGAA GGACATTTTAGGACGGGACTTGGGTGACTCTAGGCGACTG GTTTTCTTCCAGAGAGCGGAACAGCGAGGAAAAGTAGT CCCTTCTCGGCGATTCTGCGGAGGATCTCCGTGGGGCGGT GAACGCCGATGATTATATAAGGACGCGCGGGTGTGGCAC AGCTAGTTCCGTCGAGCCGGGATTTGGGTCGCGGTTCTTG TTGTGGATCGCTGTGATCGTCACTTGGTGAAGTTCGGGCT GCTGGGCTGGCCGGGGCTTTCGTGGCCGCCGGGCCGCTCG

-continued

SEQ ID NO: Description	Sequence
	GTGGGACGGAAGCGTGTGGAGAGACCGCAAGGGCTGTAG TCTGGGTCCGCGAGCAAGGTTGCCCTGAACTGGGGGTTGG GGGGAGCGCACAAAATGGCGGCTGTCCCGAGTCTTGAAT GGAAGACGCTTGTAAAGCGGGCTGTGAGGTCGTTGAAACA AGGTGGGGGGCATGGTGGGCGCAAGAACCCAAAGGCTCTG AGGCCTTCGCTAATGCGGGAAAGCTCTTATTCGGGTGAGAT GGGCTGGGGCACCCTCTGGGGACCCTGACGTGAAGTTTGT CACTGACTGGAGAACTCGGGTTTGTCTGTCTGGTTGCGGGG CGGCAGTTATGCGGTGCCGTGGGCAGTGACCCGTACCTT TGGGAGCGCGCGCCTCGTCTGTCTGTGACGTACCCGTTCT GTTGGCTTATAATGACAGGTTGGGGCACCTGCCGGTAGGT GTGCGGTAGGCTTTTCTCGTCGACGACGACGGGTTTCGGG CCTAGGTTAGGCTCTCCTGAATCGACAGGCGCCGGACCTCT GGTGGGGGAGGGATAAGTGAGGCGTCAGTTTCTTTGGTC GGTTTTATGTACCTATCTTCTTAAGTAGCTGAAGCTCCGGT TTGAACATGCGCTCGGGTTGGCGAGTGTGTTTTGTGAAG TTTTTTAGGCACCTTTTGAATGTAATCATTTGGGTCAATAT GTAATTTTCAGTGTTAGACTAGTAAA
67 Poly A; SV40	GTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCA TCACAAATTTCACAAATAAAGCATTTTTTCAGTCATTCT AGTTGTGGTTTGTCCAACTCATCAATGTATCTTATCA
68 Poly A; bGH	GACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTGCCCTC CCCCGTGCCTTCTTGACCTTGAAGGTGCCACTCCCCTG TCCTTTCTTAATAAATGAGGAAATGTCATCGCATGTCTG AGTAGGTGTCTTCTATTCTGGGGGTGGGTGGGGCAGG ACAGCAAGGGGAGGATTGGGAAGACAATAGCAGGCATG CTGGGGATGCGGTGGGCTCTATGG
69 HIV Gag; Bal	ATGGGTGCGAGAGCGTCAGTATTAAGCGGGGAGAATTAG ATAGGTGGGAAAAATTCGGTTAAGGCCAGGGGGAAAGA AAAAATATAGATTAAACATATAGTATGGGCAAGCAGGGA ACTAGAAAGATTGCGAGTCAATCCTGGCCTGTTAGAAACAT CAGAAGGCTGCAGACAAATACTGGGACAGCTACAACCATC CCTTCAGACAGGATCAGAAGAACTTAGATCATTATATAATA CAGTAGCAACCTCTATTGTGTACATCAAAAGATAGAGGT AAAAGACACCAAGGAAGCTTTAGACAAATAGAGGAAGA GCAAAACAAATGTAAGAAAAAGGCACAGCAAGCAGCAGC TGACACAGGAAACAGCGGTCAGGTCAGCCAAAATTTCCCT ATAGTGCAGAACCTCCAGGGGCAATGTTACATCAGGCCA TATCACCTAGAACTTTAATGCAATGGGTAAAGTAATAGA AGAGAAAGCTTTAGCCAGAAAGTAATACCCATGTTTTAG CATTATCAGAAGGAGCCACCCACAAAGATTTAAACACCAT GCTAAACACAGTGGGGGACATCAAGCAGCCATGCAAAATG TTAAAGAAACCCATCAATGAGGAAGCTGCAAGATGGGATA GATTGCATCCCCTGTCAGGCAGGGCCTGTTGCCACGAGCCA GATAAGAGATCCAAGGGGAAGTGACATAGCAGGAATACC AGTACCCTTCAGGAACAAATAGGATGGATGACAAGTAATC CACCTATCCAGTAGGAGAAATCTATAAAGATGGATAAT CCTGGGATTAAATAAAATAGTAAGGATGTATAGCCCTACC AGCATTTTGGACATAAGACAAGGACCAAGGAACCTTTA GAGACTATGTAGACCGGTTCTATAAACTCTAAGAGCCGA GCAAGCTTCACAGGAGGTAAAAAATTGGATGACAGAAACC TTGTTGGTCCAAAATGCGAACCCAGATTGTAAGACTATTTT AAAAGCATTTGGACACAGCAGCTACACTAGAAGAAATGATG ACAGCATGTGAGGAGTGGGAGGACCCAGCCATAAAGCAA GAATTTTGGCAGAAGCAATGAGCCAAGTAACAAATTCAGC TACCATAATGATGCAGAAAGGAATTTTAGGAACCAAGA AAGATTGTTAAATGTTTCAATTGTGGCAAGAAAGGACACA TAGCCAGAACTGCAGGGCCCTAGGAAAAGGGGCTGTTG GAAATGTGAAAGGAAGGACACCAATGAAAGACTGTACT GAGAGACAGGCTAATTTTTAGGGAAAACTTGCCCTTCCCA CAAAGGAAGGCCAGGGAATTTCTTCAGAGCAGACCAGAG CCAACAGCCCCACGACCCACAGAAAGAGAGCTTCAGGT TTGGGGAAGAGACAACAACTCCCTCTCAGAAGCAGGAGCT GATAGACAAGGAACGTATCTTTAGCTTCCTCTCAGATCAC TCTTTGGCAACGACCCCTCGTCACAATAA
70 HIV Pol; Bal	ATGAATTTGCCAGGAAGATGGAAACCAAAATGATAGGGG GAATTGGAGTTTATCAAAGTAAGACAGTATGATCAGAT ACTCATAGAAATCTGTGGACATAAAGCTATAGGTACAGTA

-continued

SEQ ID NO: Description	Sequence
	<p>TTAATAGGACCTACACCTGTCAACATAATTGAAGAAATCT GTTGACTCAGATTGGTTGCACCTTAAATTTTCCATTAGTCC TATTGAACTGTACCAGTAAATTTAAACCAGGAATGGAT GGCCCCAAAAGTTAAACAATGGCCACTGACAGAAGAAAAA TAAAAGCATTAATGGAAATCTGTACAGAAATGGAAAAGGA AGGGAAAATTTCAAAAATTGGGCTGAAAATCCATACAA ACTCCAGTATTTGCCATAAAGAAAAAGACAGTACTAAAT GGAGAAAATTAGTAGATTTCAGAGAACTTAATAAGAAAAC TCAAGACTTCTGGGAAGTACAATTAGGAATACACATCCCG CAGGGGTAAAAAGAAAAAATCAGTAACAGTACTGGATG TGGGTGATGCATATTTTTCAGTTCCTTAGATAAAGAATT AGGAAGTACTGCAATTTACCATACCTAGTATAAACAAATGA AACACCAGGGATCAGATATCAGTACAATGTACTTCCACAG GGATGGAAAGGATCACCAGCAATATTTCAAAGTAGCATGA CAAGAACTTAGAGCCTTTAGAAAACAAAATCCAGAAAT AGTGATCTATCAATACATGGATGATTTGTATGTAGGATCTG ACTTAGAAATAGGGCAGCATAGAACAAAAATAGAGGAACT GAGACAACATCTGTTGAGGTGGGATTTACCACACCAGAC AAAAAACATCAGAAAGAACCTCCATTCTTTGGATGGGTT ATGAACTCCATCCTGATAAATGGACAGTACAGCCTATAGTG CTGCCAGAAAAGACAGCTGGACTGTCAATGACATACAGA AGTTAGTGGGAAAATTGAATTGGGCAAGTCAGATTTACCC AGGAATTAAGTAAAGCAATATGTAGGCTCCTTAGGGGA ACCAAGGCATTAACAGAGTAATACCACTAACAAAAGAAA CAGAGCTAGAACTGGCAGAGAACAGGGAATTTCTAAAAGA ACCAGTACATGGGGTGATTTATGACCCATCAAAGACTTA ATAGCAGAAATACAGAGCAGGGGCAAGGCCAATGGACA TATCAAATTTATCAAGAGCCATTTAAAAATCTGAAAACAG GAAAATATGCAAGAATGAGGGGTGCCACACTAATGATGT AAAACAATTAAACAGAGGCAGTGCAAAAATAACACAGA AAGCATAGTAATATGGGAAAGACTCCTAAATTTAAACTA CCCATACAAAAAGAAACATGGGAAACATGGTGACAGAGT ATTGGCAAGCCACCTGGATTCTGTAGTGGGAGTTTGTCAAT ACCCCTCCCTTAGTGAAATTTATGGTACCAGTTAGAGAAAGA ACCCATAATAGGAGCAGAAACATTCTATGTAGATGGAGCA GCTAACCCGGGAGACTAAATTAGGAAAAGCAGGATATGTTA CTAACAGAGGAAGACAAAAGTTGTCTCCCTAACTGACAC AACAAATCAGAAAGCTGAGTTACAAGCAATTCATCTAGCT TTACAAGATTCAGGATTAGAAGTAACATAGTAACAGACT CACAAATATGCATTAGGAATCATTCAAGCACAACCAGATAA AAGTGAATCAGAGTTAGTCAGTCAAATAATAGAACAGTTA ATAAAAAAGGAAAAGGTCTACCTGGCATGGGTACCAGCGC ACAAAGGAATTGGAGGAATGAACAAGTAGATAAATTAGT CAGTACTGGAATCAGGAAAGTACTA</p>
71 HIV Integrase; Bal	<p>TTTTTAGATGGAATAGATATAGCCCAAGAAGAACATGAGA AATATCACAGTAATTTGGAGAGCAATGGCTAGTGATTTTAA CTGCCACCTGTGGTAGCAAAAGAAATAGTAGCCAGCTGTG ATAAATGTCAGCTAAAAGGAGAAGCCATGCATGGACAAGT AGACTGTAGTCCAGGAATATGGCACTAGATTGTACACATT TAGAAGGAAAAATTATCTGTGTAGCAGTTTATGTAGCCAG TGGATATATAGAAGCAGAAGTTATCCAGCAGAGACAGGG CAGGAAACAGCATACTTTCTCTTAAATTAGCAGGAAGAT GGCCAGTAAAAACAATACATACAGACAATGGCAGCAATTT CACTAGTACTACAGTCAAGGCCGCCCTGTTGGTGGGCGGG ATCAAGCAGGAATTTGGCATTCCTTACAATCCCCAAAGTCA GGGAGTAGTAGAATCTATAAATAAGAAATTAAGAAAAATT ATAGGACAGGTAAGAGATCAGGCTGAACATCTTAAACAG CAGTACAAATGGCAGTATTCATCCCAATTTTAAAGAAA AGGGGGGATTGGGGGTATAGTGCAGGGGAAAGAAATAGT AGACATAATAGCAACAGACATACAACTAAAGAATTACAA AAACAAATTACAAAATTTCAAATTTTCGGGTTTATTACAG GGACAGCAGAGATCCACTTTGGAAAGGACCAGCAAAGCTT CTCTGGAAGGTGAAGGGCAGTAGTAATACAAGATAATA GTGACATAAAAGTAGTACCAAGAAGAAAAGCAAGATCAT TAGGGATTATGGAAAACAGATGGCAGGTGATGATTGTGTG GCAAGTAGACAGGATGAGGATTAG</p>
72 Envelope; RD114	<p>ATGAACTCCCAACAGGAATGGTCATTTTATGTAGCCTAAT AATAGTTCGGGCAGGGTTTGACGACCCCGCAAGGCTATC GCATTAGTACAAAACAAATGGTAAACCATGCGAATGCA GCGGAGGGCAGGTATCCGAGGCCCCACCGAATCCATCCA</p>

-continued

SEQ ID NO: Description	Sequence
	ACAGGTAACCTTGCCAGGCAAGACGGCCTACTTAATGACC AACCAAAATGGAAATGCAGAGTCACTCCAAAAATCTCA CCCCTAGCGGGGGAAGTCCAGAACTGCCCTGTAAACAC TTTCCAGGACTCGATGCACAGTCTTGTGTATACTGAATACC GGCAATGCAGGGCGAATAATAAGACATACTACACGGCCAC CTTGCTTAAATACGGTCTGGGAGCCTCAACGAGGTACAG ATATTACAAAACCCCAATCAGCTCCTACAGTCCCCTTGTAG GGGCTCTATAAATCAGCCGTTTGTGGAGTGCCACAGCCC CCATCCATATCTCCGATGGTGGAGACCCCTCGATACTAAG AGAGTGTGGACAGTCCAAAAAGGCTAGAACAATTCATA AGGCTATGCATCCTGAACCTCAATACCAACCCCTAGCCCTG CCCAAGTCAGAGATGACCTTAGCCTTGATGCACGGACTTT TGATATCCTGAATACCACTTTTAGGTTACTCCAGATGTCCA ATTTTAGCCTTGCCCAAGATTGTTGGCTCTGTTTAAACTA GGTACCCCTACCCCTCTTGCATACCCACTCCCTCTTTAAC TACTCCCTAGCAGACTCCCTAGCGAATGCCTCCTGTCAGAT TATACCTCCCTCTTGGTTCAACCGATGCAGTCTCTCAACTC GTCCTGTTTATCTTCCCTTTTATTAAACGATACGGAACAAA TAGACTTAGGTGCAGTCACCTTTACTAAGTCACCTCTGTA GCCAATGTGAGTAGTCTTATGTGCCCTAAACGGGTGAGT CTTCTCTGTGGAAATAACATGGCATACACTATTATACCCC AAAAGTGGACAGGACTTTGCGTCCAAGCCTCCCTCCTCCCC GACATTGACATCATCCCGGGGATGAGCCAGTCCCATTC TGCCATTGATCATTATATACATAGACCTAAACGAGCTGTAC AGTTCATCCCTTTACTAGCTGGACTGGGAATCACCGCAGCA TTCACCAACGGAGCTACAGGCTAGGTGTCTCCGTCAACCA GTATACAAAATTATCCCATCAGTTAATATCTGATGTCCAAG TCTTATCCGGTACCATACAAGATTTACAAGACCAGGTAGAC TCGTTAGCTGAAGTAGTTCTCCAAATAGGAGGGGACTGG ACCTACTAACGGCAGAACAAAGGAGGAATTTGTTAGCCTT ACAAGAAAAATGCTGTTTTATGCTAACAAAGTCAGGAATTG TGAGAAACAAAATAAGAACCTTACAAGAGAATTACAAAA ACGCAGGGAAAGCCTGGCATCCAACCTCTCTGGACCGGG CTGCAGGGCTTTCTTCCGTACCTCCTACCTCTCCTGGGACCC CTACTCACCTCCTACTCATATAACCATTTGGGCCATGCGT TTTCAATCGATTGGTCCAATTGTTAAAGACAGGATCTCAG TGGTCCAGGCTCTGGTTTTGACTCAGCAATATCACAGCTA AAACCCATAGAGTACGAGCCATGA
73 Envelope; GALV	ATGCTTCTCACCTCAAGCCCGCACCACCTTCGGCACCAGAT GAGTCTGGGAGCTGGAAAAGACTGATCATCTCTTAAGCT CGGTATTGAGAGACGGCAAAACGAGTCTGCAGAAATAAGAA CCCCCACCAGCTGTGACCTCACCTGGCAGGTACTGTCCC AAACTGGGACGTTGTCTGGGACAAAAGGCAGTCCAGCC CCTTTGGACTTGGTGGCCCTCTTTACACCTGATGTATGTGC CCTGGCGGCGGTCTTGAATCCTGGGATATCCCGGGATCCG ATGTATCGTCTCTAAAAGAGTTAGACCTCCTGATTACAGAC TATACTGCCGCTTATAAGCAATCACCTGGGGAGCCATAG GGTGCAGCTACCTCGGGCTAGGACAGGATGGCAAAATTC CCCCTTCTACGTGTGTCCCGAGCTGGCCGAACCCATTACAG AAGCTAGGAGGTGTGGGGGCTAGAATCCCTATCTGTAA AGAATGGAGTTGTGAGACCACGGGTACCGTTTATTGGCAA CCCAAGTCTCATGGGACCTCATAACTGTAAATGGGACC AAAATGTGAAATGGGAGCAAAATTTCAAAAGTGTGAACA AACCAGGCTGGTGTAAACCCCTCAAGATAGACTTCACAGAA AAAGGAAAACCTCTCAGAGATTGGATAACGGAAGAAACCT GGGAATTAAGGTTCTATGTATATGGACACCCAGGCATACA GTTGACTATCCGCTTAGAGGTCACTAACATGCCGTTGTGG CAGTGGGCCAGACCTGTCTTGCAGAACAGGGACCTCCT AGCAAGCCCTCTACTCTCCCTCTCTCCCAAGGAAAGCGCC GCCCACCCCTCTACCCCGGGCGCTAGTGAGCAACCCCTG CGGTGCATGGAGAACTGTTACCTAAACTCTCCGCTCCC ACCAAGTGGGACCGACTCTTTGGCCTTGTGACGGGGCCCTT CCTAACCTTGAATGCTACCAACCCAGGGGCCACTAAGTCTT GCTGGCTCTGTTGGGCATGAGCCCCCTTATTATGAAGGG ATAGCCTCTTCAAGGAGAGGTGCTTATACCTCCAACCATAC CCGATGCCACTGGGGGCCCAAGGAAAGCTTACCTCACT GAGGTCTCCGACTCGGGTCATGCATAGGAAGGTGCCTC TTACCCATCAACATCTTTGCAACCAAGCTTACCCATCAAT TCCTCTAAAAACCATCAGTATCTGCTCCCTCAAACCATAG CTGGTGGGCTGCAGCACTGGCCTCACCCCTGCCTCTCCA CCTCAGTTTTTAATCAGTCTAAGACTTCTGTGTCCAGGTC

-continued

SEQ ID	NO: Description	Sequence
		CAGCTGATCCCCGCATCTATTACCATTCTGAAGAAACCTT GTTACAAGCCTATGACAAATCACCCCCAGGTTTAAAGA GAGCCTGCCTCACTTACCCTAGCTGTCTTCTGGGGTTAGG GATTGCGGCAGGTATAGGTACTGGCTCAACCGCCCTAATTA AAGGGCCCATAGACCTCCAGCAAGGCCTAACCAGCCTCCA AATCGCCATTGACGCTGACCTCCGGGCCCTTCAGGACTCAA TCAGCAAGCTAGAGGACTCACTGACTTCCCTATCTGAGGTA GTACTCCAAAATAGGAGAGGCCCTTGACTTACTATTCTTAA AGAAGGAGGCCTCTGCGCGGCCCTAAAGAAGAGTGCTGT TTTTATGTAGACCACTCAGGTGCAGTACGAGACTCCATGAA AAAACTTAAAGAAAGACTAGATAAAAGACAGTTAGAGCGC CAGAAAAACCAAACTGGTATGAAGGGTGGTTCAATAACT CCCCTTGGTTTACTACCCTACTATCAACCATCGCTGGGCCC CTATTGCTCCTCCTTTTGTACTCACTCTTGGGCCCTGCATC ATCAATAAATTATCCAATTCAATGATAGGATAAGTGC AGTCAAAATTTTAGTCCCTTAGACAGAAATATCAGACCCTAG ATAACGAGGAAACCTTTAA
74	Envelope; FUG	ATGGTTCCGCAGGTTCCTTTTGTGTACTCCTTCTGGGTTTT TCGTTGTGTTTCGGGAAGTTCCTTACACGATACCAGA CGAACTTGGTCCCTGGAGCCCTATTGACATACACCATCTCA GCTGTCCAAATAACCTGGTTGTGGAGGATGAAGGATGTAC CAACCTGTCCGAGTTCTCCTACATGGAACCTCAAAGTGGGAT ACATCTCAGCCATCAAAGTGAACGGGTTCACTTGACAGGT GTTGTGACAGAGGCAGAGACCTACACCACTTTGTTGGTTA TGTCAACACCATTCAGAGAAAGCATTTCCGCCCAACCC CAGACGCATGTAGAGCCGCGTATAACTGGAAGATGGCCGG TGACCCAGATATGAAGAGTCCCTACACAATCCATACCCCG ACTACCCTGGCTTCGAAGTGAAGAACCACCAAGAGTC CCTCATTATCATATCCCCAAGTGTGACAGATTGGACCCAT ATGACAAATCCCTTCACTCAAGGGTCTTCCCTGGCGGAAAG TGCTCAGGAATAACGGTGTCTCTACCTACTGCTCAACTAA CCATGATTACACCATTTGGATGCCCGAGAATCCGAGACCA AGGACACCTTGTGACATTTTACCATAGCAGAGGGAAGA GAGCATCCAACGGGAACAGACTTGGCGCTTGTGGATGA AAGAGGCCTGTATAAGTCTCTAAAAGGAGCATGCAGGCTC AAGTTATGTGGAGTTCTTGGACTTAGACTTATGGATGGAAC ATGGGTCGCGATGCAACATCAGATGAGACCAATGGTGC CCTCCAGATCAGTTGGTGAATTTGCACGACTTTCGCTCAGA CGAGATCGAGCATCTCGTTGTGGAGGAGTTAGTTAAGAAA AGAGAGGAATGTCTGGATGCATTAGAGTCCATCATGACCA CCAAGTCAGTAAGTTTCAGACGTCTCAGTCACCTGAGAAA ACTTGTCCAGGGTTTGGAAAAGCATATACCATATTCAACA AAACCTTGTAGGAGGCTGATGCTCACTACAAGTCAGTCCG GACCTGGAATGAGATCATCCCTCAAAGGGTGTGTTGAAA GTTGGAGGAAGTGCCATCCTCATGTGAACGGGGTGTGTTTT CAATGGTATAATATTAGGGCTGACGACCATGCTCCTAATCC CAGAGATGCAATCATCCCTCCTCCAGCAACATAGGAGTTG TTGGAATCTTCAGTTATCCCCCTGATGCACCCCTGGCAGA CCCTTCTACAGTTTCAAAGAAGGTGATGAGGCTGAGGATT TTGTTGAAGTTCACTCCCCGATGTGTACAAACAGATCTCA GGGGTTGACCTGGGTCTCCCGAAGTGGGAAAGTATGTATT GATGACTGCAGGGGCCATGATTGGCCTGGTGTGATATTTT CCTAATGACATGGTGCAGAGTTGGTATCCATCTTTGCATT AAATTAAAGCACACCAAGAAAAGACAGATTTATACAGACA TAGAGATGAACCGACTTGGAAGTAA
75	Envelope; LCMV	ATGGGTCAGATTGTGACAATGTTTGGAGGCTCTGCCTCACAT CATCGATGAGGTGATCAACATTGTCTATTATTGCTTATCG TGATCAGGGTATCAAGGCTGTCTACAATTTTGCCACCTGT GGGATATTTCGATTGATCAGTTTCCTACTTCTGGCTGGCAG GTCCGTGGCATGTACGGTCTTAAGGGACCCGACATTTACA AAGGAGTTTACCAATTTAAGTCAGTGGAGTTTGATATGTCA CATCTGAACCTGACCATGCCCAACGCATGTTGAGCCAACAA CTCCCACCATTACATCAGTATGGGACTTCTGGACTAGAAT TGACCTTCAACCAATGATTCCATCATCAGTCACAACCTTTGC AATCTGACCTCTGCCTTCAACAAAAGACCTTTGACCACAC ACTCATGAGTATAGTTTTCGAGCCTACACCTCAGTATCAGAG GGAACCTCAACTATAAGGCAGTATCTGCGACTTCAACAAT GGCATAACCATCCAATACAACCTTGACATTCTCAGATCGACA AAGTGCTCAGAGCCAGTGTAGAACCTTCAGAGGTAGAGTC CTAGATATGTTTGAAGTGCCTTCGGGGGGAATACATGA

-continued

SEQ ID NO: Description	Sequence
	GGAGTGGCTGGGGCTGGACAGGCTCAGATGGCAAGACCAC CTGGGTAGCCAGACGAGTTACCAATACCTGATTATACAAA ATAGAACCTGGGAAACCCTGCACATATGCAGGTCCTTTT GGGATGTCCAGGATTCTCCTTTCCCAAGAGAAGACTAAGTT CTTCACTAGGAGACTAGCGGGCACATTACCTGGACTTTGT CAGACTCTTCAGGGGTGGAGAATCCAGGTGGTTATTGCCTG ACCAAAATGGATGATTCTTGCTGCAGAGCTTAAGTGTTCGG GAACACAGCAGTTGCGAAATGCAATGTAAATCATGATGCC GAATTCTGTGACATGCTGCGACTAATTGACTACAACAAGGC TGCTTTGAGTAAGTTCAAAGAGGACGTAGAATCTGCCTTGC ACTTATTCAAAACAACAGTGAATTCTTTGATTTCAGATCAA CTACTGATGAGGAACCACTTGAGAGATCTGATGGGGTGC CATATTGCAATTACTCAAAGTTTTGGTACCTAGAACATGCA AAGACCGGCGAAACTAGTGTCCCCAAGTGTGGCTTGTCA CCAATGGTTCTTACTTAAATGAGACCCACTTCAGTGATCAA ATCGAACAGGAAGCCGATAACATGATTACAGAGATGTTGA GGAAGGATTACATAAAGAGGCGAGGGAGTACCCCTAGC ATTGATGGACCTTCTGATGTTTTCCACATCTGCATATCTAGT CAGCATCTTCTGCACCTTGTCAAATACCAACACACAGGC ACATAAAAGGTGGCTCATGTCCAAAGCCACACCGATTAAAC CAACAAAGGAATTTGTAGTTGTGGTGCAATTAAGGTGCCTG GTGTAAAAACCGTCTGGAAGAGACGCTGA
76 Envelope; FPV	ATGAACACTCAAATCCTGGTTTTTCGCCCTTGTGGCAGTCAT CCCCACAAATGCAGACAAAATTTGTCTTGGACATCATGCTG TATCAAATGGCACCAGTAAACACACTCACTGAGAGAGG AGTAGAAGTTGTCAATGCAACGGAACAGTGGAGCGGACA AACATCCCCAAAATTTGCTCAAAAGGGAAGAACCACTG ATCTTGGCCAAATGCGGACTGTTAGGGACCATTACCGGACCA CCTCAATGCGACCAATTTCTAGAATTTTCAGCTGATCTAAT AATCGAGAGACGAGAAGGAAATGATGTTTGTACCCGGGG AAGTTTGTAAATGAAGAGGCATTGCGCAAAATCCTCAGAG GATCAGGTGGGATTGACAAAGAAACAATGGGATTCACATA TAGTGGAATAAGGACCAACGGAACAACAGTGCATGTAGA AGATCAGGGTCTTCAATCTATGCAGAAATGGAGTGGCTCCT GTCAAATACAGACAATGCTGCTTTCCCAAAATGACAAA TCATACAAAACACAAGGAGAGAATCAGCTCTGATAGTCT GGGGAATCCACCATTCAGGATCAACCACCGAACAGACCAA ACTATATGGGAGTGGAAATAAACTGATAACAGTCGGGAGT TCCAAATATCATCAATCTTTGTGCCGAGTCCAGGAACACG ACCGCAGATAAATGGCCAGTCCGGACGGATTGATTTTCATT GGTTGATCTTGGATCCCAATGATACAGTTACTTTTAGTTTC AATGGGGCTTTCATAGCTCCAAATCGTGCCAGCTTCTTGAG GGGAAAGTCCATGGGGATCCAGAGCGATGTGCAGGTTGAT GCCAATTGCGAAGGGGAATGCTACCACAGTGGAGGACTA TAACAAGCAGATTGCCCTTTTCAAACATCAATAGCAGAGC AGTTGGCAAATGCCCAAGATATGTAAAACAGGAAAGTTTA TTATTGGCAACTGGGATGAAGAACGTTCCGAACCTTCCAA AAAAAGGAAAAAAGAGGCCCTGTTTGGCGCTATAGCAGGG TTTATTGAAAAATGGTTGGGAAGGTCTGGTCGACGGGTGGTA CGTTTCAGGCATCAGAAATGCACAAGGAGAAGGAACGCA GCAGACTACAAAAGCACCCAATCGGCAATTGATCAGATAA CCGAAAAGTTAAATAGACTCATTGAGAAAACCAACCAGCA ATTTGAGCTAATAGATAATGAATTCAGTGGGTGGAAG CAGATTGGCAATTTAATTAACCTGGACCAAGACTCCATCAC AGAAGTATGGTCTTACAATGCTGAACCTCTTGTGGCAATGG AAAACGACACACTATTGATTTGGCTGATTCAGAGATGAA CAAGCTGTATGAGCGAGTGAGGAAACAATTAAGGGAAAAAT GCTGAAGAGGATGGCACTGGTTGCTTTGAAATTTTTCATAA ATGTGACGATGATTGTATGGCTAGTATAAGGAACAATACTT ATGATCACAGCAAAATACAGAGAAGAAGCGATGCAAAATAG AATACAAATTGACCCAGTCAAATTGAGTAGTGGCTACAAA GATGTGATACTTTGGTTTAGCTTCGGGCATCATGCTTTTTG CTTCTTGCCATTGCAATGGGCCTTGTTTTCATATGTGTGAAG AACGGAACATGCGGTGCCTATTGTATATAA
77 Envelope; RRV	AGTGTAACAGAGCACTTTAATGTGTATAAGGCTACTAGACC ATACCTAGCACATTGCGCCGATTGCGGGGACGGGTACTTCT GCTATAGCCAGTTGCTATCGAGGAGATCCGAGATGAGGC GTCTGATGGCATGCTTAAGATCCAAGTCTCCGCCCAATAG GTCTGGACAAGGCAGGCACCCACGCCACACGAAGCTCCG ATATATGGCTGGTCATGATGTTCAGGAATCTAAGAGAGATT

-continued

SEQ ID NO: Description	Sequence
	CCTTGAGGGTGTACACGTCCGCAGCGTGCTCCATACATGGG ACGATGGGACACTTCATCGTCGCACACTGTCCACCAGGCG ACTACCTCAAGGTTTCGTTTCGAGGACGCAGATTGCGACGTG AAGGCATGTAAGGTCCAATACAAGCACAAATCCATTGCCGG TGGGTAGAGAGAAGTTCTGTGGTTAGACCACACTTTGGCGT AGAGCTGCCATGCACCTCATACCAGCTGACAACGGCTCCC ACCGACGAGGAGATTGACATGCATACACCGCCAGATATAC CGGATCGCACCTGTATCACAGACGGCGGGCAACGTCAA AATAACAGCAGGCGGCAGGACTATCAGGTACAACGTGACC TGCGGCCGTGACAACGTAGGCACTACCAGTACTGACAAGA CCATCAACACATGCAAGATTGACCAATGCCATGTGCCGTG ACCAGCCATGACAAATGGCAATTTACCTCTCCATTGTGTTCC CAGGGCTGATCAGACAGCTAGGAAAGGCAAGGTACACGTT CCGTTCCCTCTGACTAACGTACCTGCCAGTGCCGTTGGC TCGAGCGCCGATGCCACCTATGGTAAGAAGGAGGTGACC CTGAGATTACACCCAGATCATCCGACGCTCTTCTCCTATAG GAGTTTAGGAGCCGAACCGCACCCGTACGAGGAATGGGTT GACAAGTTCTCTGAGCGCATCATCCAGTGACGGAAGAAG GGATTGAGTACCAGTGGGGCAACAACCCGCCGCTGCTCCT GTGGGCGCAACTGACGACCGAGGGCAAAACCCATGGCTGG CCACATGAAATCATTAGTACTATTATGGACTATACCCGCG CGCCACTATTGCGCGAGTATCCGGGGCGAGTCTGATGGCCC TCCTAACTCTGGCGGCCACATGCTGCATGCTGGCCACCGCG AGGAGAAAGTGCCCTAACACCGTACGCCCTGACGCCAGGAG CGGTGGTACCGTTGACACTGGGGTGCTTTGCTGCGCACCG AGGGCGAATGCA
78 Envelope; MLV 10A1	AGTGTAAACAGACACTTTAATGTGTATAAGGCTACTAGACC ATACCTAGCACATTGCGCCGATTGCGGGGACGGGTACTTCT GCTATAGCCAGTTGCTATCGAGGAGATCCGAGATGAGGC GTCTGATGGCATGCTTAAGATCCAAGTCTCCGCCAAATAG GTCTGGACAAGGCAGGCCACCGCCACACGAAGCTCCG ATATATGGCTGGTCATGATGTTCAAGAACTAAGAGAGATT CTTTGAGGGTGTACACGTCCGCAGCGTGCTCCATACATGGG ACGATGGGACACTTCATCGTCGCACACTGTCCACCAGGCG ACTACCTCAAGGTTTCGTTTCGAGGACGCAGATTGCGACGTG AAGGCATGTAAGGTCCAATACAAGCACAAATCCATTGCCGG TGGGTAGAGAGAAGTTCTGTGGTTAGACCACACTTTGGCGT AGAGCTGCCATGCACCTCATACCAGCTGACAACGGCTCCC ACCGACGAGGAGATTGACATGCATACACCGCCAGATATAC CGGATCGCACCTGTATCACAGACGGCGGGCAACGTCAA AATAACAGCAGGCGGCAGGACTATCAGGTACAACGTGACC TGCGGCCGTGACAACGTAGGCACTACCAGTACTGACAAGA CCATCAACACATGCAAGATTGACCAATGCCATGTGCCGTG ACCAGCCATGACAAATGGCAATTTACCTCTCCATTGTGTTCC CAGGGCTGATCAGACAGCTAGGAAAGGCAAGGTACACGTT CCGTTCCCTCTGACTAACGTACCTGCCAGTGCCGTTGGC TCGAGCGCCGATGCCACCTATGGTAAGAAGGAGGTGACC CTGAGATTACACCCAGATCATCCGACGCTCTTCTCCTATAG GAGTTTAGGAGCCGAACCGCACCCGTACGAGGAATGGGTT GACAAGTTCTCTGAGCGCATCATCCAGTGACGGAAGAAG GGATTGAGTACCAGTGGGGCAACAACCCGCCGCTGCTCCT GTGGGCGCAACTGACGACCGAGGGCAAAACCCATGGCTGG CCACATGAAATCATTAGTACTATTATGGACTATACCCGCG CGCCACTATTGCGCGAGTATCCGGGGCGAGTCTGATGGCCC TCCTAACTCTGGCGGCCACATGCTGCATGCTGGCCACCGCG AGGAGAAAGTGCCCTAACACCGTACGCCCTGACGCCAGGAG CGGTGGTACCGTTGACACTGGGGTGCTTTGCTGCGCACCG AGGGCGAATGCA
79 Envelope; Ebola	ATGGGTGTTACAGGAATATTGCAGTTACCTCGTGATCGATT CAAGAGGACATCATCTTTCTTTGGGTAATTATCCTTTTCCA AAGAACATTTTCCATCCCACTTGGAGTCATCCACAATAGCA CATTACAGGTTAGTGATGTCGACAACTGGTTTGCCGTGAC AAAGTGTATCCACAATCAATTGAGATCAGTTGGACTGA ATCTCGAAGGGAATGGAGTGGCAACTGACGTGCCATCTGC AACTAAAGATGGGGCTTCAGGTCCGGTGTCCCAACCAAG GTGGTCAATTATGAAGCTGGTGAATGGGTGAAAACCTGCT ACAATCTTGAAATCAAAAACCTGACGGGAGTGAGTGTCT ACCAGCAGCGCCAGACGGGATTCGGGGCTTCCCCCGGTGC CGGTATGTGCACAAAGTATCAGGAACGGGACCGTGTGCCG GAGACTTTGCCCTTCCACAAGAGGGTGCTTTCTCTCTGTAT

-continued

SEQ ID NO: Description	Sequence
	GACCGACTTGCTTCCACAGTTATCTACCGAGGAACGACTTT CGCTGAAGGTGTCGTTGCATTTCTGATACTGCCCCAAGCTA AGAAGGACTTCTTCAGCTCACACCCCTTGAGAGAGCCGGTC AATGCAACGGAGGACCCGTCTAGTGGCTACTATTCTACCAC AATTAGATATCAAGCTACCGGTTTTGGAACCAATGAGACA GAGTATTTGTTTCGAGGTTGACAATTTGACCTACGTCCAAC TGAATCAAGATTCACACCACAGTTTCTGCTCCAGCTGAATG AGACAATATATACAAGTGGGAAAAGGAGCAATACCACGGG AAAACATAATTTGGAAGGTCAACCCGAAATTGATACAACA ATCGGGGAGTGGGCCTTCTGGGAACTAAAAAACCTCAC TAGAAAAATTCGAGTGAAGAGTTGTCTTTCACAGCTGTAT CAACACAGAGCCAAAACATCAGTGGTCAGAGTCCGGCGCG AACTTCTTCCGACCCAGGACCAACACAACCACTGAAGAC CACAAAATCATGGCTTCAGAAAATTCCTCTGCAATGGTTCA AGTGACAGCTCAAGGAAGGGAAGCTGCAGTGTGCGATCTG ACAACCCCTTGCCACAATCTCCACGAGTCTCAACCCCCCAC AACCAACCCAGGTCGGGACCAACAGCACCCACAATACACCC GTGTATAAACTTGACATCTCTGAGGCAACTCAAGTTGAACA ACATCACCGCAGAACAGACAACGACAGCAGCCTCCGAC ACTCCCCCGCCACGACCGCAGCCGACCCCTAAAGCAG AGAACACCAACACGAGCAAGGGTACCGACCTCTGGACCC CGCCACCACAACAAGTCCCCAAAACACAGCGAGACCGCT GGCAACAACAACACTCATCACAAGATACCGGAGAAGAGA GTGCCAGCAGCGGGAAGCTAGGCTTAATTACCAATACTATT GCTGGAGTCGCAGGACTGATCACAGGCGGGAGGAGAGCTC GAAGAGAAGCAATTGTCAATGCTCAACCCAAATGCAACCC TAATTTACATTACTGGACTACTCAGGATGAAGGTGCTGCAA TCGGACTGGCCTGGATACCATAATTCGGGCCAGCAGCCGA GGGAATTTACATAGAGGGGCTGATGCACAATCAAGATGGT TTAATCTGTGGTTGAGACAGCTGGCCAACGAGACGACTC AAGCTCTTCAACTGTTCTGAGAGCCACAACCGAGCTACGC ACCTTTTCAATCCTCAACCGTAAGGCAATTGATTTCTTGCT GCAGCGATGGGGCGGCACATGCCACATTTGGGACCGGAC TGCTGTATCGAACCACATGATTGGACCAAGAACATAACAG ACAAAATTGATCAGATTATTATGATTGTTGTGATAAAACC CTTCCGGACCGGGGACAAATGACAATTGGTGGACAGGAT GGAGACAATGGATACCGGCAGGTATTGGAGTTACAGGCGT TATAATTGCAGTTATCGCTTTATTCTGTATATGCAAAATTTGT CTTTTAG
80 Short WPRE sequence	AATCAACCTCTGGATTACAAAATTTGTGAAAGATTGACTGA TATTTCTTAATATGTTGCTCCTTTTACGCTGTGTGGATATGC TGCTTTAATGCCTCTGTATCATGCTATTGCTTCCCGTACGGC TTTCGTTTTCTCCTCCTGTATATAATCCTGGTTGCTGTCTCTT TATGAGGAGTTGTGGCCGTTGTCCGTCAACGTGGCGTGGT GTGCTCTGTGTTTGTGACGCAACCCCACTGGCTGGGGCA TTGCCACCACCTGTCAACTCCTTTCTGGGACTTTCGCTTTCC CCCTCCCGATCGCCACGGCAGAACTCATCGCCGCTGCCTT GCCCCTGCTGGACAGGGCTAGGTTGCTGGGCACGTGATA ATTCGCTGGTGTGTC
81 Primer	TAAGCAGAATTCATGAATTTGCCAGGAAGAT
82 Primer	CCATACAATGAATGGACACTAGGCGGCCGCACGAAT
83 Gag, Pol, Integrase fragment	GAATTCATGAATTTGCCAGGAAGATGGAACCAAAAATGA TAGGGGAATTGGAGGTTTTATCAAAGTAAGACAGTATGA TCAGATACTCATAGAAATCTGCGGACATAAAGCTATAGGT ACAGTATTAGTAGGACCTACACCTGTCAACATAAATTGGAA GAAATCTGTGACTCAGATTGGCTGCACTTTAAATTTTCCC ATTAGTCTATTGAGACTGTACCAGTAAAATTAAAGCCAGG AATGGATGGCCCAAAGTTAAACAATGGCCATTGACAGAA GAAAAATAAAAGCATTAGTAGAAAATTTGTACAGAAATGG AAAAGGAAGGAAAAATTTCAAAAATTGGCCCTGAAAATCC ATACAATACTCCAGTATTTGCCATAAAGAAAAAGACAGT ACTAAATGGAGAAAATTAGTAGATTTACAGAGAACTTAATA AGAGAACTCAAGATTTCTGGGAAGTTCAATTAGGAATACC ACATCCTGCAGGGTTAAAACAGAAAAAATCAGTAACAGTA CTGGATGTGGGCGATGCATATTTTCAGTTCCCTTAGATAA AGACTTCAGGAAGTATATGCAATTTACCATACTTAGTATAA ACAATGAGACACCGGGATTAGATATCAGTACAATGTGCT TCCACAGGATGGAAGGATCACCAGCAATATTCAGTGT

-continued

SEQ ID	No: Description	Sequence
		AGCATGACAAAAATCTTAGAGCCTTTTAGAAAAACAAATC CAGACATAGTCATCTATCAATACATGGATGATTGTATGTA GGATCTGACTTAGAAATAGGGCAGCATAGAACAAAAATAG AGGAACTGAGACAACATCTGTTGAGGTGGGGATTACCAC ACCAGACAAAAACATCAGAAAGAACCTCCATTCTTTGG ATGGGTTATGAACTCCATCTGTATAATGGACAGTACAGCC TATAGTGCTGCCAGAAAAGGACAGCTGGACTGTCAATGAC ATACAGAAATTAGTGGGAAAAATTGAATTGGGCAAGTCAGA TTTATGCAGGGATTAAAGTAAGGCAATTATGTAAACTTCTT AGGGGAACCAAAGCACTAACAGAAGTAGTACCCTAACAG AAGAAGCAGAGCTAGAATGGGAGAAAAAGGAGATTCT AAAAGAACCAGTACATGGAGTGATTATGACCCATCAAAA GACTTAATAGCAGAAATACAGAAGCAGGGGCAAGGCCAAT GGACATATCAAAATTTATCAAGAGCCATTTAAAAATCTGAA AACAGGAAAGTATGCAAGAATGAAGGGTCCCACTAAT GATGTGAAACAATTAAACAGAGGCAGTACAAAAATAGCCA CAGAAAGCATAGTAATATGGGAAAGACTCCTAAATTTAA ATTACCCATACAAAAGGAAACATGGGAAGCATGGTGGACA GAGTATTGGCAAGCCACCTGGATTCTGAGTGGGAGTTTGT CAATACCCCTCCCTTAGTGAAGTTATGGTACAGTTAGAGA AAGAACCATAATAGGAGCAGAACTTTCTATGTAGATGG GGCAGCCAATAGGGAACATAATAGGAAAGCAGGATAT GTAAGTGACAGAGGAAGCAAAAAGTTGTCCCTTAACGG ACACAACAATCAGAAGACTGAGTTACAAGCAATTCATCT AGCTTTGCAGGATTGGGATTAGAAGTAAACATAGTGACA GACTCACAATATGCATTGGGAATCATTCAAGCACAAACAG ATAAGAGTGAATCAGAGTTAGTCAGTCAATAATAGAGCA GTTAATAAAAAAGGAAAAAGTCTACCTGGCATGGGTACCA GCACACAAAGGAATTGGAGGAAATGAACAAGTAGATAAAT TGGTCAGTGCTGGAATCAGGAAAGTACTATTTTAGATGGA ATAGATAAGGCCCAAGAAGACATGAGAAATATCACAGTA ATTGGAGAGCAATGGCTAGTGATTTTAACTACCACCTGTA GTAGCAAAAGAAATAGTAGCCAGCTGTGATAAATGTCAGC TAAAAGGGGAAGCCATGCATGGACAAGTAGACTGTAGCCC AGGAATATGGCAGCTAGATTGTACACATTTAGAAGGAAAA GTTATCTTGGTAGCAGTTTATGTAGCCAGTGGATATATAGA AGCAGAAGTAATTCAGCAGAGACAGGGCAAGAAACAGC ATACTCTCTTAAATTAGCAGGAAGATGGCCAGTAAAA ACAGTACATACAGACAATGGCAGCAATTCACCACTACTA CAGTTAAGCCCGCTGTTGGTGGGCGGGATCAAGCAGGA ATTTGGCATTCCCTACAATCCCAAAGTCAAGGAGTAATAG AATCTATGAATAAAGAAATTAAGAAAAATTATAGGACAGGT AAGAGATCAGGCTGAACATCTTAAGACAGCAGTACAAATG GCAGTATTCATCCACAATTTTAAAGAAAAGGGGGATTG GGGGGTACAGTGCAGGGGAAAGAAATAGTAGACATAATAGC AACAGACATACAACTAAGAATTACAAAAACAAATTACA AAAATTCAAAAATTTCCGGTTTATTACAGGGACAGCAGAG ATCCAGTTTGGAAAGGACAGCAAGCTCCTCTGGAAGG TGAAGGGCAGTAGTAATACAAGATAATAGTGACATAAAA GTAGTGCCAAGAAGAAAAGCAAGATCATCAGGGATTATG GAAAACAGATGGCAGGTGATGATTGTGTGGCAAGTAGACA GGATGAGGATTAA
84	DNA Fragment containing Rev, RRE and rabbit beta globin poly A	TCTAGAATGGCAGGAAGAAGCGGAGACAGCGACGAAGAG CTCATCAGAACAGTCAGACTCATCAAGCTTCTCTATCAAAG CAACCCACCTCCCAATCCGAGGGGACCCGACAGGCCCCGA AGGAATAGAAGAAGAAGGTGGAGAGAGACAGAGACAG ATCCATTGATAGTGAACGGATCCTTGGCACTTATCTGGG ACGATCTGCGGAGCCTGTGCCTTTCAGCTACCACCGCTTG AGAGACTTACTCTGATTGTAACGAGGATTGTGGAACCTCT GGGACGCAAGGGGTGGGAAGCCCTCAAATATTGGTGGAAAT CTCCTACAATATTGGAGTCAGGAGCTAAAGAAATAGAGGAG CTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATG GGCGCAGCGTCAATGACGCTGACGGTACAGGCCAGACAAT TATTGTCTGGTATAGTGCAGCAGCAGAACAAATTTGCTGAGG GCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTG GGGCATCAAGCAGCTCCAGGCAAGAATCCTGGCTGTGGAA AGATACCTAAAGGATCAACAGCTCCTAGATCTTTTCCCTC TGCCAAAAATTTAGGGGACATCATGAAGCCCTTGAGCAT CTGACTTCTGGCTAATAAAGGAAATTTATTTTCATTGCAAT AGTGTGTTGGAATTTTGTGTCTCTCACTCGAAGGACAT ATGGGAGGGCAATCATTTAAACATCAGAAAGAGTATTT

-continued

SEQ ID NO: Description	Sequence
	GGTTTAGAGTTTGGCAACATATGCCATATGCTGGCTGCCAT GAACAAAGGTGGCTATAAGAGGTCATCAGTATATGAAAC AGCCCCCTGCTGTCCATTCTTATTCCATAGAAAAGCCTTG ACTTGAGGTTAGATTTTTTTTATATTTTGTGTTATTTT TTTCTTTAACATCCCTAAAATTTTCCTTACATGTTTACTAG CCAGATTTTTCCTCTCTCTGACTACTCCAGTCATAGCTG TCCCTCTTCTCTTATGAAGATCCCTCGACCTGCAGCCCAAG CTTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAAATT GTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG CATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAA CTCACATTAAATGCGTTGCGCTCACTGCCCGCTTCCAGTC GGGAAACCTGTGCTGCCAGCGGATCCGCATCTCAATTAGTC AGCAACCATAGTCCCGCCCCCTAATCCGCCCCATCCCGCCCC TAACTCCGCCCCAGTTCCGCCCCATTCTCGCCCCATGGCTGA CTAATTTTTTTTATTATGACAGAGCCGAGGCCCTCGGC CTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAG GCCTAGGCTTTTGCAAAAAGCTAAGTTGTTTATTGCAGCTT ATAATGGTTACAAATAAGCAATAGCATCACAAATTTAC AAATAAAGCATTTTTTCACTGCATTCTAGTTGTGGTTGTG CAAATCATCAATGTATCTATCAGCGGCCGCCCGGG
85 DNA fragment containing the CAG enhancer/ promoter/intron sequence	ACGCGTTAGTTATTAATAGTAATCAATTACGGGGTCATTAG TTCATAGCCCATATATGGAGTTCGCGTTACATAACTTACG GTAAATGGCCCCGCTGGCTGACCGCCCAACGACCCCCGCC CATTGACGTCATAATGACGTATGTTCCCATAGTAACGCCA ATAGGGACTTTCCATTGACGTCAATGGGTGGACTATTTACG GTAAACTGCCCACTTGGCAGTACATCAAGTGATCATATGC CAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCC GCCTGGCATTATGCCCAGTACATGACCTTATGGGACTTTCC TACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCA TGGGTCGAGGTGAGCCCCACGTTCTGCTTCACTCTCCCCAT CTCCCCCCCCCTCCCCACCCCCAATTTGTATTATTTATTTT TTAATTATTTTGTGACGATGGGGCGGGGGGGGGGGGG GCGCGCGCCAGGCGGGGCGGGGCGGGGCGAGGGGCGGGG CGGGGCGAGGCGGAGAGGTGCGGCGGCGAGCCAAATCAGAG CGGCGCGCTCCGAAAGTTTCTTTTATGGCGAGGCGCGGC GGCGGCGGCCCTATAAAAAGCGAAGCGCGCGGCGGGCGG GAGTCGCTGCGTTGCTTTCGCCCCGTGCCCGCTCCGCGCC GCCTCGCGCGCCCCCGCGCTCTGACTGACCGGCTTACT CCCACAGGTGAGCGGGCGGGACGGCCCTTCTCCTCCGGGC TGTAATTAGCGCTTGGTTAATGACGGCTCGTTCTTTTCTG TGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCCCTT TGTGCGGGGGGAGCGGCTCGGGGGGTGCGTGCCTGTGTG TGTGCTGGGGAGCGCGCTGCGGCGCGCGCTGCCCGGC GGCTGTGAGCGCTGCGGGCGCGGCGCGGGCTTGTGCGC TCCGCGTGTGCGCGAGGGGAGCGCGGCGGGGCGGTGCC CCGCGGTGCGGGGGGCTGCGAGGGGAACAAAGGCTGCGT GCGGGGTGTGTGCTGGGGGGGTGAGCAGGGGTGTGGC GCGGCGGTGCGGCTGTAAACCCCCCTGCACCCCCCTCCCC GAGTTGCTGAGCACGGCCCGCTTCGGGTGCGGGGCTCCG TGCGGGCGTGGCGCGGGCTCGCCGTGCGGGCGGGGG TGGCGGCAAGTGGGGTGCCGGGCGGGGCGGGCGCCCTC GGGCCGGGAGGGCTCGGGGAGGGGCGCGCGGCCCCG GAGCGCCGGCGGCTGTGAGGCGCGGCGAGCCGAGCCAT TGCTTTTATGGTAATCGTGAGAGGGGCGAGGACTTCC TTTGTCCCAATCTGGCGAGCCGAAATCTGGGAGGCGCC GCCGACCCCCCTAGCGGGCGCGGCGAAGCGGTGCGGC GCCGCGAGGAAGGAAATGGGCGGGGAGGGCTTCGTGCGT CGCCGCGCGCGCTCCCTTCTCCATCTCCAGCTCGGGG TGCCGCGAGGGGAGGCTGCTTCGGGGGGGACGGGGCAG GGCGGGGTTCCGCTTCTGCGGTGTGACCGGCGGGAATTC
86 DNA fragment containing VSV-G	GAATTCATGAAGTGCCTTTTGTACTTAGCCTTTTATTCATT GGGGTGAATTGCAAGTTCACCATAGTTTTTCCACACAACCA AAAAGGAACTGGAAAAATGTTCTTCTAATACCATTTATT GCCCGTCAAGCTCAGATTAAATTTGGCATAATGACTTAATA GGCACAGCCTTACAAGTCAAAATGCCAAGAGTCACAAGG CTATTCAAGCAGACGGTTGGATGTGTCATGCTTCCAAATGG GTCATACTTGTGATTTCGCTGGTATGGACCGAAGTATAT AACACATTCCATCCGATCCTTCACTCCATCTGTAGAACAA GCAAGGAAAGCATTGAACAAACGAACAGGAAGTGGCT GAATCCAGGCTTCCCTCCTCAAGTTGTGGATATGCAACTG

-continued

SEQ ID NO: Description	Sequence
	TGACGGATGCCGAAGCAGTGATTGTCCAGGTGACTCCTCAC CATGTGCTGGTTGATGAATACACAGGAGAATGGGTTGATTCT ACAGTTCATCAACGGAATGCAGCAATTACATATGCCCC ACTGTCCATAACTCTACAACCTGGCATTCTGACTATAAGGT CAAAGGGCTATGTGATTCTAACCTCATTTCCATGGACATCA CCTTCTTCTCAGAGGACGGAGAGCTATCATCCTGGGAAG GAGGGCACAGGGTTCAGAAGTAACACTTTGCTTATGAAA CTGGAGGCAAGGCCCTGCAAAATGCAATACTGCAAGCATTG GGGAGTCAGACTCCCATCAGGTGTCTGGTTCGAGATGGCTG ATAAGGATCTCTTTGCTGCAGCCAGATTCCCTGAATGCCCA GAAGGGTCAAGTATCTCTGCTCCATCTCAGACCTCAGTGGAA TGTAAGTCTAATTCAGGACGTTGAGAGGATCTTGATTATT CCCTCTGCCAAGAAACCTGGAGCAAAATCAGAGCGGGTCT TCCAATCTCTCCAGTGGATCTCAGCTATCTTGCTCCTAAAA ACCCAGGAACCGGTCTGCTTTTACCATAATCAATGGTACC CTAAAATACTTTGAGACCAGATACATCAGAGTCGATATTGC TGCTCCAATCCTCTCAAGAAATGGTCGGAATGATCAGTGGAA CTACCACAGAAAGGGAACGTGGGATGACTGGGCACCAT TGAAGACGTGGAATGGACCAATGGAGTTCTGAGGACC AGTTCAGGATATAAGTTTCTTTATACATGATTGGACATGG TATGTTGGACTCCGATCTTCATCTTAGCTCAAAGGCTCAGG TGTTCGAACATCCTCACATTCAAGACGCTGCTTCGCAACTT CCTGATGATGAGAGTTTATTTTTTGGTGATACTGGGCTATC CAAAAATCCAATCGAGCTTGTAAGGTTGGTTCAGTAGTT GGAAAAGCTCTATTGCCCTCTTTTTCTTTATCATAGGGTTAA TCATTGGACTATCTTGGTTCTCCGAGTTGGTATCCATCTTT GCATTAAATTAAAGCACCAAGAAAAGACAGATTTATAC AGACATAGAGATGAGAATTC
87 Helper plasmid containing RRE and rabbit beta globin poly A	TCTAGAAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAG GAAGCACTATGGGCGCAGCGTCAATGACGCTGACGGTACA GGCCAGACAATATTGTCTGGTATAGTGCAGCAGCAGAAC AATTTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCA ACTCACAGTCTGGGCGCATCAAGCAGCTCCAGGCAAGAATC CTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTAG ATCTTTTCCCTCTGCCAAAATTATGGGGACATCATGAAG CCCCTTGAGCATCTGACTTCTGGCTAATAAAGGAAATTTAT TTTCATTGCAATAGTGTGTTGGAATTTTTTGTGCTCTCTCACT CGGAAGGACATATGGGAGGGCAATCATTAAACATCAG AATGAGTATTTGGTTTAGAGTTTGGCAACATATGCCATATG CTGGCTGCCATGAACAAAGGTGGCTATAAAGAGGTCATCA GTATATGAAACAGCCCCCTGCTGTCCATTCTTATTCCTATA GAAAAGCCTTGACTTGAGGTTAGATTTTTTTTATATTTTGT TTGTGTTATTTTTTCTTTAATACCTTAAATTTTCTTAC ATGTTTACTAGCCAGATTTTCTCTCTCTGACTACTCC CAGTCATAGCTGTCCCTCTTCTCTTATGAAGATCCCTCGAC CTGCAGCCCAAGCTTGGCGTAATCATGGTCATAGCTGTTTC CTGTGTGAAATTGTTATCCGCTCACAATCCACACAACATA CGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAAT GAGTGAGCTAACTACATTAATTCGCTGCGCTCACTGCC GCTTTCCAGTCGGGAAACCTGTCGTGCCAGCGGATCCGCAT CTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCC CATCCCGCCCCCTAACTCCGCCAGTTCCGCCCATCTCCGC CCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAG GCCGCCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAG GCTTTTTTGGAGGCCCTAGGCTTTTGCAAAAAGCTAACTTGT TTATTGCAGCTTATAATGGTTACAATAAAGCAATAGCATC ACAAATTTACAAAATAAAGCATTTTTTTTACATGCATTCTAG TTGTGGTTTGTCCAAACTCATCAATGTATCTTATACCCGG G
88 RSV promoter and HIV Rev	CAATTGCGATGTACGGGCCAGATATACGCGTATCTGAGGG GACTAGGGTGTGTTTAGGCGAAAAGCGGGGCTTCGGTTGT ACGCGGTTAGGAGTCCCCCAGGATATAGTAGTTTCGCTTT TGCATAGGGAGGGGAAATGTAGTCTTATGCAATACACTT GTAGTCTTGCAACATGGTAACGATGAGTTAGCAACATGCCT TACAAGGAGAGAAAAGCACCGTGATGCGGATTGGTGGA AGTAAGGTGGTACGATCGTGCCTTATTAGGAAGGCAACAG ACAGGTCTGACATGGATTGGACGAACCACTGAATTCGCA TTGCAGAGATAATTGTATTTAAGTGCCCTAGCTCGATACAAT AAACGCCATTTGACCATTACCACATTGGTGTGCACCTCCA AGCTCGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGAC

-continued

SEQ ID NO: Description	Sequence
	GCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGAC CGATCCAGCCTCCCCTCGAAGCTAGCGATTAGGCATCTCCT ATGGCAGGAAGAAGCGGAGACAGCGACGAAGAACTCCTC AAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC ACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAAGGAAT AGAAGAAGAAGGTGGAGAGAGACAGAGACAGATCCAT TCGATTAGTGAACGGATCCTTAGCACTTATCTGGGACGATC TGCGGAGCCTGTGCCTCTTCAGCTACCACCGCTTGAGAGAC TTACTCTTGATTGTAACGAGGATTGTGGAACCTCTGGGACG CAGGGGGTGGGAAGCCCTCAAATATTGGTGGAATCTCCTA CAATATTGGAGTCAGGAGCTAAAGAATAGTCTAGA
89 Target sequence	ATGGCAGGAAGAAGCGGAG
90 shRNA sequence	ATGGCAGGAAGAAGCGGAGTTCAAGAGACTCCGCTTCTTC CTGCCATTTTTT
91 H1 promoter and shRT sequence	GAACGCTGACGTCATCAACCCGCTCCAAGGAATCGCGGGC CCAGTGTCACTAGGCGGGAAACACCCAGCGCGCTGCGCCC TGGCAGGAAGATGGCTGTGAGGGACAGGGGAGTGGCGCCC TGCAATATTTGCATGTCGCTATGTGTTCTGGGAAATCACCA TAAACGTGAAATGTCTTTGGATTGGGAATCTTATAAGTTC TGTATGAGACCACTTGGATCCGCGAGACAGCGACGAAGA GCTTCAAGAGAGCTCTTCGTGCTGTCTCCGCTTTTT
92 H1 CCR5 sequence	GAACGCTGACGTCATCAACCCGCTCCAAGGAATCGCGGGC CCAGTGTCACTAGGCGGGAAACACCCAGCGCGCTGCGCCC TGGCAGGAAGATGGCTGTGAGGGACAGGGGAGTGGCGCCC TGCAATATTTGCATGTCGCTATGTGTTCTGGGAAATCACCA TAAACGTGAAATGTCTTTGGATTGGGAATCTTATAAGTTC TGTATGAGACCACTTGGATCCGCTGTCAAGTCCAATCTATGT TCAAGAGACATAGATTGGACTTGACACTTTTT
93 Primer	AGGAATTGATGGCGAGAAGG
94 Primer	CCCCAAAGAAGGTCAAGGTAATCA
95 Primer	AGCGCGGCTACAGCTTCA
96 Primer	GGCGACGTAGCACAGCTTCT
97 AGT103 CCR5 miR30	TGTAAACTGAGCTTGCTCTA
98 AGT103-R5- 1	TGTAAACTGAGCTTGCTCGC
99 AGT103-R5- 2	CATAGATTGGACTTGACAC
100 CAG promoter	TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATA GCCCCATATATGGAGTTCGCGTTACATAACTTACGGTAAAT GGCCCGCTGGCTGACCGCCCAACGACCCCGCCCATTTGAC GTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGG ACTTTCCATTGACGTCAATGGGTGGACTATTTACGGTAAAC TGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTA CGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCTGG CATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTG GCAGTACATCTACGTATTAGTCATCGCTATTACCATGGGTC GAGGTGAGCCCCACGTTCTGCTTCACTCTCCCATCTCCCC CCCCCTCCCAACCCCAATTTGTATTTATTTATTTTAAATT ATTTTGTGCAGCGATGGGGGCGGGGGGGGGGGGCGCGC GCCAGGCGGGGCGGGGCGGGGCGAGGGGCGGGGCGGGG GAGGCGGAGAGGTGCGGCGGCGAGCCAATCAGAGCGGCGC GCTCCGAAAGTTTCCTTTATGGCGAGGCGGCGGCGGCGG GGCCCTATAAAAAGCGAAGCGCGCGCGGGGCG
101 H1 element	GAACGCTGACGTCATCAACCCGCTCCAAGGAATCGCGGGC CCAGTGTCACTAGGCGGGAAACACCCAGCGCGCTGCGCCC TGGCAGGAAGATGGCTGTGAGGGACAGGGGAGTGGCGCCC TGCAATATTTGCATGTCGCTATGTGTTCTGGGAAATCACCA

-continued

SEQ ID NO: Description	Sequence
	TAAACGTGAAATGCTTTGGATTGGGAATCTTATAAGTTC TGTATGAGACCACTT
102 3' LTR	TGGAAGGGCTAATTCACCTCCCAACGAAGATAAGATCTGCTT TTTGCTTGTACTGGGTCTCTCTGGTTAGACCAGATCTGAGC CTGGGAGCTCTCTGGCTAACTAGGGAACCACTGCTTAAGC CTCAATAAAGCTTGCCCTTGAGTGCTTCAAGTAGTGTGTGCC CGTCTGTTGTGTGACTCTGGTAACTAGAGATCCCTCAGACC CTTTTAGTCAGTGTGGAAATCTCTAGCAGTAGTAGTTCAT GTCA
103 7SK promoter	CTGCAGTATTTAGCATGCCCCACCCATCTGCAAGGCATTCT GGATAGTGTCAAAACAGCCGGAATCAAGTCCGTTTATCTC AAACTTTAGCATTTTGGGAATAAATGATATTTGCTATGCTG GTAAATTAGATTTTAGTTAAATTTCTGTCTGAAGCTCTAG TACGATAAGCAACTTGACCTAAGTGTAAAGTTGAGATTTCC TTCAGGTTTATATAGCTTGTGCGCCGCTGGCTACCTC
104 miR155 Tat	CTGGAGGCTTGCTGAAGGCTGTATGCTGTCCGCTTCTTCC GCCATAGGGTTTGGCCACTGACTGACCCATGGGGGAAGA AGCGGACAGGACACAAGGCTGTACTAGCACTCACATGG AACAAATGGCC
105 pRSV Rev	AGCGCCCAATACGCAACCGCCTCTCCCGCGCGTTGGCCG ATTCAATTAATGCAGCTGGCAGCAGGTTTCCCGACTGGAA AGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTC ACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGC TCGTATGTTGTGTGGAATTGTGAGCGGATAACAATTTCA CAGGAAACAGCTATGACCATGATTACGAATTCATGTACG GGCCAGATATACGCGTATCTGAGGGGACTAGGGTGTGTTT AGGCGAAAAGCGGGGCTTCGGTTGTACGCGTTAGGAGTC CCCTCAGGATATAGTAGTTCGCTTTTGCATAGGGAGGGGG AAATGTAGTCTTATGCAATACACTTGTAGTCTTGCAACATG GTAACGATGAGTTAGCAACATGCCTTACAAGGAGAGAAAA AGCACCGTGATGCGCGATTGGTGGAAGTAAGGTGGTACGA TCGTGCCCTTATTAGGAAGGCAACAGACAGGTCTGACATGG ATTGGACGAACCACTGAATTCGCGATTGCAGAGATAATTGT ATTTAAGTGCCTAGCTCGATACAATAAACGCCATTTGACCA TTCACCACATTGGTGTGACCTCCAAGCTCGAGCTCGTTTA GTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTT TGACCTCCATAGAAGACCCGGGACCGATCCAGCCTCCCT CGAAGCTAGTCGATTAGGCATCTCCTATGGCAGGAAGAAG CGGAGACAGCGACGAAGACCTCCTCAAGGCAGTCAGACTC ATCAAGTTTCTCTATCAAGCAACCCACCTCCCAATCCCGA GGGGACCCGACAGGCCCGAAGGAATAGAAGAAGAAGGTG GAGAGAGAGACAGAGACAGATCCATTGATTAGTGAACGG ATCCTTAGCACTTATCTGGGACGATCTGCGGAGCCTGTGCC TCTTCAGCTACCACCGCTTGAGAGACTTACTCTGATTGTA ACGAGGATTGTGGAACCTTCTGGGACGAGGGGTGGGAAG CCCTCAAAATATTGGTGGAATCTCCTACAATATTGGAGTCAG GAGCTAAAGAATAGTGTGTAGCTTGCTCAATGCCACAGC TATAGCAGTAGCTGAGGGGACAGATAGGGTTATAGAAGTA GTACAAGAAGCTTGGCACTGGCCGTCGTTTACATGATCTG AGCCTGGGAGATCTCTGGCTAACTAGGGAACCACTGCTTA AGCCTCAATAAAGCTTGCCCTTGAGTGCTTCAAGTAGTGTGT GCCCCTCTGTTGTGTGACTCTGGTAACTAGAGATCACAAAG CAACCATAGTACGCGCCCTGTAGCGGCGCATTAAAGCGCGG CGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCC AGCGCCCTAGCGCCCGCTCCTTTCGCTTCTTCCCTTCTCTT CTCGCCACGTTTCGCGGCTTTCGCCGTCAGCTCTAAATCG GGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACC TCGACCCCAAAAACTTGATTTGGGTGATGGTTCACGTAGT GGGCCATCGCCCTGATAGACGGTTTTTTCGCCCTTTGACGTT GGAGTCCACGTTCTTAAATAGTGGACTCTTGTTCCAACTG GAACAACACTCAACCTATCTCGGGCTATCTTTTGTATTTA TAAGGGATTTTCCGATTTTCGGCCTATTGGTTAAAAAATGA GCTGATTTAACAAAAATTTAACGCGAATTTTAAACAAATAT TAACGTTTACAATTTTATGGTGCACCTCTCAGTACAATCTGC TCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAAC ACCCGCTGACGCGCCCTGACGGGCTGTCTGCTCCCGCAT CCGCTTACAGACAAGCTGTGACCGTCTCGGGAGCTGCATG TGTGAGAGGTTTTCACCGTCATACCCGAACCGCGGAGAC

-continued

SEQ ID NO: Description	Sequence
	GAAAGGGCCTCGTGATACGCCTATTTTTATAGGTAAATGTC ATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCG GGGAAATGTGCGCGGAACCCCTATTGTATTATTTTCTAAA TACATTCAAATATGTATCCGCTCATGAGACAATAACCCCTGA TAAATGCTTCAATAATATTGAAAAAGGAAGATATGAGTA TTCAACATTTCGGTGTGCGCCTATTCCCTTTTTTTCGGCAT TTTGCCCTTCCTGTTTTTGCTCAGCCAGAAACGCTGGTGAAA GTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGT ACATCGAAGTGGATCTCAACAGCGGTAAGATCCTTGAGAG TTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTA AAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCC GGCAAGAGCAACTCGGTGCGCGCATACACTATTCTCAGA ATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTT ACGGATGGCATGACAGTAAGAGAATTATGTCAGTGTGCCA TAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACA ACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGACA ACATGGGGGATCATGTAACGCGCTTGATCGTTGGGAACCG GAGCTGAATGAAGCCATACCAACGACGAGCGTGACACCA CGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACTATT AACTGGCGAAGTACTTACTCTAGCTTCCCGGCAACAATTAA TAGACTGGATGGAGCGGATAAAGTTGACAGGACCACTTCT GCGCTCGGCCCTTCGCGCTGGCTGGTTTTATTGCTGATAAAT CTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGC ACTGGGGCCAGATGGTAAGCCCTCCGTATCGTAGTTATCT ACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAG ACAGATCGCTGAGTAGGTGCGCTCACTGATTAAGCATTGGT AACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGAT TTAAAACCTTCATTTTTTAATTTAAAAGGATCTAGGTGAAGAT CCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGT TTTCGTTCCACTGAGCGTCAGACCCGTAGAAAAGATCAAA GGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGC TTGCAAAACAAAAAACCCCGCTACAGCGGTGGTTTGTGTT GCCGGATCAAGAGCTACCAACTCTTTTCCGAAGGTAAGT GCTTCAGCAGAGCGCAGATACCAAACTGTCCTTCTAGTG TAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACC GCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTG CTGCCAGTGGCGATAAGTCGTGCTTACCAGGTGGACTCA AGACGATAGTTACCGGATAAGGCGCAGCGGTGGGCTGAA CGGGGGTTTCGTGCACACAGCCAGCTTGAGCGAACGAC CTACACCGAAGTGAATACCTACAGCGTGAGCTATGAGAA AGCGCCACGCTTCCCGAAGGGAGAAAGCGGACAGGTATC CGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGG AGCTTCCAGGGGAAACGCCCTGGTATCTTTATAGTCCTGTC GGGTTTCGCACCTCTGACTTGAGCGTCGATTTTGTGATG CTCGTCAGGGGGCGGAGCTTATGAAAACGCCAGCAAC GCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCT CACATGTTCTTTCCGTGCGTTATCCCTGATTCTGTGGATAAC CGTATTACCGCCTTTGAGTGAGCTGATACCGCTCGCCGAG CCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGC GGAAG
106 pCMV-VSV- G	GAGCTTGGCCCATTCATACGTTGTATCCATATCATAATAT GTACATTTATATTGGCTCATGTCCAACATTACCGCATGTT GACATTGATTATTGACTAGTTATTAATAGTAATCAATTACG GGGTCATTAGTTATAGCCCATATATGGAGTTCCGCGTTAC ATAACTTACGGTAAATGGCCGCGCTGGCTGACCGCCCAAC GACCCCGCCCATTTGACGTCAATAATGACGTATGTTCCCAT AGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTG GAGTATTACGGTAACTGCCCACTTGGCAGTACATCAAGT GTATCATATGCCAAGTACGCCCTATTGACGTCAATGACG GTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTA TGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCAT CGCTATTACCATGGTGATGCGGTTTTTGGCAGTACATCAATG GGCGTGGATAGCGGTTTGACTCACGGGGATTTCGAAGTCTC CAGCCCATTTGACGTCAATGGGAGTTTGTTTTGGCACCAAAA TCAACGGGACTTTCCAAATGTCTGAACAACCTCGCCCCAT TGACGCAATGGGCGGTAGGCGGTACGGTGGGAGGTCTA TATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGA GACGCCATCCACGCTGTTTGGACCTCCATAGAAGACACCGG GACCGATCCAGCCTCCGGTCGACCGATCCTGAGAAGTTTACG GGTGAATTTGGGACCTTGATTGTTCTTTCTTTTCGCTAT TGTAATTCATGTTATATGGAGGGGCAAGTTTTTCAGGG

-continued

SEQ ID NO: Description	Sequence
	TGTGTGTTTAGAATGGGAAGATGTCCCTTGTATCACCATGGA CCCTCATGATAATTTGTCTTCTTCACTTCTACTCTGTTGA CAACCATTGTCTCCTCTTATTTCTTTTCACTTTCTGTAACCT TTTCGTTAACTTTAGCTTGCATTTGTAACGAATTTTAAAT TCACCTTTGTTTATTGTGAGATTGTAAGTACTTTCTCTAAT CACTTTTTTTTCAAGGCAATCAGGGTATATTATATTGTACTT CAGCACAGTTTTAGAGAACAAATGTTATAATTAAATGATAA GGTAGAATATTTCTGCATATAAATTCTGGCTGGCGTGGAAA TATTCTTATTGGTAGAAACAACCTACACCTGGTCATCATCC TGCCTTTCTCTTATGTTTACAATGATATACACTGTTTGAGA TGAGGATAAAATACTCTGAGTCCAAACCGGGCCCCCTGCT AACCATGTTTCATGCCTTCTCTCTTCTACAGCTCCTGGGC AACGTGCTGGTTGTTGTGCTGTCTCATCATTTTGGCAAAGA ATTCCTCGACGGATCCGCCATGAAGTGCCTTTTGTACTTAG CCTTTTTATTCTATTGGGGTGAATTGCAAGTTCAACATAGTTT TTCCACACAACCAAAAGGAACTGGAAAAATGTTCTCTTC TAATTACCATTTATGCCCGTCAAGCTCAGATTAAATTTGGC ATAATGACTTAATAGGCACAGCCTTACAAGTCAAAATGCC CAAGAGTCACAAGGCTATTCAAGCAGACGGTTGGATGTGT CATGCTTCCAAATGGGTCACTACTTGTGATTTCCGCTGGTA TGGACCGAAGTATATAACACATTCATCCGATCCTTCATC CATCTGTAGAACAATGCAAGGAAAGCATTGAACAAACGAA ACAAGGAACCTTGGCTGAATCCAGGCTTCCCTCCTCAAAGTT GTGGATATGCAACTGTGACGGATGCCGAAGCAGTGATTGT CCAGGTGACTCCTCACCATGTGCTGGTTGATGAATACACAG GAGAATGGGTGATTACAGTTTCATCAACGGAATATGCAG CAATTACATATGCCCCACTGTCCATACTCTACAACCTGGC ATTCTGACTATAAGGTCAAAGGGCTATGTGATTCTAACCTC ATTTCCATGGACATCACCTTCTCTCAGAGGACGGAGAGCT ATCATCCCTGGGAAAGGAGGGCACAGGGTTCAAGTAAC TACTTTGCTTATGAACTGGAGGCAAGGCTGCAAAATGC AATCTGCAAGCATTGGGGAGTCAGACTCCCATCAGGTGT CTGGTTCGAGATGGCTGATAAGGATCTCTTGTGTCAGCCA GATTCCCTGAATGCCCAGAAGGGTCAAGTATCTCTGCTCCA TCTCAGACCTCAGTGGATGTAAGTCTAATTCAGGACGTGA GAGGATCTTGGATTATTCCTCTGCCAAGAACTGGAGCA AAATCAGAGCGGGTCTTCCAATCTCTCAGTGGATCTCAGC TATCTTGCTCCTAAAAACCCAGGAACCGGTCTGCTTTTAC CATAATCAATGGTACCTTAAATACTTTGAGACCAGATACA TCAGAGTCGATATTGCTGCTCCAATCCTCTCAAGAATGGTC GGAATGATCAGTGGAACTACCACAGAAAGGGAACGTGGG ATGACTGGGCACCATATGAAGCGTGGAAATTGGACCCAA TGGAGTTCTGAGGACCAGTTCAAGGATATAAGTTTCTTTAT ACATGATTGGACATGGTATGTTGGACTCCGATCTTCTATCT AGCTCAAAGGCTCAGGTGTTTGAACATCTCAGATTCAAGA CGCTGCTTCGCAACTTCTGATGATGAGAGTTATTTTTTGG TGATACTGGGTATCCAAAAATCCAAATCGAGCTGTAGAA GGTGGTTCAGTAGTTGGAAGCTCTATTGCTCTTTTTTC TTTATCATAGGGTTAATCATTTGACTATTCTTGGTTCTCCGA GTTGGTATCCATCTTTGCATTAATTAAGCACACCAAGAA AAGACAGATTTATACAGACATAGAGATGAACCGACTTGA AAGTGATAAGGATCCGTCGAGGAATTCATCTCTCAGGTGC AGGCTGCCATCAGAAGGTGGTGGCTGGTGTGGCCAATGC CCTGGCTCACAATAACCACTGAGATCTTTTCCCTCTGCCA AAAATTATGGGACATCATGAAGCCCTTGAGCATCTGACT TCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTG TTGGAATTTTTTGTCTCTCACTCGGAAGGACATATGGGA GGGCAAAATCATTTAAACATCAGAAAGATTTTGGTTTAG AGTTTGGCAACATATGCCCATATGCTGGCTGCCATGAACAA AGGTTGGCTATAAAGAGGTCATCAGTATATGAACAGCCC CCTGCTGTCCATTCCTTATTCATAGAAAAGCCTTGACTTG AGGTTAGATTTTTTTTATATTTGTTTGTGTTATTTTTTCT TTAACATCCCTAAAAATTTTCTTACATGTTTACTAGCCAGA TTTTTCTCTCTCTGACTACTCCAGTCATAGCTGTCCCT CTTCTCTTATGGAGATCCCTCGACGGATCGGCCGAATTCG TAATCATGTATAGCTGTTTCTGTGTGAAATTTGTATCCGC TCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTG TAAAGCCTGGGGTGCTAATGAGTGAGCTAACTCACATTA ATTGCGTTGCGCTCACTGCCCGCTTTCAGTCGGGAAACCT GTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGG AGAGGCGGTTTGGCTATTGGGCGCTCTTCGCTTCTCGCT CACTGACTCGCTGCGCTCGGTCGTTCCGCTGCGGCGAGCGG

-continued

SEQ ID NO: Description	Sequence
	TATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGA ATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGG CCAGCAAAAGGCCAGGAACCGTAAAGGCGCGTTGCTG GCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACA AAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGA CTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGT GCGCTCTCCTGTTCCGACCCCTGCCGCTTACCGGATACCTGT CCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGC TCACGCTGTAGGTATCTCAGTTCGTGTAGGTGCTCGCTC CAAGCTGGGCTGTGTGCACGAACCCCGCTTCAGCCCGACC GCTGCGCTTATCCGGTAACATCTGCTTGTAGTCCAACCCG GTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTA ACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGTACAGA GTTCTTGAAGTGGTGGCTAACTACGGCTACACTAGAAGA ACAGTATTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTT CGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAACAAACC ACCGCTGGTAGCGGTGGTTTTTTGTGTTGCAAGCAGCAGAT TACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTGATC TTTTCTACGGGCTGTGACGCTCAGTGAACGAAAACTCACG TTAAGGGATTTTGGTCATGAGATTATCAAAAGGATCTTCA CCTAGATCCTTTTAAATTAAAAATGAAGTTTAAATCAATC TAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATG CTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTC GTTTCATCCATAGTTGCGTGAATCCCGTCTGTAGATAACT ACGATACGGAGGGCTTACCATCTGGCCCGAGTGTGCAA TGATACCGCGAGACCCACGCTACCGGCTCCAGATTATCA GCAATAAACGAGCCAGCCGGAAGGCGGAGCGCAGAAGT GGTCCTGCAACTTTATCCGCTCCATCCAGTCTATTAATTGT TGCCGGGAAGCTAGAGTAAGTAGTTCCGCGAGTTAATAGTTT GCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTAC GCTCGTCTGTTGGTATGGCTTCATTGAGTCCGCTTCCCAA CGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAA AAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAGT AAGTTGGCCGAGTGTATCACTCATGGTTATGGCAGCACT GCATAATCTCTTACTGTATGCCATCCGTAAGATGCTTTTC TGTGACTGGTGAGTACTCAACCAAGTCACTTGAGAATAGT GTATGCGGCGACCGAGTTGCTCTTGGCCGCGTCAATACGG GATAATACCGCGCCACATAGCAGAACTTAAAGTGCTCA TCATTGGAACGTTCTTCGGGCGAAACTCTCAAGGATC TTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGC ACCCAACTGATCTTCAGCATCTTTTACTTTCACCGCGTTTC TGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAA GGGAATAAGGGCGACACGGAATGTTGAATACTCATACTC TTCCTTTTCAATATATTGAAGCATTATCAGGGTTATTGT CTCATGAGCGGATACATATTGAATGATTTAGAAAAATAA ACAAATAGGGGTTCCGCGCACATTTCCCGAAAAGTGCCA CCTAAATGTGAAGCGTTAATATTTTGTAAAAATTCGCGTTA AATTTTGTAAATCAGCTCATTTTTTAAACCAATAGGCCGA AATCGGCAAAATCCCTTATAAATCAAAAGAAATAGACCGAG ATAGGGTTGAGTGTGTTCCAGTTTGGAAACAAGATCCACT ATTAAAGAACGTGGACTCCAACGTCAAAGGGCGAAAAACC GTCTATCAGGGCGATGGCCCACTACGTGAACCATCACCTTA ATCAAGTTTTTGGGGTTCAGGTGCGGTAAAGCACTAAATC GGAACCTTAAAGGGAGCCCCGATTAGAGCTTGACGGGG AAAGCCGGCGAACGTGGCGAGAAAGGAAGGAAGAAAGC GAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTC ACGCTGCGCGTAACCAACACACCCCGCGCTTAATGCGCC GCTACAGGGCGCGTCCCATTCGCGATTACGGTGTGCGCACT GTTGGGAAGGGCGATCGGTGCGGGCTCTTCGCTATTACGC CAGCTGGCGAAAGGGGATGTGCTGCAAGGCGATTAAAGTT GGGTAACGCGAGGGTTTTCCAGTCACGACGTTGTAAAC GACGGCCAGTGAGCGCGCTAATACGACTCACTATAGGGC GAATTGGAGCTCCACCGCGTGGCGGCGCTCTAGA
107 PSPAX2 delta Rev	GTCGACATTGATTATTGACTAGTTATTAATAGTAATCAATT ACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGT TACATAACTTACGGTAAATGGCCCGCTGGCTGACGCCCA ACGACCCCGCCCATTTGACGTCAATAATGACGTATGTTCCC ATAGTAACGCCAATAGGGACTTTCATTGACGTCAATGGGT GGACTATTTACGGTAAATGCCCACTTGGCAGTACATCAAG TGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGAC GGTAAATGGCCCGCTGGCATATTGCCCGAGTACATGACCTT

-continued

SEQ ID NO: Description	Sequence
	ATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCA TCGCTATTACCATGGGTCGAGGTGAGCCCCACGTTCTGCTT CACTCTCCCATCTCCCCCCTCCCCACCCCAATTTTGTA TTTATTTATTTTAAATTATTTTGTGCAGCGATGGGGCGGG GGGGGGGGGGCGCGCGCCAGGCGGGCGGGCGGGCG AGGGCGGGGGCGGGCGAGGCGGAGAGGTGCGGCGGCAG CCAATCAGAGCGGCGCGCTCCGAAAGTTTCCTTTTATGGCG AGGCGGCGGGCGGGCGGGCTATAAAAAGCGAAGCGCGC GGCGGGCGGGAGTCTGCTGCGTTGCTTCGCCCGTGCCCG CTCCGCGCGCGCTCGCGCGCGCCCGCCCGGCTCTGACTGAC CGCGTTACTCCACAGGTGAGCGGGCGGGACGGCCCTTCTC CTCCGGGCTGTAATTAGCGCTTGTTTAAATGACGGCTCGTT TCTTTTCTGTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGA GGGCCCTTTGTGCGGGGGGAGCGGCTCGGGGGGTGCGTG CGTGTGTGTGTGCGTGGGGAGCGCGCGTGCGGCCGCGC TGCCCGGCGGCTGTGAGCGCTGCGGGCGGGCGCGGGGCT TTGTGCGCTCCGCGTGTGCGCGAGGGGAGCGCGCGGGG GCGGTGCCCGCGGTGCGGGGGGCTGCGAGGGGAACAAA GGCTGCGTGCGGGGTGTGTGCGTGGGGGGTGAAGAGGG GTGTGGGCGCGCGGTGCGGGCTGAACCCCCCTGCACCC CCCTCCCCGAGTTGCTGAGCACGGCCCGGCTTCGGGTGCGG GGCTCCGTGCGGGGCGTGCGCGGGGCTCGCGTGCCGGG CGGGGGGTGGCGGCGAGGTGGGGGTGCCGGGCGGGCGGG GCCGCTCGGGCGGGGAGGGCTCGGGGAGGGCGCGGC GGCCCCGAGCGCGCGGCGCTGTGAGGCGCGGCGAGCCG CAGCCATTGCTTTTATGTAATCGTGCGAGAGGGCGCAGG GACTTCCTTTGTCCAAATCTGGCGGAGCCGAAATCTGGGA GGCGCGCGCGCACCCCTCTAGCGGGCGGGCGAAGCGG TGCGCGCGCGGAGGAAAGAAATGGCGGGGAGGGCCTTC GTGCGTTCGCGCGCGCGCGCTCCCTTCTCATCTCCAGCCT CGGGGCTGCGCGAGGGGACGGCTGCCCTCGGGGGGACG GGGCGGGGCGGGGTTCGGCTTCTGCGGTGTGACGGCGGC TCTAGAGCCTCTGCTAACCATGTTTATGCTTCTTTTCTT CTACAGCTCCTGGCAACGTGCTGGTTATTTGTGCTGTCTCA TCATTTTGGCAAAGAATTGCGGCCGCGCGCTTGACGCGCA CGGCAAGAGGCGAGGGCGCGCACTGGTGAGAGATGGGT GCGAGAGCGTCAGTATTAAGCGGGGAGAATTAGATCGAT GGGAAAAAATTCGTTAAGGCGAGGGGAAAGAAAAAAT ATAAATTAAACATATAGTATGGGCAAGCAGGGAGCTAGA ACGATTTCGAGTTAATCCTGGCCTGTAGAAACATCAGAAG GCTGTAGACAAATACTGGGACAGCTACAACTTCCCTTCA GACAGGATCAGAAGAACTTAGATCATTATATAATACAGTA GCAACCTCTATTGTGTGCATCAAAGGATAGAGATAAAG ACACCAAGGAAGCTTTAGACAAGATAGAGGAAGAGCAA ACAAAAGTAAGAAAAAGCACAGCAAGCAGCAGCTGACA CAGGACACAGCAATCAGGTCAGCCAAAATTACCTATAGT GCAGAACATCCAGGGGCAAAATGGTACATCAGGCCATATCA CCTAGAACTTTAAATGCATGGGTAAAAGTAGTAGAAGAGA AGGCTTTCAGCCAGAAGTGATACCCATGTTTTCAGCATTA TCAGAAAGGAGCACCCCAAGATTAAACACCATGCTAA ACACAGTGGGGGACATCAAGCAGCCATGCAAAATGTTAAA AGAGACCATCAATGAGGAAGCTGCAGAAATGGGATAGAGTG CATCCAGTGATGCAGGGCTTATGCACCGAGCCAGATGA GAGAACCAAGGGGAAGTGACATAGCAGGAATACTAGTAC CCTTCAGGAACAAATAGGATGGATGACACATAATCCACCT ATCCAGTAGGAGAAATCTATAAAGATGGATAATCCTGG GATTAAATAAAATAGTAAGAATGTATAGCCCTACCAGCAT TCTGGACATAAGACAAGGACCAAAGGAACCTTTAGAGAC TATGTAGACCGATTCTATAAACTCTAAGAGCCGAGCAAG CTTCACAAAGAGTAATAAATGGATGACAGAAACCTTGT GGTCCAAATGCGAACCCAGATTGTAAGACTATTTTAAA GCATTGGGACCAAGGAGCGACACTAGAAGAAATGATGACAG CATGTCAGGGAGTGGGGGACCCGGCCATAAAGCAAGAGT TTTGGCTGAAGCAATGAGCCAAGTAACAAATCCAGCTACC ATAATGATACAGAAAGGCAATTTTAGGAACCAAAGAAAGA CTGTTAAGTGTTCAATTGTGGCAAGAGGGCACATAGCC AAAAATTGCAAGGCCCTTAGGAAAAAGGGCTGTGGAAT GTGGAAGGAAGGACACCAATGAAAGATTGTAAGTGTGAGAG ACAGGCTAATTTTTAGGGAAGATCTGGCTTCCACCAAGG GAAGGCCAGGAATTTCTTCAGAGCAGACAGAGCCCAAC AGCCCCACAGAAGAGAGCTTCAGGTTTGGGAAGAGACA ACAACCTCCCTCTCAGAAGCAGGAGCCGATAGACAAGGAAC

-continued

SEQ ID NO: Description	Sequence
	TGTATCCTTTAGCTTCCCTCAGATCACTCTTTGGCAGCGACC CCTCGTCACAATAAAGATAGGGGGGCAATTAAAGGAAGCT CTATTAGATACAGGAGCAGATGATACAGTATTAGAAGAAA TGAATTTGCCAGGAAGATGGAACCAAAAATGATAGGGGG AATTGGAGGTTTTATCAAAGTAGGACAGTATGATCAGATA CTCATAGAATCTGCGGACATAAAGCTATAGGTACAGTATT AGTAGGACCTACACCTGTCAACATAAATTGGAAGAAATCTG TTGACTCAGATTGGCTGCACTTAAATTTCCCATTAGTCCT ATTGAGACTGTACCAGTAAAATTAAAGCCAGGAATGGATG GCCCCAAAGTTAAACAATGGCCATTGACAGAAGAAAAAT AAAAGCATTAGTAGAAATTTGTACAGAAATGGAAGGAA GGAAAAATTTCAAAAATTGGGCTGAAATCCATACAATA CTCCAGTATTTGCCATAAAGAAAAAGACAGTACTAAATG GAGAAAAATTAGTAGATTTTACAGAACTTAATAAGAGAACT CAAGATTTCTGGGAAGTTCAATTAGGAATACCACATCCTGC AGGGTTAAACAGAAAAATCAGTAACAGTACTGGATGTG GGCGATGCATATTTTTCAGTTCCTTAGATAAAGACTTCAG GAAGTATACTGCATTTACCATACCTAGTATAAACATGAGA CACCAGGGATTAGATATCAGTACAATGTGCTTCCACAGGG ATGGAAGGATCACCAGCAATATTCAGTGTAGCATGACA AAAACTTAGAGCCTTTTAGAAAACAAAATCCAGACATAG TCATCTATCAATACATGGATGATTTGTATGTAGGATCTGAC TTAGAAATAGGGCAGCATAGAACAAAAATAGAGGAAGTGA GACAACTCTGTTGAGGTGGGATTACCACACCAGACAA AAAACATCAGAAAGAACCTCCATTCTTTGGATGGGTATG AACTCCATCCTGATAAATGGACAGTACAGCCTATAGTGCTG CCAGAAAAGGACAGCTGGACTGTCAATGACATACAGAAAT TAGTGGGAAAAATGAATTGGGCAAGTCAGATTTATGCAGG GATTAAGTAAGGCAATTATGTAACCTCTTAGGGGAACC AAAGCACTAACAGAGTAGTACCCTAACAGAAAGAGCAG AGCTAGAACTGGCAGAAAAACAGGGAGATTCTAAAAGAAC GGTACATGGAGTGTATTATGACCCATCAAAAGACTTAATA GCAGAAATACAGAAAGCAGGGGCAAGGCCAATGGACATATC AAATTTATCAAGAGCCATTTAAAAATCTGAAAAACAGGAAA ATATGCAAGAAATGAAGGGTGCCACACTAATGATGTGAAA CAATTAACAGAGGCAGTACAAAAATAGCCACAGAAAGCA TAGTAATATGGGAAAGACTCCTAAATTTAAATTACCCATA CAAAAGGAACATGGGAAGCATGGTGGACAGAGTATTGGC AAGCCACTGGATTCTGAGTGGGAGTTTGTCAATACCCCT CCCTTAGTGAAGTTATGGTACCAGTTAGAGAAAGAACCCA TAATAGGAGCAGAACTTTCTATGTAGATGGGGCAGCCAA TAGGGAAACTAAATTAGGAAAAGCAGGATATGTAAGTAC AGAGGAAGACAAAAAGTTGTCCCTTAACGGACACAACAA ATCAGAAGACTGAGTTACAAGCAATTCATCTAGCTTTGCAG GATTCGGGATTAGAAGTAAACATAGTGACAGACTCACAAAT ATGCATTGGGAATCATTCAAGCACAAACAGATAAGAGTGA ATCAGAGTTAGTCAGTCAAAATATAGAGCAGTTAATAAAA AAGGAAAAAGTCTACCTGGCATGGGTACCAGCACACAAAG GAATTGGAGGAATGAACAAGTAGATGGGTGGTCAAGTGC TGGAATCAGGAAAGTACTATTTTATAGTGAATAGATAAG GCCCAAGAAGAACATGAGAAATATCACAGTAATTGGAGAG CAATGGCTAGTGATTTTAACCTACCACCTGTAGTAGCAAAA GAAATAGTAGCCAGCTGTGATAAATGTAGCTAAAGGGG AAGCCATGCATGGACAAGTAGACTGTAGCCAGGAATATG GCAGCTAGATTGTACACATTTAGAAGGAAAAGTTATCTTGG TAGCAGTTTCATGTAGCCAGTGGATATATAGAAGCAGAAAT AATTCCAGCAGAGACAGGGCAAGAAACAGCATACTTCCTC TTAAATTAGCAGGAAGATGGCCAGTAAAAACAGTACATA CAGACAATGGCAGCAATTCACCAGTACTACAGTTAAGGC CGCTGTGTGGTGGCGGGGATCAAGCAGGAATTTGGCATT CCCTACAATCCCCAAAGTCAAGGAGTAATAGAACTATGA ATAAAGAAATTAAGAAAAATTATAGGACAGGTAAAGATCA GGCTGAACATCTTAAGACAGCAGTACAAATGGCAGTATTC ATCCACAATTTTAAAGAAAAAGGGGGATTGGGGGTACA GTGCAGGGGAAAGAATAGTAGACATAATAGCAACAGACAT ACAAACTAAAGAATTACAAAAACAAATTACAAAAATTCAA AATTTTCGGGTTTATTACAGGACAGCAGAGATCCAGTTTG GAAAGGACCAGCAAAGCTCCTCTGGAAAGGTGAAGGGGCA GTAGTAATACAAGATAATAGTGACATAAAGTAGTGCCAA GAAGAAAAGCAAAGATCATCAGGGATTATGGAACACAGAT GGCAGGTGATGATTGTGTGGCAAGTAGACAGGATGAGGAT TAACACATGGAATTCGCAACACTGCTGTTTATCCATTTT

-continued

SEQ ID NO: Description	Sequence
	AGAAATGGAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGC AGGAAGCACTATGGGCGCAGCGTCAATGACGCTGACGGTA CAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGA ACAATTTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTG CAACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGAA TCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCT GGGGATTTGGGGTTGCTCTGGA AAACTCATTGCA CCACTG CTGTGCCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAA CAGATTTGGAATCACACGACCTGGATGGAGTGGGACAGAG AAATTAACAATTACACAAGCTTGCTAGCAGATCTTTTCCC TCTGCCAAAAATATGGGGACATCATGAAGCCCCCTTGAGC ATCTGACTTCTGGCTAATAAAGGAATTTATTTTCATTGCA ATAGTGTGTGGAAATTTTGTGTCTCTCACTCGGAAGGAC ATATGGGAGGGCAAATCATTAAAAACATCAGAATGAGTAT TTGGTTTAGAGTTTGGCAACATATGCCATATGCTGGCTGCC ATGAACAAAGGTGGCTATAAAGAGGTCATCAGTATATGAA ACAGCCCCCTGCTGTCCATTCTTATTCATAGAAAAGCCT TGACTTGAGGTTAGATTTTTTTTATATTTGTTTGTGTATT TTTTTCTTTAACATCCCTAAAAATTTCTTACATGTTTACT AGCCAGATTTTTCTCTCTCTCTGACTACTCCAGTCATAGC TGTCCTCTCTCTCTTATGAAGATCCCTCGACCTGCAGCCCA AGCTTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAAA TTGTTATCCGCTCACAATCCACACAACATACGAGCCGGAA GCATAAAGTGTAAGCCTGGGGTGCTAATGAGTGAGCTA ACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTCCAGT CGGGAACCTGTGCTGCCAGCGGATCCGCATCTCAATTAGT CAGCAACCATAGTCCCGCCCCTAACCTCGCCCATCCCGCCC CTAACTCCGCCAGTCCGCCCATTTCTCGCCCCATGGCTG ACTAATTTTTTTTATTTATGCAAGGCGGAGGCGCGCTCGG CCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGA GGCCTAGGCTTTTGCAAAAAGCTAACTGTTTATTGCAGCT TATAATGGTTACAAATAAAGCAATAGCATCAAAATTTTAC AAATAAAGCATTTTTTCTACTGCATCTAGTTGTGGTTGTG CAAACATCAATGTATCTTATCATGTCTGGATCCGCTGCA TTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGGT ATTGGGCGCTCTTCGCTTCTCGCTCACTGACTCGCTGCG CTCGGTCGTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAA AGGCGGTAAATCGGTTATCCACAGAATCAGGGGATAACGC AGGAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAG GAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGC TCCGCCCCCTGACGAGCATCAAAAAATCGACGCTCAAG TCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAG GCGTTTCCCCCTGGAAGCTCCCTCGTGGCTCTCTGTTCG ACCTGCGCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCG GGAAGCGTGGCGCTTTCTCAATGCTCAGCTGTAGGTATCT CAGTTCCGTTGTAGGTCGTTCCGCTCCAAGCTGGGCTGTGTG ACGAACCCCCGTTTACGCGGACCGCTGCGCTTATCCGGT AACTATCGTCTTGAGTCCAACCGGTAAGACACGACTTATC GCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCG AGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTGGTGGC CTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGC GCTCTGCTGAAGCCAGTTACCTTCGGA AAAAGAGTTGGTA GCTCTGTATCCGGCAAAACAAACACCGCTGGTAGCGGTGG TTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAG GATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGAC GCTCAGTGGAACGAAAACTCAGTTAAGGGATTTTGGTCAT GAGATTATCAAAAAGGATCTTACCTAGATCCTTTTAAAT AAAAATGAAGTTTTAAATCAATCTAAGTATATATGAGTA AACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCAC CTATCTCAGCGATCTGTCTATTTCTGTTATCCATAGTTGCCT GACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTT ACCATCTGGCCCCAGTGTGCAATGATACCGCGAGACCCA CGCTCACC GGCTCCAGATTTATCAGCAATAAACCGACGCAGC CGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTTATCC GCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGT AAGTAGTTTCGCGAGTTAATAGTTTGGCGAACGTTGTTGCCA TTGCTACAGGCATCGTGGTGTACGCTCGTCTGTTGGTATG GCTTCATTACAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCG GTCTCCGATCGTTGTGCAAGTAAGTTGGCGCAGTGTTA TCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT CATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACT

-continued

SEQ ID NO:	Description	Sequence
		CAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAG TTGCTCTTGCCCGCGCTCAATACGGGATAATACCGCGCCAC ATAGCAGAACTTTAAAAGTGCTCATCATTTGAAAACGTTCT TCGGGGCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATC CAGTTCGATGTAACCCACTCGTGACCCCAACTGATCTTCAG CATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAACA GGAAGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACA CGGAAATGTTGAATACTCATACTCTTCTTTTCAATATTAT TGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATACAT ATTTGAATGTATTTAGAAAAATAACAAATAGGGGTTCCG CGCACATTCCCGAAAAGTGCCACCTG

[0412] While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be

limited to such embodiments. Various modifications may be made thereto without departing from the scope and spirit of the present invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 107

<210> SEQ ID NO 1
 <211> LENGTH: 118
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: miR30 CCR5

<400> SEQUENCE: 1

aggtatattg ctgttgacag tgagcgactg taaactgagc ttgctctact gtgaagccac 60
 agatgggtag agcaagcaca gtttaccgct gcctactgcc tcggacttca aggggctt 118

<210> SEQ ID NO 2
 <211> LENGTH: 116
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: miR21 Vif

<400> SEQUENCE: 2

catctccatg gctgtaccac cttgtcggg gatgtgtact tctgaacttg tgttgaatct 60
 catggagttc agaagaacac atccgactg acattttgggt atctttcatc tgacca 116

<210> SEQ ID NO 3
 <211> LENGTH: 114
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: miR185 Tat

<400> SEQUENCE: 3

gggcctggct cgagcagggg gcgagggatt ccgcttcttc ctgccatagc gtggccccct 60
 cccctatggc aggcagaagc ggcaccttcc ctcccaatga ccgcgtcttc gtcg 114

<210> SEQ ID NO 4
 <211> LENGTH: 1104
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Elongation Factor-1 alpha (EF1-alpha) promoter

-continued

<400> SEQUENCE: 4

```

ccggtgccta gagaaggtgg cgcggggtaa actgggaaag tgatgtcgtg tactggctcc    60
gcctttttcc cgagggtggg ggagaaccgt atataagtgc agtagtcgcc gtgaacgttc    120
tttttcgcaa cggttttgcc gccagaacac aggtaagtgc cgtgtgtggt tccgcgggc    180
ctggcctctt tacgggttat ggcccttgcg tgccttgaat tacttccacg cccctggctg    240
cagtacgtga ttcttgatcc cgagcttcgg gttggaagtg ggtgggagag ttcgaggcct    300
tgcgcttaag gagecccttc gctcgtgct tgagttgagg cctggcctgg gcgctggggc    360
cgccgcgtgc gaatctgggt gcaccttcgc gcctgtctcg ctgctttcga taagtctcta    420
gccatttaaa atttttgatg acctgctgcg acgctttttt tctggcaaga tagtcttgta    480
aatgcggggc aagatctgca cactggtatt tcggtttttg gggccgcggg cggcgacggg    540
gcccgtgcgt cccagcgcac atgttcggcg aggcggggcc tgcgagcgcg gccaccgaga    600
atcggacggg ggtagtctca agctggcccg cctgctctgg tgccctggcct cgcgcgcggc    660
tgtatcgccc cgccctgggc ggcaaggctg gcccggtcgg caccagtgc gtgagcgga    720
agatggccgc tcccgggccc tgctgcaggg agctcaaaat ggaggacgcg gcgctcgga    780
gagcggggcg gtgagtcaac cacacaaagg aaaagggcct tccgtcctc agcgcgcgt    840
tcattgtgact ccacggagta ccgggcgcgg tccaggcacc tcgattagtt ctcgagcttt    900
tggagtacgt cgtctttagg ttggggggag gggttttatg cgatggagtt tccccacact    960
gagtgggtgg agactgaagt taggccagct tggcacttga tgtaattctc cttggaattt   1020
gccctttttg agtttggatc ttggttcatt ctcaagctc agacagtgg tcaaagtttt   1080
tttttccat ttcaggtgtc gtga                                           1104

```

<210> SEQ ID NO 5

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: CCR5 target sequence

<400> SEQUENCE: 5

```

gagcaagctc agtttaca                                           18

```

<210> SEQ ID NO 6

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Vif target sequence

<400> SEQUENCE: 6

```

gggatgtgta cttctgaact t                                           21

```

<210> SEQ ID NO 7

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Tat target sequence

<400> SEQUENCE: 7

```

tccgcttctt cctgccatag                                           20

```

-continued

<210> SEQ ID NO 8
<211> LENGTH: 126
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: TAR decoy sequence

<400> SEQUENCE: 8

cttgcaatga tgcgtaatt tgcgtcttac ctcgttctcg acagcgacca gatctgagcc 60
tgggagctct ctggctgtca gtaagctggg acagaagggt gacgaaaatt cttactgagc 120
aagaaa 126

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Rev/Tat target sequence

<400> SEQUENCE: 9

gcggagacag cgacgaagag c 21

<210> SEQ ID NO 10
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Rev/Tat shRNA sequence

<400> SEQUENCE: 10

gcggagacag cgacgaagag cttcaagaga gctcttcgtc gctgtctccg cttttt 56

<210> SEQ ID NO 11
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Gag target sequence

<400> SEQUENCE: 11

gaagaaatga tgacagcat 19

<210> SEQ ID NO 12
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Gag shRNA sequence

<400> SEQUENCE: 12

gaagaaatga tgacagcatt tcaagagaat gctgtcatca tttcttcttt tt 52

<210> SEQ ID NO 13
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Pol target sequence

<400> SEQUENCE: 13

caggagcaga tgatacag 18

-continued

<210> SEQ ID NO 14
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Pol shRNA sequence

<400> SEQUENCE: 14

caggagatga tacagttcaa gagactgtat catctgctcc tggtttt 47

<210> SEQ ID NO 15
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 target sequence #1

<400> SEQUENCE: 15

gtgtcaagtc caatctatg 19

<210> SEQ ID NO 16
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 shRNA sequence #1

<400> SEQUENCE: 16

gtgtcaagtc caatctatgt tcaagagaca tagattggac ttgacacttt tt 52

<210> SEQ ID NO 17
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 target sequence #2

<400> SEQUENCE: 17

gagcatgact gacatctac 19

<210> SEQ ID NO 18
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 shRNA sequence #2

<400> SEQUENCE: 18

gagcatgact gacatctact tcaagagagt agatgtcagt catgctcttt tt 52

<210> SEQ ID NO 19
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 target sequence #3

<400> SEQUENCE: 19

gtagctctaa caggttgga 19

<210> SEQ ID NO 20
<211> LENGTH: 52

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 shRNA sequence #3

<400> SEQUENCE: 20

gtagctctaa caggttggat tcaagagatc caacctgtta gagctacttt tt 52

<210> SEQ ID NO 21
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 target sequence #4

<400> SEQUENCE: 21

gttcagaaac tacctcttta 19

<210> SEQ ID NO 22
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 shRNA sequence #4

<400> SEQUENCE: 22

gttcagaaac tacctcttat tcaagagata agaggtagtt tctgaacttt tt 52

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 target sequence #5

<400> SEQUENCE: 23

gagcaagctc agtttacacc 20

<210> SEQ ID NO 24
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 shRNA sequence #5

<400> SEQUENCE: 24

gagcaagctc agtttacacc ttcaagagag gtgtaaactg agcttgctct tttt 54

<210> SEQ ID NO 25
<211> LENGTH: 141
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 gene, sequence 1

<400> SEQUENCE: 25

atggattatc aagtgtcaag tccaatctat gacatcaatt attatacatc ggagccctgc 60

caaaaaatca atgtgaagca aatcgagcc cgcctcctgc ctccgctcta ctcaactgggtg 120

ttcatctttg gttttgtggg c 141

<210> SEQ ID NO 26
<211> LENGTH: 633

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 gene, sequence 2

<400> SEQUENCE: 26

aacatgctgg tcatcctcat cctgataaac tgcaaaaggc tgaagagcat gactgacatc	60
tacctgctca acctggccat ctctgacctg ttttctcttc ttactgtccc cttctgggct	120
cactatgctg ccgcccagtg ggacttttga aatacaatgt gtcaactctt gacagggctc	180
tattttatag gcttctcttc tggaaatctc ttcacatccc tctgacaat cgataggtac	240
ctggctgtcg tccatgctgt gtttgcttta aaagccagga cggtcacctt tgggggtgtg	300
acaagtgtga tcacttgggt ggtggctgtg ttgctgtctc tcccaggaat catctttacc	360
agatctcaaa aagaaggctc tcattacacc tgcagctctc attttccata cagtcagtat	420
caattctgga agaatttcca gacattaaag atagtcactc tggggctggc cctgccgctg	480
cttgtcatgg tcactctgta ctcggaatc ctaaaaactc tgcttcgggt tcgaaatgag	540
aagaagaggc acagggtgt gaggttctc ttcaccatca tgattgttta tttctcttc	600
tgggctccct acaacattgt ccttctctg aac	633

<210> SEQ ID NO 27
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 gene, sequence 3

<400> SEQUENCE: 27

acctccagg aattcttttg cctgaataat tgcagtagct ctaacagggt ggaccaagct	60
atgcagggtga	70

<210> SEQ ID NO 28
<211> LENGTH: 140
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 gene, sequence 4

<400> SEQUENCE: 28

cagagactct tgggatgacg cactgctgca tcaaccccat catctatgcc ttgtcgggg	60
agaagtccag aaactacctc ttagtcttct tccaaaagca cattgccaaa cgttcttgca	120
aatgctgttc tattttccag	140

<210> SEQ ID NO 29
<211> LENGTH: 75
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CCR5 gene, sequence 5

<400> SEQUENCE: 29

caagaggctc ccgagcgagc aagctcagtt tacacccgat ccactgggga gcaggaaata	60
tctgtgggct tgtga	75

<210> SEQ ID NO 30
<211> LENGTH: 541

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CD4 promoter sequence

<400> SEQUENCE: 30

tggtggggtt caaatgtgag cccagctgt tagccctctg caaagaaaa aaaaaaaaaa    60
aaagaacaaa gggcctagat ttccctctg agccccacc taagatgaag cctctctttt    120
caagggagtg ggggtggggt ggaggcgat cctgtcagct ttgctctctc tgtggctggc    180
agtttctcca aagggttaaca ggtgtcagct ggctgagcct aggetgaacc ctgagacatg    240
ctacctctgt cttctcatgg ctggaggcag cctttgtaag tcacagaaag tagctgaggg    300
gctctggaag aaagacagcc aggggtggagg tagattggtc ttgactcct gatttaagcc    360
tgattctgct taactttttc ccttgacttt ggcattttca ctttgacatg ttccctgaga    420
gctggggggg tggggaaccc agctccagct ggtgacgttt ggggccggcc caggcctagg    480
gtgtggagga gccttgccat cgggcttctt gtctctcttc atttaagcac gactctgcag    540
a                                                                    541

```

```

<210> SEQ ID NO 31
<211> LENGTH: 359
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miR30-CCR5/miR21-Vif/miR185 Tat microRNA
cluster sequence

```

```

<400> SEQUENCE: 31

aggatatatt ctgttgacag tgagcgactg taaactgagc ttgctctact gtgaagccac    60
agatgggtag agcaagcaca gtttaccgct gcctactgcc tcggacttca aggggcttcc    120
cgggcatctc catggctgta ccaccttgtc gggggatgtg tacttctgaa cttgtgttga    180
atctcatgga gttcagaaga acacatccgc actgacattt tggatatctt catctgacca    240
gctagcgggc ctggctcgag caggggggga gggattccgc ttcttctgcg catagcgtgg    300
tcccctcccc tatggcaggc agaagcggca ccttccctcc caatgaccgc gtcttcgctc    359

```

```

<210> SEQ ID NO 32
<211> LENGTH: 590
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Long WPRE sequence

```

```

<400> SEQUENCE: 32

aatcaacctc tgattacaaa atttgtgaaa gattgactgg tattcttaac tatgttgctc    60
cttttacgct atgtggatac gctgctttta tgcctttgta tcatgctatt gcttcccgta    120
tggttttcat tttctctcc ttgtataaat cctggttgct gtctctttat gaggagtgtg    180
ggcccgttgt caggcaacgt ggcgtgggtg gcactgtgtt tgctgacgca acccccactg    240
gttggggcat tgccaccacc tgctagctcc ttccgggac ttctgcttcc cccctcccta    300
ttgccacggc ggaactcatc gccgcctgcc ttgcccgctg ctggacaggg gctcggtgtg    360
tgggcactga caattccgtg gtgttgctcg ggaaatcatc gtccttctct tggtgctcg    420
cctgtgttgc cacctggatt ctgcgcggga cgtccttctg ctacgtccct tcggccctca    480

```

-continued

atccagcggga ccttccctcc cgcggcctgc tgccggctct gcggcctctt ccgcgtcttc	540
gccttcgccc tcagacgagt cggatctccc ttggggcgc ctccccgcct	590

<210> SEQ ID NO 33
 <211> LENGTH: 1469
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Elongation Factor-1 alpha (EF1-alpha)
 promoter - miR30CCR5 - miR21Vif - miR185 Tat

<400> SEQUENCE: 33

ccggtgccta gagaaggtgg cgcggggtaa actgggaaag tgatgtcgtg tactggctcc	60
gcctttttcc cgagggtggg ggagaaccgt atataagtc agtagtcgcc gtgaacgttc	120
tttttcgcaa cgggtttgcc gccagaacac aggtaagtc cgtgtgtggt tcccgcgggc	180
ctggcctctt tacgggttat ggcccttcgc tgccctgaat tacttccacg cccctggctg	240
cagtacgtga ttcttgatcc cgagcttcgg gttggaagtg ggtgggagag ttcgaggcct	300
tgcgcttaag gagecccttc gcctcgtgct tgagttgagg cctggcctgg gcgctggggc	360
cgcgcgctgc gaatctggtg gcaccttcgc gcctgtctcg ctgctttcga taagtctcta	420
gccatttaaa atttttgatg acctgctgcg acgctttttt tctggcaaga tagtcttgta	480
aatgcggggc aagatctgca cactggtatt tcggtttttg gggccgcggg cggcgacggg	540
gcccgtgcgt cccagcgcac atgttcggcg aggcggggcc tgcgagcgcg gccaccgaga	600
atcgagcggg ggtagtctca agctggcccg cctgctctgg tgccctggcct cgcgcgcgcg	660
tgtatcgccc cgcctcgggc ggcaaggctg gcccggtcgg caccagttgc gtgagcggaa	720
agatggccgc tcccgcgcc tgctgcaggg agctcaaaat ggaggacgcg gcgctcggga	780
gagcgggcgg gtgagtcacc cacacaaagg aaaagggcct ttccgtcttc agcgcgtcgt	840
tcattgtgact ccacggagta ccgggcgcgcg tccaggcacc tcgattagtt ctcgagcttt	900
tggagtacgt cgtcttttag ttggggggag gggttttatg cgatggagtt tccccacact	960
gagtgggtgg agactgaagt taggccagct tggcacttga tgtaattctc cttggaattt	1020
gccctttttg agtttgatc ttggttcatt ctcaagctc agacagtggg tcaaagtttt	1080
tttcttccat ttcaggtgct gtgatgtaca aggtatattg ctgttgacag tgagcgactg	1140
taaactgagc ttgctctact gtgaagccac agatgggtag agcaagcaca gtttaccgct	1200
gcctactgcc tcggacttca aggggcttcc cgggcacttc catggctgta ccaccttgct	1260
gggggatgtg tacttctgaa cttgtgttga atctcatgga gttcagaaga acacatccgc	1320
actgacattt tggtatcttt catctgacca gctagcgggc ctggctcgag cagggggcga	1380
gggattccgc ttcttctgc catagcgtgg tcccctcccc tatggcaggc agaagcggca	1440
ccttccctcc caatgaccgc gtcttcgctc	1469

<210> SEQ ID NO 34
 <211> LENGTH: 228
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Rous Sarcoma virus (RSV) promoter

<400> SEQUENCE: 34

gtagtcttat gcaatactct tgtagtcttg caacatggta acgatgagtt agcaacatgc	60
---	----

-continued

cttacaagga gagaaaaagc accgtgcatg cggattgggtg gaagtaaggt ggtacgatcg 120
tgccttatta ggaaggcaac agacgggtct gacatggatt ggacgaacca ctgaattgcc 180
gcattgcaga gatattgtat ttaagtgcct agctcgatac aataaacg 228

<210> SEQ ID NO 35
<211> LENGTH: 180
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' Long terminal repeat (LTR)

<400> SEQUENCE: 35

ggtctctctg gttagaccag atctgagcct gggagctctc tggctaacta gggaaccac 60
tgcttaagcc tcaataaagc ttgccttgag tgcttcaagt agtgtgtgcc cgtctgttgt 120
gtgactctgg taactagaga tccctcagac ccttttagtc agtgtggaaa atctctagca 180

<210> SEQ ID NO 36
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Psi Packaging signal

<400> SEQUENCE: 36

tacgccaaaa attttgacta gcggaggcta gaaggagaga g 41

<210> SEQ ID NO 37
<211> LENGTH: 233
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Rev response element (RRE)

<400> SEQUENCE: 37

aggagctttg ttccttgggt tcttgggagc agcaggaagc actatgggcg cagcctcaat 60
gacgtgacg gtacaggcca gacaattatt gtctggata gtgcagcagc agaacaattt 120
gctgagggct attgaggcgc aacagcatct gttgcaactc acagtctggg gcatcaagca 180
gctccaggca agaatcctgg ctgtggaaag atacctaaag gatcaacagc tcc 233

<210> SEQ ID NO 38
<211> LENGTH: 118
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Central polypurine tract (cPPT)

<400> SEQUENCE: 38

ttttaaaaga aaagggggga ttggggggta cagtgcaggg gaaagaatag tagacataat 60
agcaacagac atacaaacta aagaattaca aaaacaaatt acaaaattca aaatttta 118

<210> SEQ ID NO 39
<211> LENGTH: 250
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' delta LTR

<400> SEQUENCE: 39

-continued

tggaagggtc aattcactcc caacgaagat aagatctgct ttttgcttgt actgggtctc	60
tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta	120
agcctcaata aagcttgctc tgagtgtctc aagtagtgtg tgcccgctcg ttgtgtgact	180
ctggttaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtagta	240
gttcatgtca	250

<210> SEQ ID NO 40
 <211> LENGTH: 352
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper/Rev - CMV early (CAG) enhancer - EnhanceTranscription

<400> SEQUENCE: 40

tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata tggagttccg	60
cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc ccgccccatt	120
gacgtcaata atgacgtatg ttcccatagt aacccaata gggactttcc attgacgtca	180
atgggtggac tatttacggt aaactgccc cttggcagta catcaagtgt atcatatgcc	240
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta	300
catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tc	352

<210> SEQ ID NO 41
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper/Rev - Chicken beta actin (CAG) promoter - Transcription

<400> SEQUENCE: 41

gctattacca tgggtcgagg tgagccccac gttctgcttc actctcccca tctccccccc	60
ctcccccccc ccaattttgt atttatattat tttttaatta ttttgtgcag cgatgggggc	120
gggggggggg ggggcgcgcg ccaggcgggg cggggcgggg cgaggggcgg ggcggggcga	180
ggcggagagg tgcggcgcca gccaatcaga gcggcgcgct ccgaaagttt ccttttatgg	240
cgaggcgggc gcggcgggcg ccctataaaa agcgaagcgc gcggcgggcg	290

<210> SEQ ID NO 42
 <211> LENGTH: 960
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper/Rev - Chicken beta actin intron - Enhance gene expression

<400> SEQUENCE: 42

ggagtgcgtg cgttgccctc gccccgtgcc ccgctccgcg ccgcctcgcg ccgcccgcgc	60
cggtctgac tgaccgcgtt actccacag gtgagcgggc gggacggccc ttctcctccg	120
ggctgtaatt agcgtttggt ttaatgacgg ctcgtttctt ttctgtgggt gcgtgaaagc	180
cttaaagggc tccgggaggg ccctttgtgc gggggggagc ggctcggggg gtgcgtgcgt	240
gtgtgtgtgc gtggggagcg ccgcgtgcgg cccgcgctgc ccggcggctg tgagcgctgc	300

-continued

gggcgcggcg cggggctttg tgcgctccgc gtgtgcgcga ggggagcgcg gccgggggcg	360
gtgccccgcg gtgcgggggg gctgcgaggg gaacaaaggc tgcgtgcggg gtgtgtgcgt	420
gggggggtga gcagggggtg tgggcgcggc ggtegggctg taaccccccc ctgcaccccc	480
ctccccgagt tgctgagcac ggccccgctt cgggtgcggg gctccgtgcg gggcgtggcg	540
cggggctcgc cgtgccgggc ggggggtggc ggcaggtggg ggtgccgggc ggggcggggc	600
cgcctcgggc cggggagggc tcgggggagg ggcgcggcgg ccccgagcg cggcggtcg	660
tcgaggcgcg gcgagccgca gccattgctt tttatggtaa tcgtgcgaga gggcgaggg	720
acttcctttg tcccaaatct ggcgagcg aaatctggga ggcgccgcg caccctctct	780
agcgggcgcg ggcgaagcgg tcgggcgcg gcaggaagga aatgggcggg gagggccttc	840
gtgcgtcgcc gcgccgcgt ccccttctcc atctccagcc tcggggctgc cgcaggggga	900
cggctgcctt cgggggggac ggggcagggc ggggttcggc ttctggcgtg tgaccggcg	960

<210> SEQ ID NO 43

<211> LENGTH: 1503

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Helper/Rev - HIV Gag - Viral capsid

<400> SEQUENCE: 43

atgggtgcga gacgctcagt attaagcggg ggagaattag atcgatggga aaaaattcgg	60
ttaaggccag ggggaaagaa aaaatataaa ttaaaacata tagtatgggc aagcagggag	120
ctagaacgat tcgcagttaa tcctggcctg ttagaaacat cagaaggctg tagacaaata	180
ctgggacagc tacaaccatc ccttcagaca ggatcagaag aacttagatc attatataat	240
acagtagcaa cctctattg tgtgcatcaa aggatagaga taaaagacac caaggaagct	300
ttagacaaga tagaggaaga gcaaaacaaa agtaagaaaa aagcacagca agcagcagct	360
gacacaggac acagcaatca ggtcagccaa aattacccta tagtgagaa catccagggg	420
caaatggtac atcaggccat atcacctaga actttaaatg catgggtaaa agtagtagaa	480
gagaaggctt tcagcccaga agtgataccc atgttttcag cattatcaga aggagccacc	540
ccacaagatt taaacaccat gctaaacaca gtggggggac atcaagcagc catgcaaatg	600
ttaaaagaga ccatcaatga ggaagctgca gaatgggata gagtgcattc agtgcattgca	660
gggcctattg caccaggcca gatgagagaa ccaaggggaa gtgacatagc aggaactact	720
agtacccttc aggaacaaat aggatggatg acacataatc cacctatccc agtaggagaa	780
atctataaaa gatggataat cctgggatta aataaaatag taagaatgta tagccctacc	840
agcattctgg acataagaca aggaccaaag gaacccttta gagactatgt agaccgattc	900
tataaaactc taagagccga gcaagcttca caagaggtaa aaaattggat gacagaaacc	960
ttgttggtcc aaaatgcgaa ccagattgt aagactattt taaaagcatt gggaccagga	1020
gcgacactag aagaaatgat gacagcatgt caggagtggt ggggacccgg ccataaagca	1080
agagtttttg ctgaagcaat gagccaagta acaaatccag ctaccataat gatacagaaa	1140
ggcaatttta ggaaccaaag aaagactgtt aagtgtttca attgtggcaa agaagggcac	1200
atagccaaaa attgcagggc ccctaggaaa aagggtgtgt ggaaatgtgt aaaggaagga	1260
caccaaatga aagattgtac tgagagacag gctaattttt tagggaagat ctggccttcc	1320

-continued

cacaaggga ggccaggga tttcttcag agcagaccag agccaacagc cccaccagaa	1380
gagagcttca gggttgaggga agagacaaca actccctctc agaagcagga gccgatagac	1440
aaggaaactgt atccttttagc ttccctcaga tcaactcttg gcagcgaccc ctggtcacia	1500
taa	1503

<210> SEQ ID NO 44
 <211> LENGTH: 1872
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper/Rev - HIV Pol - Protease and reverse transcriptase

<400> SEQUENCE: 44

atgaatttgc caggaagatg gaaacaaaa atgatagggg gaattggagg ttttatcaaa	60
gtaggacagt atgatcagat actcatagaa atctgaggac ataaagctat aggtacagta	120
ttagtaggac ctacacctgt caacataatt ggaagaaatc tgttgactca gattggctgc	180
actttaaatt ttcccattag tcctattgag actgtaccag taaaattaaa gccaggaatg	240
gatggcccaa aagttaaaca atggccattg acagaagaaa aaataaaagc attagtagaa	300
atttgtacag aaatggaaaa ggaaggaaaa atttcaaaaa ttgggcctga aaatccatac	360
aatactccag tatttgcct aaagaaaaaa gacagtacta aatggagaaa attagtagat	420
ttcagagaac ttaataagag aactcaagat ttctgggaag ttcaattagg aataccacat	480
cctgcagggt taaaacagaa aaaatcagta acagtactgg atgtgggcca tgcataat	540
tcagttccct tagataaaga cttcaggaag tatactgcat ttaccatacc tagtataaac	600
aatgagacac cagggttag atatacgtac aatgtgcttc cacagggatg gaaaggatca	660
ccagcaatat tccagtgtag catgacaaaa atcttagagc ctttttagaaa acaaaatcca	720
gacatagtca tctatcaata catggatgat ttgtatgtag gatctgactt agaaataggg	780
cagcatagaa caaaaataga ggaactgaga caacatctgt tgagggtggg atttaccaca	840
ccagacaaaa aacatcagaa agaactcca ttcccttgga tgggttatga actccatcct	900
gataaatgga cagtacagcc tatagtgtg ccagaaaagg acagctggac tgtcaatgac	960
atacagaaat tagtgggaaa attgaattgg gcaagtcaga tttatgcagg gattaaagta	1020
aggcaattat gtaaaacttct taggggaacc aaagcactaa cagaagtagt accactaaca	1080
gaagaagcag agctagaact ggcagaaaaac agggagattc taaaagaacc ggtacatgga	1140
gtgtattatg acccatcaaa agacttaata gcagaaatac agaagcaggg gcaaggccaa	1200
tggacatatc aaatttatca agagccattt aaaaatctga aaacaggaaa atatgcaaga	1260
atgaagggtg cccacactaa tgatgtgaaa caattaacag aggcagtaca aaaaatagcc	1320
acagaaagca tagtaatatg gggaaagact cctaaattta aattacccat acaaaaggaa	1380
acatgggaag catggtggac agagtattgg caagccacct ggattcctga gtgggagttt	1440
gtcaataccc ctcccttagt gaagttagg taccagttag agaaagaacc cataatagga	1500
gcagaaactt tctatgtaga tggggcagcc aatagggaaa ctaaattagg aaaagcagga	1560
tatgtaactg acagaggaag acaaaaagtt gtccccctaa cggacacaac aaatcagaag	1620
actgagttac aagcaattca tctagctttg caggattcgg gattagaagt aaacatagtg	1680
acagactcac aatatgcatt gggaaatcatt caagcacacac cagataagag tgaatcagag	1740

-continued

```

ttagtcagtc aaataataga gcagttaata aaaaaggaaa aagtctacct ggcatgggta 1800
ccagcacaca aaggaattgg aggaaatgaa caagtagatg ggttggtcag tgctggaatc 1860
aggaaagtac ta 1872

```

```

<210> SEQ ID NO 45
<211> LENGTH: 867
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper Rev - HIV Integrase - Integration of
viral RNA

```

```

<400> SEQUENCE: 45

```

```

tttttagatg gaatagataa ggcccaagaa gaacatgaga aatatcacag taattggaga 60
gcaatggcta gtgatttttaa cctaccacct gtagtagcaa aagaaatagt agccagctgt 120
gataaatgtc agctaaaagg ggaagccatg catggacaag tagactgtag ccaggaata 180
tggcagctag attgtacaca tttagaagga aaagttatct tggtagcagt tcatgtagcc 240
agtggatata tagaagcaga agtaattcca gcagagacag ggcaagaaac agcatacttc 300
ctcttaaaat tagcaggaag atggccagta aaaacagtac atacagacaa tggcagcaat 360
ttcaccagta ctacagttaa ggccgcctgt tgggtggcgg ggatcaagca ggaatttggc 420
attccctaca atcccaaag tcaaggagta atagaatcta tgaataaaga attaaagaaa 480
attataggac aggtgaagaga tcagggtgaa catcttaaga cagcagtaca aatggcagta 540
ttcatccaca attttaaaag aaaagggggg attggggggg acagtgcagg ggaaagaata 600
gtagacataa tagcaacaga catacaaact aaagaattac aaaaacaaat tacaaaaatt 660
caaaattttc gggtttatta cagggacagc agagatccag tttggaaagg accagcaaag 720
ctcctctgga aaggtgaagg ggcagtagta atacaagata atagtacat aaaagtagtg 780
ccaagaagaa aagcaaagat catcagggat tatggaaaac agatggcagg tgatgattgt 840
gtggcaagta gacaggatga ggattaa 867

```

```

<210> SEQ ID NO 46
<211> LENGTH: 234
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper/Rev - HIV RRE- Binds Rev element

```

```

<400> SEQUENCE: 46

```

```

aggagctttg ttccttgggt tcttgggagc agcaggaagc actatgggag cagcgtcaat 60
gacgctgacg gtacaggcca gacaattatt gtctggtata gtgcagcagc agaacaattt 120
gctgagggct attgaggcgc aacagcatct gttgcaactc acagtctggg gcatcaagca 180
gctccaggca agaatcctgg ctgtggaaag atacctaaag gatcaacagc tcct 234

```

```

<210> SEQ ID NO 47
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper/Rev - HIV Rev - Nuclear export and
stabilize viral mRNA

```

```

<400> SEQUENCE: 47

```

-continued

atggcaggaa gaagcggaga cagcgacgaa gaactcctca aggcagtcag actcatcaag 60
tttctctatc aaagcaaccc acctcccaat cccgagggga cccgacaggc ccgaaggaat 120
agaagaagaa ggtggagaga gagacagaga cagatccatt cgattagtga acggatcctt 180
agcacttata tgggacgata tgcggagcct gtgcctcttc agctaccacc gcttgagaga 240
cttactcttg attgtaacga ggattgtgga acttctggga cgcagggggg gggaagccct 300
caaataattg tggaatctcc tacaataattg gagtcaggag ctaaagaata g 351

<210> SEQ ID NO 48
<211> LENGTH: 448
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper/Rev - Rabbit beta globin poly A - RNA
stability

<400> SEQUENCE: 48

agatcttttt ccctctgcca aaaattatgg ggacatcatg aagccccttg agcatctgac 60
ttctggctaa taaaggaaat ttattttcat tgcaatagtg tgttggaatt ttttgtgtct 120
ctcactcgga aggacatatg ggagggcaaa tcatttaaaa catcagaatg agtatttggt 180
ttagagtgtg gcaacatatg ccatatgctg gctgccatga acaaaggtgg ctataaagag 240
gtcatcagta tatgaaacag cccctgctg tccattcctt attccataga aaagccttga 300
cttgaggtta gatttttttt atattttgtt ttgtgttatt tttttcttta acatccctaa 360
aattttcctt acatgtttta ctagccagat ttttcctcct ctctgacta ctcccagtca 420
tagctgtccc tcttctctta tgaagata 448

<210> SEQ ID NO 49
<211> LENGTH: 352
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper - CMV early (CAG) enhancer -
Enhancetranscription

<400> SEQUENCE: 49

tagttattaa tagtaatcaa ttacggggtc attagtcatg agcccatata tggagttccg 60
cgttacataa cttacggtaa atggcccgcc tggctgacgg cccaacgacc ccgcccatt 120
gacgtcaata atgacgtatg tccccatagt aacgccaata gggactttcc attgacgtca 180
atgggtggac tatttacggg aaactgccc cttggcagta catcaagtg atcatatgcc 240
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgcccagta 300
catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tc 352

<210> SEQ ID NO 50
<211> LENGTH: 290
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Helper - Chicken beta actin (CAG) promoter -
Transcription

<400> SEQUENCE: 50

gctattacca tgggtcgagg tgagcccccac gttctgcttc actctcccca tctccccccc 60

-continued

ctcccccccc ccaattttgt atttatttat tttttaatta ttttgtgcag cgatgggggc	120
gggggggggg ggggcgcgcg ccaggcgggg cggggcgggg cgaggggcgg ggcggggcga	180
ggcggagagg tgcggcgcca gccaatcaga gcggcgcgct ccgaaagttt ccttttatgg	240
cgaggcggcg gcggcggcgg ccctataaaa agcgaagcgc gcggcggggc	290

<210> SEQ ID NO 51
 <211> LENGTH: 960
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper - Chicken beta actin intron - Enhance gene expression

<400> SEQUENCE: 51

ggagtcgctg cgttgccctt gccccgtgcc ccgctccgcg ccgcctcgcg ccgcccgccc	60
cggtcttgac tgaccgcggt actccacag gtgagcgggc gggacggccc ttctcctccg	120
ggctgtaatt agcgttgggt ttaatgacgg ctcgtttctt ttctgtggct gcgtgaaagc	180
cttaaggggc tccgggaggg ccttttgtgc gggggggagc ggctcggggg gtgcgtgcgt	240
gtgtgtgtgc gtggggagcg ccgcgtgcgg ccgcgcgtgc ccggcggctg tgagcgtgc	300
gggcgcggcg cggggccttg tgcgctccgc gtgtgcgcga ggggagcgcg gccggggggc	360
gtgccccgcg gtgcgggggg gctgcgaggg gaacaaagcg tgcgtgcggg gtgtgtgcgt	420
gggggggtga gcaggggggt tgggcgcggc ggtcgggctg taaccccccc ctgcaccccc	480
ctccccgagt tgctgagcac ggccccgctt cgggtgcggg gctccgtgcg gggcgtggcg	540
cggggctcgc cgtgccgggc ggggggtggc ggcaggtggg ggtgccgggc ggggcggggc	600
cgctccgggc cggggagggc tcgggggagg ggcgcggcgg ccccgagcgg ccggcggctg	660
tcgaggcgcg gcgagccga gccattgcct tttatggtaa tcgtgcgaga gggcgcaggg	720
acttcctttg tcccaaatct ggccgagcgg aaatctggga ggcgccgcgg cccccctct	780
agcgggcgcg ggcgaagcgg tgcggcgcgg gcaggaagga aatgggcggg gagggccttc	840
gtgcgtgcgc gcgcgcgct cccctctccc atctccagcc tcggggctgc cgcaggggga	900
cggctgcctt cggggggggc ggggcagggc ggggttcggc ttctggcgtg tgaccggcgg	960

<210> SEQ ID NO 52
 <211> LENGTH: 1503
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper - HIV Gag - Viral capsid

<400> SEQUENCE: 52

atgggtgcga gagcgtcagt attaagcggg ggagaattag atcgatggga aaaaattcgg	60
ttaaggccag ggggaaagaa aaaatataaa ttaaaacata tagtatgggc aagcaggag	120
ctagaacgat tcgcagtaa tcctggcctg ttagaaacat cagaagcgtg tagacaaata	180
ctgggacagc tacaaccatc ccttcagaca ggatcagaag aacttagatc attatataat	240
acagtagcaa ccctctattg tgtgcatcaa aggatagaga taaaagacac caagggaagct	300
ttagacaaga tagaggaaga gcaaaacaaa agtaagaaaa aagcacagca agcagcagct	360
gacacaggac acagcaatca ggtcagccaa aattacccta tagtgcagaa catccagggg	420
caaatggtac atcaggccat atcacctaga actttaaatg catgggtaaa agtagtagaa	480

-continued

gagaaggctt tcagcccaga agtgataccc atgttttcag cattatcaga aggagccacc	540
ccacaagatt taaacacccat gctaaacaca gtggggggac atcaagcagc catgcaaattg	600
ttaaagaga ccatcaatga ggaagctgca gaatgggata gagtgcaccc agtgcacatga	660
gggcctattg caccaggcca gatgagagaa ccaaggggaa gtgacatagc aggaactact	720
agtacccttc aggaacaaat aggatggatg acacataatc cacctatccc agtaggagaa	780
atctataaaa gatggataat cctgggatta aataaaatag taagaatgta tagccctacc	840
agcattcttg acataagaca aggaccaaag gaacccttta gagactatgt agaccgattc	900
tataaaactc taagagccga gcaagcttca caagaggtaa aaaattggat gacagaaacc	960
ttgttggtcc aaaatgcgaa cccagattgt aagactatct taaaagcatt gggaccaggga	1020
gagacactag aagaaatgat gacagcatgt caggagatgg ggggacccgg ccataaagca	1080
agagtttttg ctgaagcaat gagccaagta acaaatccag ctaccataat gatacagaaa	1140
ggcaatttta ggaaccaaag aaagactgtt aagtgtttca attgtggcaa agaagggcac	1200
atagccaaaa attgcagggc cctaggaaa aagggtgtgt ggaaatgtgg aaaggaagga	1260
caccaaataa aagattgtac tgagagacag gctaattttt tagggaagat ctggccttcc	1320
cacaagggaa ggccagggaa ttttcttcag agcagaccag agccaacagc cccaccagaa	1380
gagagcttca ggtttgggga agagacaaca actccctctc agaagcagga gccgatagac	1440
aaggaaactgt atccttttagc ttccctcaga tcactctttg gcagcgaccc ctctgcacaa	1500
taa	1503

<210> SEQ ID NO 53

<211> LENGTH: 1872

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Helper - HIV Pol - Protease and reverse transcriptase

<400> SEQUENCE: 53

atgaatttgc caggaagatg gaaacaaaa atgatagggg gaattggagg ttttatcaaa	60
gtaggacagt atgatcagat actcatagaa atctgcggac ataaagctat aggtacagta	120
ttagtaggac ctacacctgt caacataatt ggaagaaatc tgttgactca gattggctgc	180
actttaaatt ttcccattag tcctattgag actgtaccag taaaattaaa gccaggaatg	240
gatggcccaa aagttaaaca atggccattg acagaagaaa aaataaaagc attagtagaa	300
atttgtacag aaatggaaaa ggaaggaaaa atttcaaaaa ttgggcctga aaatccatac	360
aatactccag tatttgccat aaagaaaaaa gacagtacta aatggagaaa attagtagat	420
ttcagagaac ttaataagag aactcaagat ttctgggaag ttcaattagg aataccacat	480
cctgcagggt taaaacagaa aaaatcagta acagtactgg atgtgggcga tgcatatttt	540
tcagttccct tagataaaga cttcaggaag tatactgcat ttaccatacc tagtataaac	600
aatgagacac cagggtattg atatcagtag aatgtgcttc cacagggatg gaaaggatca	660
ccagcaatat tccagtgtag catgacaaaa atcttagagc cttttagaaa acaaaatcca	720
gacatagtca tctatcaata catggatgat ttgtatgtag gatctgactt agaaataggg	780
cagcatagaa caaaaataga ggaactgaga caacatctgt tgagggtggg atttaccaca	840

-continued

ccagacaaaa aacatcagaa agaacctcca ttccttttga tgggttatga actccatcct	900
gataaatgga cagtacagcc tatagtgtg ccagaaaagg acagctggac tgtcaatgac	960
atacagaaat tagtgggaaa attgaattgg gcaagtcaga tttatgcagg gattaaagta	1020
aggcaattat gtaaaattct taggggaacc aaagcactaa cagaagtagt accactaaca	1080
gaagaagcag agctagaact ggcagaaaac agggagattc taaaagaacc ggtacatgga	1140
gtgtattatg acccatcaaa agacttaata gcagaaatc agaagcagg gcaaggccaa	1200
tggacatatc aaatttatca agagccattt aaaaatctga aaacaggaaa atatgcaaga	1260
atgaaggggtg cccacactaa tgatgtgaaa caattaacag aggcagtaca aaaaatagcc	1320
acagaaagca tagtaatatg gggaaagact cctaaattta aattacccat aaaaaggaa	1380
acatgggaag catggtggac agagtattgg caagccacct ggattcctga gtgggagttt	1440
gtcaataccc ctcccttagt gaagttagt taccagttag agaagaacc cataatagga	1500
gcagaaactt tctatgtaga tggggcagcc aatagggaaa ctaaattagg aaaagcagga	1560
tatgtaactg acagaggaag acaaaaagtt gtccccctaa cggacacaac aaatcagaag	1620
actgagttac aagcaattca tctagctttg caggattcgg gattagaagt aaacatagtg	1680
acagactcac aatatgcatt gggaaatcatt caagcacac cagataagag tgaatcagag	1740
ttagtcagtc aaataataga gcagttaata aaaaaggaaa aagtctacct ggcatgggta	1800
ccagcacaca aaggaattgg aggaaatgaa caagtagatg gggtggtcag tgctggaatc	1860
aggaaagtac ta	1872

<210> SEQ ID NO 54

<211> LENGTH: 867

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Helper - HIV Integrase - Integration of viral RNA

<400> SEQUENCE: 54

tttttagatg gaatagataa ggccaagaa gaacatgaga aatatcacag taattggaga	60
gcaatggcta gtgattttta cctaccacct gtagtagcaa aagaaatagt agccagctgt	120
gataaatgtc agctaaaagg ggaagccatg catggacaag tagactgtag cccaggaata	180
tggcagctag attgtacaca tttagaagga aaagtattct tggtagcagt tcatgtagcc	240
agtggatata tagaagcaga agtaattcca gcagagacag ggcaagaaac agcatacttc	300
ctcttaaaat tagcaggaag atggccagta aaaacagtac atacagacaa tggcagcaat	360
ttcaccagta ctacagttaa ggcgcctgt tggtagggcg ggatcaagca ggaatttggc	420
attccctaca atccccaag tcaaggagta atagaatcta tgaataaaga attaaagaaa	480
attataggac aggtaagaga tcaggctgaa catcttaaga cagcagtaca aatggcagta	540
ttcatccaca attttaaaag aaaagggggg attggggggg acagtgcagg ggaaagaata	600
gtagacataa tagcaacaga catacaaact aaagaattac aaaaacaaat tacaaaaatt	660
caaaattttc gggtttatta cagggacagc agagatccag tttggaaagg accagcaaag	720
ctcctctgga aaggtgaagg ggcagtagta atacaagata atagtgcacat aaaagtagtg	780
ccaagaagaa aagcaaagat catcagggat tatggaaaac agatggcagg tgatgattgt	840
gtggcaagta gacaggatga ggattaa	867

-continued

<210> SEQ ID NO 55
 <211> LENGTH: 234
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper - HIV RRE - Binds Rev element

<400> SEQUENCE: 55

aggagcctttg ttccttgggt tcttgggagc agcaggaagc actatgggcg cagcgtaaat	60
gacgctgacg gtacaggcca gacaattatt gtctggata gtgcagcagc agaacaattt	120
gctgagggct attgaggcgc aacagcatct gttgcaactc acagtctggg gcatcaagca	180
gtccaggca agaactctgg ctgtggaaag atacctaaag gatcaacagc tcct	234

<210> SEQ ID NO 56
 <211> LENGTH: 448
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helper - Rabbit beta globin poly A - RNA stability

<400> SEQUENCE: 56

agatcttttt ccctctgcc aaaaattatgg ggacatcatg aagccccttg agcatctgac	60
ttctggctaa taaaggaaat ttattttcat tgcaatagtg tgttggaatt ttttgtgtct	120
ctcactcgga aggacatag ggagggcaaa tcatttaaaa catcagaatg agtatttggt	180
ttagagtttg gcaacatag ccatatgctg gctgccatga acaaagggtg ctataaagag	240
gtcatcagta tatgaaacag cccctgctg tccattcctt attccataga aaagccttga	300
cttgagggtta gatttttttt atattttgtt ttgtgttatt tttttcttta acatccctaa	360
aattttcctt acatgtttta ctagccagat ttttcctcct ctctgacta ctcccagtca	420
tagctgtccc tcttctctta tgaagatc	448

<210> SEQ ID NO 57
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Rev - RSV promoter - Transcription

<400> SEQUENCE: 57

atggcaggaa gaagcggaga cagcgacgaa gaactcctca aggcagtcag actcatcaag	60
tttctctatc aaagcaaccc acctcccaat cccgagggga cccgacaggc ccgaaggaa	120
agaagaagaa ggtggagaga gagacagaga cagatccatt cgattagtga acggatcctt	180
agcacttata tgggacgata tgccggagcct gtgcctcttc agctaccacc gcttgagaga	240
cttactcttg attgtaacga ggattgtgga acttctggga cgcagggggg gggaaagccct	300
caaatatttg tggaatctcc tacaatattg gagtcaggag ctaaagaata g	351

<210> SEQ ID NO 58
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Rev - HIV Rev- Nuclear export and stabilize viral mRNA

-continued

<400> SEQUENCE: 58

```
atggcaggaa gaagcggaga cagcgacgaa gaactcctca aggcagtcag actcatcaag    60
tttctctatc aaagcaaccc acctcccaat cccgagggga cccgacaggc ccgaaggaat    120
agaagaagaa ggtggagaga gagacagaga cagatccatt cgattagtga acggatcctt    180
agcacttatc tgggacgata tgcggagcct gtgcctcttc agctaccacc gcttgagaga    240
cttactcttg attgtaacga ggattgtgga acttctggga cgcagggggg gggaagccct    300
caaatattgg tggaatctcc tacaatattg gagtcaggag ctaaagaata g            351
```

<210> SEQ ID NO 59

<211> LENGTH: 450

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Rev- Rabbit beta globin poly A- RNA stability

<400> SEQUENCE: 59

```
agatcttttt cctctgccca aaaattatgg ggacatcatg aagcccttg agcatctgac    60
ttctggctaa taaaggaaat ttattttcat tgcaatagtg tgttgaatt ttttgtgtct    120
ctcactcgga aggacatatg ggagggcaaa tcatttaaaa catcagaatg agtatttggt    180
ttagagtttg gcaacatatg cccatatgct ggctgccatg aacaaagggt ggctataaag    240
aggtcatcag tatatgaaac agccccctgc tgtccattcc ttattccata gaaaagcctt    300
gacttgaggt tagatttttt ttatatattg ttttgtgta ttttttctt taacatccct    360
aaaattttcc ttacatgttt tactagccag atttttcttc ctctcctgac tactcccagt    420
catagctgtc cctcttctct tatggagatc            450
```

<210> SEQ ID NO 60

<211> LENGTH: 577

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- CMV promoter- Transcription

<400> SEQUENCE: 60

```
acattgatta ttgactagtt attaatagta atcaattacg gggtcattag ttcatagccc    60
atatatggag ttccgcgtta cataacttac ggtaaatggc ccgcctggct gaccgcccaa    120
cgacccccgc ccattgaagt caataatgac gtatgttccc atagtaacgc caatagggac    180
tttccattga cgtcaatggg tggagtattt acggtaaaact gccacttg cagtacatca    240
agtgtatcat atgccaaagta cgcacctat tgacgtcaat gacggtaaat ggccgcctg    300
gcattatgcc cagtacatga ctttatggga ctttctact tggcagtaca tctacgtatt    360
agtcatcgct attaccatgg tgatgcggtt ttggcagtac atcaatgggc gtggatagcg    420
gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga gtttgtttg    480
gcacccaaat caacgggact ttccaaaatg tcgtaacaac tccgccccat tgacgcaaat    540
gggcggtagg cgtgtacggt gggaggtcta tataagc            577
```

<210> SEQ ID NO 61

<211> LENGTH: 573

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- Beta globin intron- Enhance gene expression

<400> SEQUENCE: 61

```

gtgagtttgg ggacccttga ttgttctttc ttttctgcta ttgtaaaatt catgttatat    60
ggaggggggca aagttttcag ggtgttggtt agaatgggaa gatgtccctt gtatcaccat    120
ggaccctcat gataattttg tttctttcac tttctactct gttgacaacc attgtctcct    180
cttattttct tttcattttc tgtaactttt tcgttaaact ttagcttgca ttgtaacga    240
atttttaaat tcacttttgt ttatttgtca gattgtaagt actttctcta atcacttttt    300
tttcaaggca atcaggggat attatattgt acttcagcac agtttttagag aacaattggt    360
ataattaaat gataaggtag aatatttctg catataaatt ctggctggcg tggaaatatt    420
cttattggta gaaacaacta caccctggtc atcatcctgc ctttctcttt atggttacaa    480
tgatatacac tgtttgagat gaggataaaa tactctgagt ccaaaccggg cccctctgct    540
aaccatgttc atgccttctt ctctttccta cag                                573

```

<210> SEQ ID NO 62

<211> LENGTH: 1519

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- VSV-G- Glycoprotein envelope-cell entry

<400> SEQUENCE: 62

```

atgaagtgcc ttttgtactt agccttttta ttcattgggg tgaattgcaa gttcaccata    60
gtttttccac acaacaaaaa aggaaactgg aaaaatgttc cttctaatta ccattattgc    120
ccgtcaagct cagatttaaa ttggcataat gacttaatag gcacagcctt acaagtcaaa    180
atgcccaga gtcacaaggc tattcaagca gacggttgga tgtgtcatgc ttccaaatgg    240
gtcactactt gtgatttccg ctggtatgga ccgaagtata taacacattc catccgatcc    300
ttcactccat ctgtagaaca atgcaaggaa agcattgaac aaacgaaaca aggaacttgg    360
ctgaatccag gcttcctctc tcaaagtgtt ggatatgcaa ctgtgacgga tgccgaagca    420
gtgattgtcc aggtgactcc tcaccatgtg ctggttgatg aatacacagg agaatgggtt    480
gattcacagt tcactcaacg aaaatgcagc aattacatat gcccactgt ccataactct    540
acaacctggc attctgacta taaggtaaaa gggctatgtg attctaacct catttccatg    600
gacatcacct tcttctcaga ggacggagag ctatcatccc tgggaaagga gggcacaggg    660
ttcagaagta actactttgc ttatgaaact ggaggcaagg cctgcaaaaat gcaatactgc    720
aagcattggg gagtcagact cccatcaggt gtctgggtcg agatggctga taaggatctc    780
tttgctgcag ccagattccc tgaatgccc gaagggtcaa gtatctctgc tccatctcag    840
acctcagtgg atgtaagtct aattcaggac gttgagagga tcttgatta ttccctctgc    900
caagaaacct ggagcaaaaat cagagcgggt cttccaatct ctccagtga tctcagctat    960
cttgctccta aaaaccaggg aaccggctct gctttcacca taatcaatgg taccctaaaa    1020
tactttgaga ccagatacat cagagtcgat attgtgctc caatcctctc aagaatggtc    1080
ggaatgatca gtggaactac cacagaaagg gaactgtggg atgactgggc accatatgaa    1140
gacgtggaaa ttggacccaa tggagttctg aggaccagtt caggatataa gtttccttta    1200

```

-continued

tacatgattg gacatggtat gttggactcc gatcttcac	ttagctcaaa ggctcagggtg	1260
ttcgaacatc ctcacattca agacgctgct tcgcaacttc	ctgatgatga gagtttattt	1320
tttggtgata ctgggctatc caaaaatcca atcgagcttg	tagaagggtg gttcagtagt	1380
tggaaaagct ctattgcctc ttttttcttt atcatagggt	taatcattgg actattcttg	1440
gttctccgag ttggtatcca tctttgcatt aaattaaagc	acaccaagaa aagacagatt	1500
tatacagaca tagagatga		1519

<210> SEQ ID NO 63
 <211> LENGTH: 450
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Envelope- Rabbit beta globin poly A- RNA stability

<400> SEQUENCE: 63

agatcttttt ccctctgcca aaaattatgg ggacatcatg	aagccccttg agcatctgac	60
ttctggctaa taaaggaaat ttattttcat tgcaatagtg	tgttggaatt ttttgtgtct	120
ctcactcgga aggacatagtg ggagggcaaa tcatttaaaa	catcagaatg agtatttggt	180
ttagagtgtg gcaacatagtg cccatagctt ggctgccatg	aacaaagggt ggctataaag	240
aggatcatcag tatatgaaac agccccctgc tgtccattcc	ttattccata gaaaagcctt	300
gacttgagggt tagatttttt ttatattttg ttttgtgtta	ttttttctt taacatccct	360
aaaattttcc ttacatgttt tactagccag atttttcctc	ctctcctgac tactcccagt	420
catagctgtc cctcttctct tatggagatc		450

<210> SEQ ID NO 64
 <211> LENGTH: 1104
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Promoter- EF-1

<400> SEQUENCE: 64

cgggtgcta gagaagggtg cgcggggttaa actgggaaag	tgatgtcgtg tactggctcc	60
gcctttttcc cgagggtggg ggagaaccgt atataagtgc	agtagtcgcc gtgaacgttc	120
tttttcgcaa cggttttgcc gccagaacac aggttaagtgc	cgtgtgtggt tcccgcgggc	180
ctggcctctt tacgggttat ggcccttgcg tgccttgaat	tacttccacg cccctggctg	240
cagtacgtga ttcttgatcc cgagcttcgg gttggaagtg	gggtgggagag ttcgaggcct	300
tgcgcttaag gagccccttc gcctcgtgct tgagttgagg	cctggcctgg gcgctggggc	360
cgccgcgtgc gaatctgggt gcaccctcgc gcctgtctcg	ctgctttcga taagtctcta	420
gccatttaaa atttttgatg acctgtgcg acgttttttt	tctggcaaga tagtcttgta	480
aatgcggggc aagatctgca cactggtatt tcggtttttg	gggcccgggg cggcgacggg	540
gccctgtcgt cccagcgcac atgttcggcg aggcggggcc	tgcgagcgcg gccaccgaga	600
atcgacgggg ggtagtctca agctggccgg cctgctctgg	tgctggcct cgcgccggcg	660
tgtatcgccc cgccctgggc ggcaaggctg gcccggtcgg	caccagttgc gtgagcgga	720
agatggccgc ttcccggccc tgctgcaggg agctcaaaat	ggaggacgcg gcgctcgga	780

-continued

gagcgggagg	gtgagtcacc	cacacaaagg	aaaagggcct	ttccgtcctc	agccgtcgct	840
tcatgtgact	ccacggagta	ccggggcgccg	tccaggcacc	tcgattagtt	ctcgagcttt	900
tggagtacgt	cgtcttttagg	ttgggggggag	gggttttatg	cgatggagtt	tccccacact	960
gagtgggtgg	agactgaagt	taggccagct	tggcacttga	tgtaattctc	cttgggaattt	1020
gccctttttg	agtttgatc	ttggttcatt	ctcaagcctc	agacagtggg	tcaaagtttt	1080
tttcttccat	ttcaggtgtc	gtga				1104

<210> SEQ ID NO 65
 <211> LENGTH: 511
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Promoter- PGK

<400> SEQUENCE: 65

gggggtgggg	ttgcgccttt	tccaaggcag	ccctgggttt	gcgcagggac	gcggctgctc	60
tgggcgtggt	tccgggaaac	gcagcggcgc	cgaccctggg	tctcgacat	tcttcacgtc	120
cgttcgcagc	gtcaccggga	tcttcgccgc	tacccttggt	ggccccccgg	cgacgcttcc	180
tgtccgccc	ctaagtcggg	aaggttcctt	gcggttcgcg	gcgtgccgga	cgtgacaaac	240
ggaagccgca	cgtctcacta	gtaccctcgc	agacggacag	cgccaggagg	caatggcagc	300
gcgcgcagcg	cgatgggctg	tggccaatag	cggtgctca	gcagggcgcg	ccgagagcag	360
cggccgggaa	ggggcggtgc	gggaggcggg	gtgtggggcg	gtagtgtggg	ccctgttcct	420
gcccgcgcgg	tgttccgcat	tctgcaagcc	tccggagcgc	acgtcggcag	tcggctccct	480
cgttgaccga	atcaccgacc	tctctcccca	g			511

<210> SEQ ID NO 66
 <211> LENGTH: 1162
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Promoter- UbC

<400> SEQUENCE: 66

gcgcggggtt	ttggcgctc	ccgcggggcg	ccccctcctc	acggcgagcg	ctgccacgtc	60
agacgaaggg	cgcaggagcg	ttcctgatcc	ttccgcccg	acgtcagga	cagcggccc	120
ctgctcataa	gactcggcct	tagaacccca	gtatcagcag	aaggacattt	taggacggga	180
cttgggtgac	tctagggcac	tggttttctt	tccagagagc	ggaacaggcg	aggaaaagta	240
gtcccttctc	ggcgattctg	cggagggtac	tccgtggggc	ggtgaacgcc	gatgattata	300
taaggacgcg	ccgggtgtgg	cacagctagt	tccgtcgag	ccgggatttg	ggtcgcggtt	360
cttggtttgt	gatecgtgtg	atcgtcactt	ggtgagttgc	gggtcgttgg	gctggccggg	420
gctttcgtgg	ccgcggggcc	gctcgtggg	acggaagcgt	gtggagagac	cgccaagggc	480
tgtagtctgg	gtccgcgagc	aaggttgccc	tgaactgggg	gttgggggga	gcgcacaaaa	540
tggcggtgtg	tcccagtgct	tgaatggaag	acgcttgtaa	ggcgggctgt	gaggtcgttg	600
aaacaagggtg	gggggcatgg	tgggcggcaa	gaacccaagg	tcttgaggcc	ttcgctaata	660
cgggaaagct	cttattcggg	tgagatgggc	tggggcacca	tctggggacc	ctgacgtgaa	720
gtttgtcact	gactggagaa	ctcgggtttg	tcgtctggtt	gcggggcgcg	cagttatgcg	780

-continued

gtgccgttgg gcagtgcacc cgtacctttg ggagcgcgcg cctcgtcgtg tcgtgacgtc	840
acccgtttctg ttggttata atgcaggggtg gggccacctg ccggtaggtg tgcggtaggc	900
ttttctccgt cgcaggacgc aggggttcggg cctagggtag gctctcctga atcgacaggc	960
gccggacctc tgggtagggg agggataagt gaggcgtcag tttctttggt cggttttatg	1020
tacctatctt cttaagtgc tgaagctccg gttttgaact atgcgcctcg ggttggcgag	1080
tgtgttttgt gaagtttttt aggcaccttt tgaaatgtaa tcatttgggt caatatgtaa	1140
ttttcagtgt tagactagta aa	1162

<210> SEQ ID NO 67
 <211> LENGTH: 120
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Poly A- SV40

<400> SEQUENCE: 67

gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt tcacaaataa	60
agcatttttt tcaactgcatt ctagtgtggg ttgtccaaa ctcacaaatg tatcttatca	120

<210> SEQ ID NO 68
 <211> LENGTH: 227
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Poly A- bGH

<400> SEQUENCE: 68

gactgtgcct tctagttgcc agccatctgt tgtttgcctc tccccgtgc cttocttgac	60
cctggaaggt gccactccca ctgtccttcc ctaataaaat gaggaattg catcgcatg	120
tctgagtagg tgtcattcta ttctgggggg tgggggtggg caggacagca agggggagga	180
ttgggaagac aatagcaggc atgctgggga tgcgggtggg tctatgg	227

<210> SEQ ID NO 69
 <211> LENGTH: 1512
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: HIV Gag- Bal

<400> SEQUENCE: 69

atgggtgcga gacgtcagc attaagcggg ggagaattag atagggtggg aaaaattcgg	60
ttaaggccag ggggaaagaa aaaatataga ttaaaacata tagtatgggc aagcagggaa	120
ctagaaagat tcgcagtcac tcttgccctg ttagaaacat cagaaggctg cagacaaata	180
ctgggacagc tacaaccatc ccttcagaca ggatcagaag aacttagatc attatataat	240
acagtagcaa ccctctattg tgtacatcaa aagatagagg taaaagacac caaggagct	300
ttagacaaaa tagaggaaga gcaaaacaaa tgtaagaaaa aggcacagca agcagcagct	360
gacacaggaa acagcgttca ggtcagccaa aatttcctta tagtgagaa cctccagggg	420
caaatggtac atcaggccat atcacctaga actttaaatg catgggtaaa agtaatagaa	480
gagaaagctt tcagcccaga agtaataccc atgttttcag cattatcaga aggagccacc	540
ccacaagatt taaacaccat gctaaacaca gtggggggac atcaagcagc catgcaaatg	600

-continued

ttaaaagaac ccatcaatga ggaagctgca agatgggata gattgcatcc cgtgcaggca	660
gggcctgttg caccaggcca gataagagat ccaaggggaa gtgacatagc aggaactacc	720
agtacccttc aggaacaaat aggatggatg acaagtaatc cacctatccc agtaggagaa	780
atctataaaa gatggataat cctgggatta aataaaatag taaggatgta tagccctacc	840
agcatttttg acataagaca aggaccaaag gaacccttta gagactatgt agaccggttc	900
tataaaactc taagagccga gcaagcttca caggaggtaa aaaattggat gacagaaacc	960
ttgttggtcc aaaatgcgaa ccagattgt aagactattt taaaagcatt gggaccagca	1020
gtacactag aagaaatgat gacagcatgt cagggagtgg gaggaccag ccataaagca	1080
agaatttttg cagaagcaat gagccaagta acaaattcag ctaccataat gatgcagaaa	1140
ggcaatttta ggaaccaaag aaagattgtt aaatgtttca attgtggcaa agaagggcac	1200
atagccagaa actgcagggc ccttaggaaa aggggctgtt ggaaatgtgg aaaggaagga	1260
caccaaatga aagactgtac tgagagacag gctaattttt tagggaaaat ctggccttcc	1320
cacaaaggaa ggccagggaa ttccttcag agcagaccag agccaacagc cccaccagcc	1380
ccaccagaag agagcttcag gtttggggaa gagacaacaa ctccctctca gaagcaggag	1440
ctgatagaca aggaactgta tcctttagct tccctcagat cactctttgg caacgacccc	1500
tcgtcacaat aa	1512

<210> SEQ ID NO 70

<211> LENGTH: 1872

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: HIV Pol- Bal

<400> SEQUENCE: 70

atgaatttgc caggaagatg gaaacaaaa atgatagggg gaattggagg ttttatcaaa	60
gtaagacagt atgatcagat actcatagaa atctgtggac ataaagctat aggtacagta	120
ttaataggac ctacacctgt caacataatt ggaagaaatc tgttgactca gattggttgc	180
actttaaatt tcccattag tcctattgaa actgtaccag taaaattaaa accaggaatg	240
gatggcccaa aagttaaaca atggccactg acagaagaaa aaataaaagc attaattggaa	300
atctgtacag aaatggaaaa ggaagggaaa atttcaaaaa ttgggcctga aaatccatac	360
aatactccag tatttgccat aaagaaaaaa gacagtacta aatggagaaa attagtagat	420
ttcagagAAC ttaataagaa aactcaagac ttctgggaag tacaattagg aatacacatc	480
ccgcaggggt taaaaaagaa aaaatcagta acagtactgg atgtgggtga tgcataatTT	540
tcagttccct tagataaaga attcaggaag tatactgcat ttaccatacc tagtataaac	600
aatgaaacac cagggatcag atatcagtac aatgtacttc cacagggatg gaaaggatca	660
ccagcaatat ttcaagtag catgacaaga atcttagagc cttttagaaa acaaaatcca	720
gaaatagtga tctatcaata catggatgat ttgtatgtag gatctgactt agaaataggg	780
cagcatagaa caaaaataga ggaactgaga caacatctgt tgagggtggg atttaccaca	840
ccagacaaaa aacatcagaa agaacctcca ttcctttgga tgggttatga actccatcct	900
gataaatgga cagtacagcc tatagtgtg cagaaaaag acagctggac tgtcaatgac	960
atacagaagt tagtgggaaa attgaattgg gcaagtcaga tttaccagg aattaaagta	1020

-continued

aagcaattat gtaggctcct taggggaacc aaggcattaa cagaagtaat accactaaca	1080
aaagaaacag agctagaact ggagagagac agggaaattc taaaagaacc agtacatggg	1140
gtgtattatg acccatcaaa agacttaata gcagaaatac agaagcaggg gcaaggccaa	1200
tggacatatc aaatttatca agagccattt aaaaatctga aaacaggaaa atatgcaaga	1260
atgaggggtg cccacactaa tgatgtaaaa caattaacag aggcagtgcg aaaaataacc	1320
acagaaagca tagtaatatg gggaaagact cctaaattta aactacccat aaaaaagaa	1380
acatgggaaa catggtggac agagtattgg caagccacct ggattcctga gtgggagttt	1440
gtcaataccc ctcccttagt gaaattatgg taccagttag agaagaacc cataatagga	1500
gcagaaacat tctatgtaga tggagcagct aaccgggaga ctaaattagg aaaagcagga	1560
tatgttacta acagaggaag acaaaaagtt gtctccctaa ctgacacaac aaatcagaag	1620
actgagttac aagcaattca tctagcttta caagattcag gattagaagt aaacatagta	1680
acagactcac aatatgcatt aggaatcatt caagcacac cagataaaaag tgaatcagag	1740
ttagtcagtc aaataataga acagttaata aaaaaggaaa aggtctacct ggcatgggta	1800
ccagcgaca aaggaattgg aggaaatgaa caagtagata aattagtcag tactggaatc	1860
aggaagtac ta	1872

<210> SEQ ID NO 71
 <211> LENGTH: 867
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: HIV Integrase- Bal
 <400> SEQUENCE: 71

tttttagatg gaatagatat agcccaagaa gaacatgaga aatatcacag taattggaga	60
gcaatggcta gtgattttta cctgccacct gtggtagcaa aagaaatagt agccagctgt	120
gataaatgtc agctaaaagg agaagccatg catggacaag tagactgtag tccaggaata	180
tggcaactag attgtacaca tttagaagga aaaattatcc tggtagcagt tcatgtagcc	240
agtggatata tagaagcaga agttattcca gcagagacag ggcaggaaac agcatacttt	300
ctcttaaaat tagcaggaag atggccagta aaaacaatac atacagacaa tggcagcaat	360
ttcactagta ctacagtcga ggcgcctgt tggtagggcg ggatcaagca ggaatttggc	420
attccctaca atcccaaag tcaggagta gtagaatcta taaataaaga attaaagaaa	480
attataggac aggtgaagaga tcaggctgaa catcttaaaa cagcagtaca aatggcagta	540
ttcatccaca attttaaaag aaaagggggg attggggggg atagtgcagg ggaagaata	600
gtagacataa tagcaacaga catacaaact aaagaattac aaaaacaaat tacaaaaatt	660
caaaattttc gggtttatta cagggacagc agagatccac tttggaagg accagcaaag	720
cttctctgga aaggtgaagg ggcagtagta atacaagata atagtgcacat aaaagtagta	780
ccaagaagaa aagcaaagat cattagggat tatggaaaac agatggcagg tgatgattgt	840
gtggcaagta gacaggatga ggattag	867

<210> SEQ ID NO 72
 <211> LENGTH: 1695
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Envelope- RD114

<400> SEQUENCE: 72

```

atgaaactcc caacaggaat ggtcatttta tgtagcctaa taatagtctg ggcagggttt      60
gacgaccccc gcaaggctat cgcattagta caaaaacaac atggtaaacc atgcgaatgc      120
agcggaggggc aggtatccga ggccccaccg aactccatcc aacaggtaac ttgccaggc      180
aagacggcct acttaatgac caacaaaaaa tggaaatgca gagtcaactcc aaaaaatctc      240
acccttagcg ggggagaact ccagaactgc cctgtgaaca cttccagga ctcgatgcac      300
agtctctgtt atactgaata ccggcaatgc agggcggaata ataagacata ctacacggcc      360
accttgctta aaatacggtc tgggagcctc aacgaggtag agatattaca aaaccccaat      420
cagctcctac agtccccttg taggggctct ataaatcagc ccgtttgctg gagtgcacaca      480
gccccatcc atactcctga tggtaggagga cccctcgata ctaagagagt gtggacagtc      540
caaaaaaggc tagaacaat tcataaggct atgcacctc aacttaata ccaccctta      600
gccctgccc aagtcagaga tgaccttagc ctgatgcac ggacttttga tatcctgaat      660
accactttta ggttactcca gatgtccaat tttagccttg cccaagattg ttggctctgt      720
ttaaaactag gtaccctac cctcttgctg ataccactc cctctttaac ctactcccta      780
gcagactccc tagcgaatgc ctctgtcag attatactc cctcttggg tcaaccgatg      840
cagttctcca actcgtcctg tttatcttc ccttccatta acgatacga acaaatagac      900
ttagtgtagc tcacctttac taactgcacc tctgtagcca atgtcagtag tcctttatgt      960
gccctaaacg ggtcagtctt cctctgtgga aataacatgg catacaccta ttaccccaa      1020
aactggacag gactttgcgt ccaagcctcc ctctccccc acattgacat catcccgggg      1080
gatgagccag tccccattcc tgccattgat cattatatac atagacctaa acgagctgta      1140
cagttcatcc ctttactagc tggactggga atcacgcag cattcaccac cgagactaca      1200
ggcctagggt tctccgtcac ccagtataca aaattatccc atcagttaat atctgatgtc      1260
caagtcttat ccggtaccat acaagattta caagaccagg tagactcgtt agctgaagta      1320
gttctccaaa ataggagggg actggacctc ctaacggcag aacaaggagg aatttgttta      1380
gccttacaag aaaaatgctg tttttatgct aacaagtcag gaattgtgag aaacaaaata      1440
agaaccttac aagaagaatt acaaaaacgc agggaaagcc tggcatccaa ccctctctgg      1500
accgggctgc agggctttct tccgtacctc ctacctctc tgggacccct actcaccctc      1560
ctactcatac taaccattgg gccatgcgtt ttcaatcgat tggccaatt tgttaaagac      1620
aggatctcag tggtcaggc tctggttttg actcagcaat atcaccagct aaaaccata      1680
gagtacgagc catga                                          1695

```

<210> SEQ ID NO 73

<211> LENGTH: 2013

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- GALV

<400> SEQUENCE: 73

```

atgcttctca cctcaagccc gcaccacctt cggcaccaga tgagtcttg gagctggaaa      60
agactgatca tcctcttaag ctgcgtattc ggagacggca aaacgagtct gcagaataag      120

```


-continued

aacccccacc agcctgtgac cctcacctgg caggtactgt cccaaactgg ggacgttgtc	180
tgggacaaaa aggcagtcaca gcccctttgg acttggtggc cctctcttac acctgatgta	240
tgtgcccctgg cggcgggtct tgagtcctgg gatatcccg gatccgatgt atcgctctct	300
aaaagagtta gacctctga ttcagactat actgccgctt ataagcaaat cacctgggga	360
gcataggggt gcagctaccc tcgggctagg accaggatgg caaattcccc cttctacgtg	420
tgtccccgag ctggccgaac ccattcagaa gctaggagggt gtgggggggt agaatacccta	480
tactgtaaag aatggagttg tgagaccacg ggtaccgttt attggcaacc caagtcctca	540
tgggacctca taactgtaaa atgggaccaa aatgtgaaat gggagcaaaa atttcaaaag	600
tgtgaacaaa cggcgtggtg taacccctc aagatagact tcacagaaaa aggaaaactc	660
tccagagatt ggataacgga aaaaacctgg gaattaaggt tctatgtata tggacacca	720
ggcatacagt tgactatccg cttagaggtc actaacatgc cggttgtggc agtgggcccc	780
gacctgttcc ttgcggaaca gggacctct agcaagcccc tcaactctcc tctctcccc	840
cggaaagcgc cgccacccc tctaccccg gcggctagt agcaaacc tgcggtgcat	900
ggagaaactg ttaccctaaa ctctccgct cccaccagt gcgaccgact ctttggcctt	960
gtgcaggggg ccttcctaac cttgaatgct accaaccag gggccactaa gtcttgctgg	1020
ctctgtttgg gcatgagccc cccttattat gaagggatag cctcttcagg agaggctgct	1080
tatacctcca accataccg atgccactgg ggggccccag gaaagcttac cctcactgag	1140
gtctccggac tcgggtcatg catagggaag gtgcctctta cccatcaaca tctttgcaac	1200
cagaccttac ccatcaattc ctctaaaaac catcagtatc tgctccctc aaaccatagc	1260
tgggtgggct gcagcactgg cctcaccccc tgctctccca cctcagtttt taatcagtct	1320
aaagacttct gtgtccaggt ccagctgac ccccgcatct attaccattc tgaagaaacc	1380
ttgttacaag cctatgacaa atcaccccc aggtttaaaa gagagcctgc ctcacttacc	1440
ctagctgtct tcctgggggt agggattgct gcaggtatag gtactggctc aacggcccta	1500
attaaagggc ccatagacct ccagcaaggc ctaaccagcc tccaaatcgc cattgacgct	1560
gacctccggg cccttcagga ctcaatcagc aagctagagg actcactgac ttccctatct	1620
gaggtagtac tccaaaatag gagaggcctt gacttactat tccttaaaga aggaggcctc	1680
tgcgggggcc taaaagaaga gtgctgtttt tatgtagacc actcaggtgc agtacgagac	1740
tccatgaaaa aacttaaaga aagactagat aaaagacagt tagagcgcca gaaaaaccaa	1800
aactgggatg aagggtggtt caataactcc ccttgggtta ctaccctact atcaaccatc	1860
gctggggccc tattgtctct ccttttggtta ctactcttg ggcctgcat catcaataaa	1920
ttaatccaat tcataatga taggataagt gcagtcaaaa ttttagtcct tagacagaaa	1980
tatcagaccc tagataacga ggaaaacctt taa	2013

<210> SEQ ID NO 74

<211> LENGTH: 1530

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- FUG

<400> SEQUENCE: 74

atggttccgc aggttctttt gttgtactc cttctgggtt tttcgttggtg tttcgggaag	60
---	----

-continued

ttccccattt acacgatacc agacgaactt ggtccctgga gccctattga catabaccat	120
ctcagctgtc caaataacct ggttgtggag gatgaaggat gtaccaacct gtccgagttc	180
tcctacatgg aactcaaagt gggatacatc tcagccatca aagtgaacgg gttcacttgc	240
acaggtgttg tgacagaggg agagacctac accaactttg ttggttatgt cacaaccaca	300
ttcaagagaa agcattttccg cccaccccca gacgcatgta gagccgcgta taactggaag	360
atggccgggtg accccagata tgaagagtcc ctacacaatc cataccccga ctaccactgg	420
cttcgaactg taagaaccac caaagagtcc ctcatatca tatccccaag tgtgacagat	480
ttggacccat atgacaaatc ccttcactca agggctcttc ctggcggaaa gtgctcagga	540
ataacggtgt cctctaccta ctgctcaact aaccatgatt acaccatttg gatgcccgag	600
aatccgagac caaggacacc ttgtgacatt ttaccaata gcagagggaa gagagcatcc	660
aacgggaaca agacttgcgg ctttgtggat gaaagaggcc tgtataagtc tctaaaagga	720
gcctgcagggc tcaagttatg tggagttctt ggacttagac ttatggatgg aacatgggtc	780
gcgatgcaaa catcagatga gaccaaatgg tgcctccag atcagttggt gaatttgac	840
gactttcgct cagacgagat cgagcatctc gttgtggagg agttagttaa gaaaagagag	900
gaatgtctgg atgcattaga gtccatcatg accaccaagt cagtaagttt cagacgtctc	960
agtcacctga gaaaacttgt cccagggttt ggaaaagcat ataccatatt caacaaaacc	1020
ttgatggagg ctgatgtcca ctacaagtca gtccggacct ggaatgagat catccctca	1080
aaagggtgtt tgaaagttgg aggaagggtc catcctcatg tgaacggggt gtttttcaat	1140
ggtataatat tagggcctga cgaccatgtc ctaatccag agatgcaatc atccctctc	1200
cagcaacata tggagttgtt ggaatcttca gttatcccc tgatgcacc cctggcagac	1260
ccttctacag ttttcaaaga aggtgatgag gctgaggatt ttgttgaagt tcacctcccc	1320
gatgtgtaca aacagatctc aggggttgac ctgggtctcc cgaactgggg aaagtatgta	1380
ttgatgactg caggggcat gattggcctg gtgttgatat ttccctaata gacatgggtc	1440
agagttggtg tccatctttg cattaaatta aagcacacca agaaaagaca gatttatata	1500
gacatagaga tgaaccgact tggaaagtaa	1530

<210> SEQ ID NO 75

<211> LENGTH: 1497

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- LCMV

<400> SEQUENCE: 75

atgggtcaga ttgtgacaat gtttgaggct ctgcctcaca tcatcgatga ggtgatcaac	60
attgtcatta ttgtgcttat cgtgatcacg ggtatcaagg ctgtctacaa ttttgccacc	120
tgtgggatat tcgcattgat cagtttctca cttctggctg gcaggctctg tggcatgtac	180
ggtcttaagg gacccgacat ttacaaagga gtttaaccaat ttaagtcagt ggagtttgat	240
atgtcacatc tgaacctgac catgccccac gcatgttcag ccaacaactc ccaccattac	300
atcagtatgg ggacttcttg actagaattg accttcacca atgattccat catcagtcac	360
aacttttgca atctgacctc tgccttcaac aaaaagacct ttgaccacac actcatgagt	420
atagtttcga gcctacacct cagtatcaga gggaactcca actataaggc agtatcctgc	480

-continued

gacttcaaca atggcataac catccaatac aacttgacat tctcagatcg acaaagtgt	540
cagagccagt gtagaacctt cagaggtaga gtcctagata tgtttagaac tgccttcggg	600
gggaaataca tgaggagtgg ctggggctgg acaggctcag atggcaagac cacctggtgt	660
agccagacga gttaccaata cctgattata caaaatagaa cctgggaaaa ccactgcaca	720
tatgcaggtc cttttgggat gtccaggatt ctctttccc aagagaagac taagttcttc	780
actaggagac tagcgggcac attcacctgg actttgtcag actcttcagg ggtggagaat	840
ccagggtggt attgcctgac caaatggatg attcttctg cagagcttaa gtgtttcggg	900
aacacagcag ttgcgaaatg caatgtaaat catgatgccg aattctgtga catgctgcga	960
ctaattgact acaacaaggc tgctttgagt aagttcaaag aggacgtaga atctgccttg	1020
cacttattca aaacaacagt gaattctttg atttcagatc aactactgat gaggaaccac	1080
ttgagagatc tgatgggggt gccatattgc aattactcaa agttttggtg cctagaacat	1140
gcaaagaccg gcgaaactag tgccccaaag tgctggcttg tcaccaatgg ttcttactta	1200
aatgagaccc acttcagtga tcaaatcgaa caggaagccg ataacatgat tacagagatg	1260
ttgaggaagg attacataaa gaggcagggg agtacccttc tagcattgat ggaccttctg	1320
atgttttcca catctgcata tctagtcagc atcttctgc acctgtcaa aataccaaca	1380
cacaggcaca taaaagggtg ctcatgtcca aagccacacc gattaaccaa caaaggaatt	1440
tgtagtgtg gtgcatttaa ggtgcctggt gtaaaaaccg tctggaaaag acgctga	1497

<210> SEQ ID NO 76

<211> LENGTH: 1692

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- FPV

<400> SEQUENCE: 76

atgaacactc aaatcctggt tttcgccctt gtggcagtc tccccacaaa tgcagacaaa	60
atttgtcttg gacatcatgc tgtatcaaat ggcaccaaag taaacacact cactgagaga	120
ggagtagaag ttgtcaatgc aacggaaaca gtggagcggg caaacatccc caaaatttgc	180
tcaaaaggga aaagaaccac tgatcttggc caatgcggac tgtaggggac cattaccgga	240
ccacctcaat gcgaccaatt tctagaattt tcagctgac taataatcga gagacgagaa	300
ggaaatgatg tttgttaccg ggggaagttt gttaatgaag aggcattgcg acaaatcttc	360
agaggatcag gtgggattga caaagaaaca atgggattca catatagtgg aataaggacc	420
aacggaacaa ctagtgcag tagaagatca gggctttcat tctatgcaga aatggagtgg	480
ctcctgtcaa atacagacaa tgctgctttc ccacaaatga caaaatcata caaaaacaca	540
aggagagaat cagctctgat agtctgggga atccaccatt caggatcaac caccgaacag	600
accaaactat atgggagtgg aaataaactg ataacagtcg ggagttccaa atatcatcaa	660
tcttttgtgc cgagtccagg aacacgaccg cagataaatg gccagtccgg acggattgat	720
tttcattggt tgatcttggg tcccaatgat acagttactt ttagtttcaa tggggctttc	780
atagctccaa atcgtgccag cttcttgagg ggaaagtcca tggggatcca gagcgatgtg	840
caggttgatg ccaattgcga aggggaatgc taccacagtg gagggactat aacaagcaga	900
ttgccttttc aaaacatcaa tagcagagca gttggcaaat gcccaagata tgtaaaacag	960

-continued

gaaagtttat	tattggcaac	tgggatgaag	aacgttcccg	aaccttccaa	aaaaaggaaa	1020
aaaagaggcc	tgtttggcgc	tatagcaggg	tttattgaaa	atggttggga	aggtctggtc	1080
gacgggtggt	acggtttcag	gcatcagaat	gcacaaggag	aaggaactgc	agcagactac	1140
aaaagcacc	aatcggaat	tgatcagata	accgaaagt	taaatagact	cattgagaaa	1200
accaaccagc	aatttgagct	aatagataat	gaattcactg	aggtggaaaa	gcagattggc	1260
aatttaatta	actggaccaa	agactccatc	acagaagtat	ggctttacaa	tgctgaactt	1320
cttggtggca	tggaaaacca	gcacactatt	gatttggctg	attcagagat	gaacaagctg	1380
tatgagcgag	tgaggaaaaca	attaagggaa	aatgctgaag	aggatggcac	tggttgcttt	1440
gaaatttttc	ataaatgtga	cgatgattgt	atggctagta	taaggaacaa	tacttatgat	1500
cacagcaaat	acagagaaga	agcgatgcaa	aatagaatac	aaattgacct	agtcaaattg	1560
agtagtggtc	acaagatgt	gatactttgg	ttagcttcg	gggcatcatg	ctttttgctt	1620
cttgccattg	caatgggcct	tgttttcata	tgtgtgaaga	acggaaacat	gcgggtgcact	1680
atttgatat	aa					1692

<210> SEQ ID NO 77

<211> LENGTH: 1266

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- RRV

<400> SEQUENCE: 77

agtgtaacag	agcacttta	tgtgtataag	gctactagac	catacctagc	acattgcgcc	60
gattgcgggg	acgggtactt	ctgctatagc	ccagttgcta	tcgaggagat	ccgagatgag	120
gcgtctgatg	gcattgctaa	gatccaagtc	tccgccccaa	taggtctgga	caaggcaggc	180
acccacgccc	acacgaagct	ccgatatatg	gctgggtcatg	atgttcagga	atctaagaga	240
gattccttga	gggtgtacac	gtccgcagcg	tgctccatac	atgggacgat	gggacacttc	300
atcgctgcac	actgtccacc	aggcgactac	ctcaagggtt	cggttcagga	cgcagattcg	360
cacgtgaagg	catgtaaggt	ccaatacaag	cacaatccat	tgccgggtggg	tagagagaag	420
ttcggtggtt	gaccacactt	tggcgtagag	ctgcccagca	cctcatacca	gctgacaacg	480
gctcccaccg	acgaggagat	tgacatgcat	acaccgccag	atataccgga	tcgcaccctg	540
ctatcacaga	cggcggggcaa	cgtaaaaata	acagcaggcg	gcaggactat	caggtacaac	600
tgtacctgcg	gccgtgacaa	cgtaggcact	accagtactg	acaagaccat	caacacatgc	660
aagattgacc	aatgccatgc	tgcgcgcacc	agccatgaca	aatggcaatt	tacctctcca	720
tttgttccca	gggctgatca	gacagctagg	aaaggcaagg	tacacgttcc	gttccctctg	780
actaacgtca	cctgccgagt	gccgttggct	cgagcgccgg	atgccaccta	tggttaagaag	840
gaggtgaccc	tgagattaca	cccagatcat	ccgacgctct	tctcctatag	gagtttagga	900
gccgaaccgc	acccgtacga	ggaatggggt	gacaagttct	ctgagcgcat	catcccagtg	960
acggaagaag	ggattgagta	ccagtggggc	aacaacccgc	cggctctgct	gtgggcgcaa	1020
ctgacgacgc	agggcaaac	ccatggctgg	ccacatgaaa	tcattcagta	ctattatgga	1080
ctataccccc	ccgcacat	tgcgcagta	tccggggcga	gtctgatggc	cctcctaact	1140
ctggcggcca	catgctgcat	gctggccacc	gcgaggagaa	agtgccctaac	accgtacgcc	1200

-continued

ctgacgccag gagcgggtggt accgttgaca ctgggggtgc tttgctgcgc accgagggcg 1260

aatgca 1266

<210> SEQ ID NO 78

<211> LENGTH: 1266

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- MLV 10A1

<400> SEQUENCE: 78

agtgtaacag agcacttta tgtgtataag gctactagac catacctagc acattgcgcc 60

gattgcgggg acgggtactt ctgctatagc ccagttgcta tcgaggagat ccgagatgag 120

gcgtctgatg gcatgcttaa gatccaagtc tccgccc aaa taggtctgga caaggcaggc 180

acccacgccc acacgaagct ccgatatatg gctggatcatg atgttcagga atctaagaga 240

gattccttga ggggtgtacac gtccgcagcg tgctccatac atgggacgat gggacacttc 300

atcgctgcac actgtccacc aggcgactac ctcaagggtt cggtcgagga cgcagattcg 360

cacgtgaagg catgtaaggt ccaatacaag cacaatccat tgccgggtggg tagagagaag 420

ttcgtggtta gaccacactt tggcgtagag ctgccatgca cctcatacca gctgacaacg 480

gctcccacgg acgaggagat tgacatgcat acaccgccag atataccgga tcgcaccctg 540

ctatcacaga cggcggggcaa cgtcaaaaata acagcaggcg gcaggactat cagggtacaac 600

tgtacctgag gccgtgacaa cgtaggcact accagtactg acaagaccat caacacatgc 660

aagattgacc aatgccatgc tgccgtcacc agccatgaca aatggcaatt tacctctcca 720

tttgttccca gggctgatca gacagctagg aaaggcaagg tacacgttcc gttccctctg 780

actaacgtca cctgccgagt gccgttggt cgagcgccgg atgccaccta tggttaagaag 840

gaggtgaccc tgagattaca ccagatcat ccgacgctct tctcctatag gagtttagga 900

gccgaaccgc acccgtagca ggaatgggtt gacaagttct ctgagcgcat catcccagtg 960

acggaagaag ggattagata ccagtggggc aacaaccgc cggtctgcct gtgggcgcaa 1020

ctgacgacgg agggcaaac ccattggctgg ccacatgaaa tcattcagta ctattatgga 1080

ctataccccg ccgcactat tgccgcagta tccggggcga gtctgatggc cctcctaact 1140

ctggcgccca catgctgcat gctggccacc gcgaggagaa agtgccctaac accgtacgcc 1200

ctgacgccag gagcgggtggt accgttgaca ctgggggtgc tttgctgcgc accgagggcg 1260

aatgca 1266

<210> SEQ ID NO 79

<211> LENGTH: 2030

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Envelope- Ebola

<400> SEQUENCE: 79

atgggtgtta caggaatatt gcagttacct cgtgatcgat tcaagaggac atcattcttt 60

ctttgggtaa ttatcctttt ccaagaaca ttttccatcc cacttgagat catccacaat 120

agcacattac aggttagtga tgcgcacaaa ctgggttgcc gtgacaaact gtcattccaca 180

aatcaattga gatcagttgg actgaatctc gaagggaatg gaggggcaac tgacgtgcca 240

-continued

tctgcaacta aaagatgggg cttcaggtcc ggtgtccac caaagtggt caattatgaa	300
gctggtgaat gggctgaaaa ctgctacaat cttgaaatca aaaaacctga cgggagtgag	360
tgtctaccag cagcgccaga cgggattcgg ggcttcccc ggtgccggt tgtgcacaaa	420
gtatcaggaa cgggaccgtg tgccggagac ttgctctcc acaaagagg tgctttcttc	480
ctgtatgacc gacttgcttc cacagttatc taccgaggaa cgactttcgc tgaagggtgc	540
gttgcatctc tgatactgcc ccaagctaag aaggacttct tcagctcaca ccccttgaga	600
gagccggtca atgcaacgga ggaccctct agtggtact attctaccac aattagatat	660
caagtaccg gttttggaac caatgagaca gagtatttgt tcgaggtga caatttgacc	720
tacgtccaac ttgaatcaag attcacacca cagttttcgc tccagctgaa tgagacaata	780
tatacaagtg gaaaaaggag caataccacg gaaaaactaa tttggaagg caaccccgaa	840
attgatacaa caatcgggga gtgggccttc tgggaaacta aaaaacctc actagaaaaa	900
ttcgagtgag agagttgtct ttcacagctg tatcaaacag agccaaaaac atcagtggtc	960
agagtcgggc gcgaacttct tccgaccag ggaccaacac aacaactgaa gaccacaaaa	1020
tcagtgcttc agaaaattcc tctgcaatgg ttcaagtga cagtcaagga agggaagctg	1080
cagtgctgca tctgacaacc ctggccacaa tctccacgag tcctcaacct cccacaacca	1140
aaccaggtcc ggacaacagc acccacaata caccctgta taaacttgac atctctgagg	1200
caactcaagt tgaacaacat caccgcagaa cagacaacga cagcacagcc tccgacactc	1260
cccccgccac gaccgcagcc ggaccctaa aagcagagaa caccaacacg agcaagggtg	1320
ccgacctctt ggaccccgcc accacaacaa gtccccaaaa ccacagcgag accgctggca	1380
acaacaacac tcatacccaa gataccggag aagagagtg cagcagcggg aagctaggct	1440
taattaccaa tactattgct ggagtcgag gactgatcac aggcgggagg agagctcgaa	1500
gagaagcaat tgtcaatgct caaccctaat gcaaccctaa tttacattac tggactactc	1560
aggatgaagg tgctgcaatc ggactggcct ggataccata tttcgggcca gcagccgagg	1620
gaatttatcat agaggggctg atgcacaatc aagatgggtt aatctgtggg ttgagacagc	1680
tggccaacga gacgactcaa gctcttcaac tgttcctgag agccacaacc gagctacgca	1740
ccttttcaat cctcaaccgt aaggcaattg atttcttctg gcagcgatgg ggccggacat	1800
gccacatttt gggaccggac tgctgtatcg aaccacatga ttggaccaag aacataacag	1860
acaaaattga tcagattatt catgattttg ttgataaaac ccttcggagc cagggggaca	1920
atgacaattg gtggacagga tggagacaat ggataccggc aggtattgga gttacaggcg	1980
ttataattgc agttatcgct ttattctgta tatgcaaat tgtcttttag	2030

<210> SEQ ID NO 80

<211> LENGTH: 389

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Short WPRE sequence

<400> SEQUENCE: 80

aatcaacctc tggattacaa aatttgtgaa agattgactg atattcttaa ctatgttgct	60
ccttttacgc tgtgtggata tgctgcttta atgcctctgt atcatgctat tgcttcccg	120
acggctttcg tttctctctc cttgtataaa tcctgggtgc tgtctcttta tgaggagttg	180

-continued

tgccccgttg tccgtcaacg tggcgtggtg tgctctgtgt ttgtgacgc aacccccact	240
ggctggggca ttgccaccac ctgtcaactc ctttctggga ctttgcgttt cccctcccg	300
atcgccacgg cagaactcat cgccgcctgc cttgcccgct gctggacagg ggctagggtg	360
ctgggcactg ataattccgt ggtgtgtgc	389

<210> SEQ ID NO 81
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 81

taagcagaat tcatgaattt gccaggaaga t	31
------------------------------------	----

<210> SEQ ID NO 82
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 82

ccatacaatg aatggacact aggcggccgc acgaat	36
---	----

<210> SEQ ID NO 83
 <211> LENGTH: 2745
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Gag, Pol, Integrase fragment

<400> SEQUENCE: 83

gaattcatga atttgcagg aagatggaaa ccaaaaatga tagggggaat tggaggtttt	60
atcaaagtaa gacagtatga tcagatactc atagaaatct gcggacataa agctataggt	120
acagtattag taggacctac acctgtcaac ataattggaa gaaatctgtt gactcagatt	180
ggctgcactt taaattttcc cattagtcct attgagactg taccagtaaa attaaagcca	240
ggaatggatg gcccaaaagt taaacaatgg ccattgacag aagaaaaaat aaaagcatta	300
gtagaaattt gtacagaaat ggaaaaggaa ggaaaaattt caaaaattgg gcctgaaaat	360
ccatacaata ctccagtatt tgccataaag aaaaagaca gtactaaatg gagaaaatta	420
gtagatttca gagaacttaa taagagaact caagatttct gggaagtcca attaggaata	480
ccacatctg cagggttaaa acagaaaaaa tcagtaacag tactggatgt gggcgatgca	540
tatttttcag ttcccttaga taaagacttc aggaagtata ctgcatttac catacctagt	600
ataaacaatg agacaccagg gattagatat cagtacaatg tgcttcaca gggatggaaa	660
ggatcaccag caatattcca gtgtagcatg acaaaaatct tagagccttt tagaaaacaa	720
aatccagaca tagtcatcta tcaatacatg gatgatttgt atgtaggatc tgacttagaa	780
atagggcagc atagaacaaa aatagaggaa ctgagacaac atctgttgag gtggggattt	840
accacaccag acaaaaaaca tcagaaagaa cctccattcc tttggatggg ttatgaactc	900
catctgata aatggacagt acagcctata gtgctgccag aaaaggacag ctggactgtc	960
aatgacatac agaaattagt gggaaaattg aattgggcaa gtcagattta tgcagggtt	1020

-continued

aaagtaaggc aattatgtaa acttcttagg ggaaccaaag cactaacaga agtagtacca	1080
ctaacagaag aagcagagct agaactggca gaaaacaggg agattctaaa agaaccggta	1140
catggagtgt attatgacct atcaaaagac ttaatagcag aaatacagaa gcaggggcaa	1200
ggccaatgga catatcaaat ttatcaagag ccatttaaaa atctgaaaac aggaaagtat	1260
gcaagaatga aggggtgccca cactaatgat gtgaacaat taacagaggc agtacaaaaa	1320
atagccacag aaagcatagt aatatgggga aagactccta aatttaaatt acccatataa	1380
aaggaaacat ggggaagcatg gtggacagag tattggcaag ccacctggat tcctgagtgg	1440
gagtttgta ataccctcc cttagtgaag ttatggtacc agttagagaa agaaccata	1500
ataggagcag aaactttcta tgtagatggg gcagccaata gggaaactaa attaggaaaa	1560
gcaggatatg taactgacag aggaagacaa aaagttgtcc ccctaacgga cacaacaaat	1620
cagaagactg agttacaagc aattcatcta gctttgcagg attcgggatt agaagtaaac	1680
atagtgcag actcacaata tgcattggga atcattcaag cacaaccaga taagagtga	1740
tcagagttag tcagtcaaat aatagagcag ttaataaaaa aggaaaaagt ctacctggca	1800
tgggtaccag cacacaaagg aattggagga aatgaacaag tagataaatt ggtcagtgt	1860
ggaatcagga aagtactatt tttagatgga atagataagg cccaagaaga acatgagaaa	1920
tatcacagta attggagagc aatggctagt gattttaacc taccacctgt agtagcaaaa	1980
gaaatagtag ccagctgtga taaatgtcag ctaaaagggg aagccatgca tggacaagta	2040
gactgtagcc caggaatatg gcagctagat tgtacacatt tagaaggaaa agttatcttg	2100
gtagcagttc atgtagccag tggatatata gaagcagaag taattccagc agagacaggg	2160
caagaaacag catacttctt cttaaaatta gcaggaagat ggccagtaaa aacagtacat	2220
acagacaatg gcagcaatgt caccagtact acagttaagg ccgcctgttg gtgggcgggg	2280
atcaagcagg aatttggeat tccctacaat ccccaaagtc aaggagtaat agaattctatg	2340
aataaagaat taaagaaat tataggacag gtaagagatc aggctgaaca tcttaagaca	2400
gcagtacaaa tggcagtatt catccacaat tttaaaagaa aaggggggat tgggggggtac	2460
agtcagggg aaagaatagt agacataata gcaacagaca tacaactaa agaattacaa	2520
aaacaaatta caaaaattca aaattttcgg gtttattaca gggacagcag agatccagtt	2580
tggaaaggac cagcaaagct cctctggaaa ggtgaagggg cagtagtaat acaagataat	2640
agtgacataa aagtagtgcc aagaagaaaa gcaaagatca tcagggatta tggaaaacag	2700
atggcagggtg atgatttgtg ggcaagtaga caggatgagg attaa	2745

<210> SEQ ID NO 84

<211> LENGTH: 1586

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: DNA Fragment containing Rev, RRE and rabbit
beta globin poly A

<400> SEQUENCE: 84

tctagaatgg caggaagaag cggagacagc gacgaagagc tcatcagaac agtcagactc	60
atcaagcttc tctatcaaag caaccacct cccaatcccg aggggacccg acaggcccga	120
aggaatagaa gaagaagggtg gagagagaga cagagacaga tccattcgat tagtgaacgg	180
atccttgga cttatctggg acgatctgag gagcctgtgc ctcttcagct accaccgctt	240

-continued

gagagactta ctcttgattg taacgaggat tgtggaactt ctgggacgca gggggtggga	300
agccctcaaa tattggtgga atctctaca atattggagt caggagctaa agaatagagg	360
agctttgttc cttgggttct tgggagcagc aggaagcact atgggcgcag cgtcaatgac	420
gctgacggta caggccagac aattattgtc tggatatgtg cagcagcaga acaatttgct	480
gagggctatt gaggcgcaac agcatctgtt gcaactcaca gtctggggca tcaagcagct	540
ccaggcaaga atcctggctg tggaaagata cctaaggat caacagctcc tagatctttt	600
tccctctgcc aaaaattatg gggacatcat gaagccctt gagcatctga cttctggcta	660
ataaaggaaa tttattttca ttgcaatagt gtgttggaat tttttgtgtc tctcactcgg	720
aaggacatat gggagggcaa atcatttaaa acatcagaat gagtatttgg tttagagttt	780
ggcaacatat gccatagct ggctgccatg aacaaagggtg gctataaaga ggctcatcagt	840
atatgaaaca gccccctgct gtccattcct tattccatag aaaagccttg acttgagggt	900
agattttttt tataattttgt tttgtgttat ttttttcttt aacatcccta aaattttcct	960
tacatgtttt actagccaga tttttcctcc tctcctgact actcccagtc atagctgtcc	1020
ctcttctctt atgaagatcc ctcgacctgc agcccaagct tggcgtaatc atggtcatag	1080
ctgtttctctg tgtgaaattg ttatccgctc acaattccac acaacatacg agccggaagc	1140
ataaagtgtg aagcctgggg tgccaatga gtgagctaac tcacattaat tgcgttgccg	1200
tcactgcccg ctttccagtc gggaaacctg tcgtgccagc ggatccgcat ctcaattagt	1260
cagcaaccat agtcccggcc ctaactccgc ccatcccgc cctaactccg cccagttccg	1320
cccattctcc gcccctggc tgactaattt tttttattta tgcagaggcc gaggcgcct	1380
cggcctctga gctattccag aagtagtgag gaggcttttt tggaggccta ggcttttgca	1440
aaaagctaac ttgtttattg cagcttataa tgggtacaaa taaagcaata gcatcacaaa	1500
tttcacaaat aaagcatttt tttcactgca ttctagtgtg ggtttgtcca aactcatcaa	1560
tgtatcttat cagcggccgc cccggg	1586

<210> SEQ ID NO 85

<211> LENGTH: 1614

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: DNA fragment containing the CAG
enhancer/promoter/intron sequence

<400> SEQUENCE: 85

acgcgttagt tattaatagt aatcaattac ggggtcatta gttcatagcc catatatgga	60
gttccgcgtt acataactta cggtaaatgg cccgcctggc tgaccgcccc acgacccccg	120
cccattgacg tcaataatga cgtatgttcc catagtaacg ccaataggga ctttccattg	180
acgtcaatgg gtggactatt tacggtaaac tgcccacttg gcagtacatc aagtgtatca	240
tatgccaaagt acgcccccta ttgacgtcaa tgacggtaaa tggcccgctt ggcatatatgc	300
ccagtacatg accttatggg actttcttac ttggcagtac atctacgtat tagtcatcgc	360
tattaccatg ggtcgagggt agccccacgt tctgcttcac tctccccatc tccccccct	420
ccccaccccc aattttgtat ttattttatt tttaattatt ttgtgcagcg atggggcgcg	480
gggggggggg ggcgcgcgcc aggcggggcg gggcggggcg aggggcgggg cggggcgagg	540

-continued

cggagaggtg cggcggcagc caatcagagc ggcgcgctcc gaaagtttcc ttttatggcg	600
aggcggcggc ggcggcggcc ctataaaaag cgaagcgcgc ggcggggcgg agtcgctgcg	660
ttgccttcgc cccgtgcccc gctccgcgcc gcctcgccgc gcccgccccg gctctgactg	720
accgcgttac tcccacaggt gagcgggcgg gacggccctt ctctccggg ctgtaattag	780
cgcttggttt aatgacggct cgtttctttt ctgtggctgc gtgaaagcct taaagggtc	840
cgggagggcc ctttgtgcgg gggggagcgg ctcggggggt gcgtgcgtgt gtgtgtgcgt	900
ggggagcgcc gcgtgcggcc cgcgctgccc ggcggtgtg agcgtgcgg gcgcggcgcg	960
gggctttgtg cgctcccgct gtgcgcgagg ggagcgcggc cgggggcgggt gcccccggt	1020
gcgggggggc tgccaggga acaaaaggct gcgtcggggt gtgtgcgtgg gggggtgagc	1080
agggggtgtg ggcgcggcgg tcgggctgta accccccct gcacccccct ccccgagttg	1140
ctgagcacgg cccggcttcg ggtgcggggc tccgtgcggg gcgtggcgcg gggctcgccg	1200
tgccggcgcg ggggtggcgg caggtggggg tgccggcgcg ggcggggcgg cctcgggcgg	1260
gggagggctc gggggagggg cgcggcggcc ccggagcgcc ggcggtgtc gaggcgcggc	1320
gagccgcagc cattgccttt tatggtaatc gtgcgagagg gcgcagggaac ttcctttgtc	1380
ccaaatctgg cggagccgaa atctgggagg cgcgcgcga cccctctag cgggcgcggg	1440
cgaagcgtg cgcgcgcggc aggaagaaa tgggcgggga gggccttcgt gcgtcgccgc	1500
gccgcctcc cctctccat ctccagcctc ggggctgcg cagggggacg gctgccttcg	1560
ggggggacgg ggcaggcggg ggttcggctt ctggcggtgt accggcggga attc	1614

<210> SEQ ID NO 86

<211> LENGTH: 1531

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: DNA fragment containing VSV-G

<400> SEQUENCE: 86

gaattcatga agtgcctttt gtacttagcc tttttattca ttggggtgaa ttgcaagttc	60
accatagttt ttccacacaa ccaaaaagga aactggaaaa atgttccttc taattacat	120
tattgcccggt caagctcaga tttaaattgg cataatgact taataggcac agccttaca	180
gtcaaaatgc ccaagagtca caaggctatt caagcagacg gttggatgtg tcatgcttcc	240
aaatgggtca ctacttgtga tttccgctgg tatggaccga agtatataac acattccatc	300
cgatccttca ctccatctgt agaacaatgc aaggaaagca tgaacaaac gaaacaagga	360
acttggtgta atccaggctt cctctctcaa agttgtggat atgcaactgt gacggatgcc	420
gaagcagtga ttgtccaggt gactcctcac catgtgctgg ttgatgaata cacaggagaa	480
tgggttgatt cacagtccat caacggaaaa tgcagcaatt acatatgcc cactgtccat	540
aactctacaa cctggcattc tgactataag gtcaaaaggc tatgtgattc taacctcatt	600
tccatggaca tcaccttctt ctccagggac ggagagctat catccctggg aaaggagggc	660
acagggttca gaagtaacta ctttgcttat gaaactggag gcaaggcctg caaaatgcaa	720
tactgcaagc attggggagt cagactccca tcaggtgtct ggttcgagat ggctgataag	780
gatctctttg ctgcagccag attccctgaa tgcccagaag ggtcaagtat ctctgctcca	840
tctcagacct cagtggatgt aagtctaatt caggacgttg agaggatctt ggattattcc	900

-continued

ctctgccaaag aaacctggag caaaatcaga gcgggtcttc caatctctcc agtggatctc	960
agctatcttg ctctaaaaa cccaggaacc ggtcctgctt tcaccataat caatgggtacc	1020
ctaaaatact ttgagaccag atacatcaga gtcgatattg ctgctccaat cctctcaaga	1080
atggctcgaa tgatcagtgg aactaccaca gaaagggaaac tgtgggatga ctgggcacca	1140
tatgaagacg tggaaattgg acccaatgga gttctgagga ccagttcagg atataagttt	1200
cctttatata tgattggaca tggatgttg gactccgac ttcattcttag ctcaaaggct	1260
cagggtgttcg aacatctctc cattcaagac gctgcttcgc aacttctga tgatgagagt	1320
ttattttttg gtgatactgg gctatccaaa aatccaatcg agcttgtaga aggttggttc	1380
agtagttgga aaagctctat tgctctttt ttctttatca tagggttaat cattggacta	1440
ttcttggttc tccgagttgg tatccatctt tgcattaaat taaagcacac caagaaaaga	1500
cagatttata cagacataga gatgagaatt c	1531

<210> SEQ ID NO 87

<211> LENGTH: 1227

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Helper plasmid containing RRE and rabbit beta globin poly A

<400> SEQUENCE: 87

tctagaagga gctttgttcc ttgggttctt gggagcagca ggaagcacta tgggcgcagc	60
gtcaatgacg ctgacggtac aggccagaca attattgtct ggtatagtgc agcagcagaa	120
caatttctg agggctattg aggcgcaaca gcatctgttg caactcacag tctggggcat	180
caagcagctc caggcaagaa tcctggctgt ggaaagatac ctaaaggatc aacagctcct	240
agatcttttt cctctgcca aaaattatgg ggacatcatg aagcccttg agcatctgac	300
ttctggctaa taaaggaaat ttattttcat tgcaatagtg tgttggaatt ttttgtgtct	360
ctcactcgga aggacatag ggagggcaaa tcatttaaaa catcagaatg agtatttgg	420
ttagagtttg gcaacatag ccatatgctg gctgccatga acaaagggtg ctataaagag	480
gtcatcagta tatgaaacag cccctgctg tccattcctt attccataga aaagccttga	540
cttgaggta gatttttttt atattttgtt ttgtgttatt tttttctta acatccctaa	600
aattttcctt acatgtttta ctagccagat ttttcctct cctctgacta ctcccagtca	660
tagctgtccc tcttctctta tgaagatccc tcgacctgca gcccaagctt ggcgtaatca	720
tggatcatagc tgtttcctgt gtgaaattgt tatccgctca caattccaca caacatacga	780
gccggaagca taaagtgtaa agcctgggt gcctaagag tgagctaact cacattaatt	840
gcgttgccgt cactgcccgc ttccagctg ggaaacctgt cgtgccagcg gatccgcac	900
tcaattagtc agcaaccata gtcccgcgcc taactccgcc catcccgcgc ctaactccgc	960
ccagttccgc ccattctccg ccccatggct gactaatttt ttttatttat gcagaggccg	1020
aggccgcctc ggctctgag ctattccaga agtagtgagg aggccttttt ggaggcctag	1080
gcttttgcaa aaagctaact tgtttattgc agcttataat gggtacaaat aaagcaatag	1140
catcacaat ttcacaaata aagcattttt ttcaactgcat tctagttgtg gtttgtccaa	1200
actcatcaat gtatcttata acccggg	1227

-continued

```

<210> SEQ ID NO 88
<211> LENGTH: 884
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RSV promoter and HIV Rev

<400> SEQUENCE: 88

caattgcat gtacgggcca gatatacgcg tatctgaggg gactaggggtg tgtttaggcg      60
aaaagcgggg cttcggttgt acgcgggttag gagtcccctc aggatatagt agtttcgctt      120
ttgcataggg agggggaaat gtagtcttat gcaatacact tgtagtcttg caacatggta      180
acgatgagtt agcaacatgc cttacaagga gagaaaaagc accgtgcatg ccgattggtg      240
gaagtaaggt ggtacgatcg tgccttatta ggaaggcaac agacagggtct gacatggatt      300
ggacgaacca ctgaattccg cattgcagag ataattgtat ttaagtgcct agctcgatac      360
aataaacgcc atttgacatc tcaccacatt ggtgtgcacc tccaagctcg agctcgttta      420
gtgaaccgtc agatcgccctg gagacgcatc ccacgctgtt ttgacctcca tagaagacac      480
cgggaccgat ccagcctccc ctggaagcta gcgattaggg atctcctatg gcaggaagaa      540
gcggagacag cgacgaagaa ctctcaagg cagtcagact catcaagttt ctctatcaaa      600
gcaaccacc tcccaatccc gaggggaccc gacaggcccg aaggaataga agaagaaggt      660
ggagagagag acagagacag atccattcga ttagtgaacg gatccttagc acttatctgg      720
gacgatctgc ggagcctgtg cctcttcagc taccaccgct tgagagactt actcttgatt      780
gtaacgagga ttgtggaact tctgggacgc aggggggtggg aagccctcaa atattggtgg      840
aatctctac aatattggag tcaggagcta aagaatagtc taga                        884

```

```

<210> SEQ ID NO 89
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence

<400> SEQUENCE: 89

atggcaggaa gaagcggag                        19

```

```

<210> SEQ ID NO 90
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: shRNA sequence

<400> SEQUENCE: 90

atggcaggaa gaagcggagt tcaagagact ccgcttcttc ctgccatttt tt      52

```

```

<210> SEQ ID NO 91
<211> LENGTH: 279
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: H1 promoter and shRT sequence

<400> SEQUENCE: 91

gaacgctgac gtcatcaacc cgctccaagg aatcgcgggc ccagtgtcac taggcgggaa      60
caccagcgc gcgtgcgcc tggcaggaag atggctgtga gggacagggg agtggcgccc      120

```

-continued

tgcaatatatt gcatgtcgct atgtgttctg ggaaatcacc ataaacgtga aatgtctttg	180
gatttgggaa tcttataagt tctgtatgag accacttgga tccgcggaga cagcgacgaa	240
gagcttcaag agagctcttc gtcgctgtct ccgcttttt	279

<210> SEQ ID NO 92
 <211> LENGTH: 275
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: H1 CCR5 sequence

<400> SEQUENCE: 92

gaacgctgac gtcatcaacc cgctccaagg aatcgcgggc ccagtgtcac taggcgggaa	60
caccagcgcg gcgtgcgccc tggcaggaag atggctgtga gggacagggg agtggcgccc	120
tgcaatatatt gcatgtcgct atgtgttctg ggaaatcacc ataaacgtga aatgtctttg	180
gatttgggaa tcttataagt tctgtatgag accacttgga tccgtgtcaa gtccaatcta	240
tgttcaagag acatagattg gacttgacac ttttt	275

<210> SEQ ID NO 93
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 93

aggaattgat ggcgagaagg	20
-----------------------	----

<210> SEQ ID NO 94
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 94

ccccaagaa ggtcaaggta atca	24
---------------------------	----

<210> SEQ ID NO 95
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 95

agcgcggtca cagcttca	18
---------------------	----

<210> SEQ ID NO 96
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (20)..(20)
 <223> OTHER INFORMATION: n = p

<400> SEQUENCE: 96

-continued

ggcgacgtag cacagcttcn 20

<210> SEQ ID NO 97
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AGT103 CCR5 miR30

<400> SEQUENCE: 97

tgtaaaactga gcttgctcta 20

<210> SEQ ID NO 98
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AGT103-R5-1

<400> SEQUENCE: 98

tgtaaaactga gcttgctcgc 20

<210> SEQ ID NO 99
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AGT103-R5-2

<400> SEQUENCE: 99

catagattgg acttgacac 19

<210> SEQ ID NO 100
<211> LENGTH: 642
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CAG promoter

<400> SEQUENCE: 100

tagttattaa tagtaatcaa ttacgggggc attagtcat agcccatata tggagttccg 60

cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc ccgcccatt 120

gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca 180

atgggtggac tatttacggt aaactgccca cttggcagta catcaagtgt atcatatgcc 240

aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta 300

catgacctta tgggactttc ctacttgga gtacatctac gtattagtca tcgctattac 360

catgggtcga ggtgagcccc acgtttctgt tcaactctccc catctcccc ccctccccac 420

ccccaat ttt gtatttat ttttttaaat tttttgtgc agcgatgggg gcgggggggg 480

gggggggcgc cgccaggcgg ggcggggcgg ggcgaggggc gggcgggggc gaggcggaga 540

ggtgcggcgg cagccaatca gagcggcgcg ctccgaaagt ttccttttat ggcgaggcgg 600

cggcggcggc ggccctataa aaagcgaagc gcgcggcggc cg 642

<210> SEQ ID NO 101
<211> LENGTH: 217
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: H1 element

<400> SEQUENCE: 101

gaacgctgac gtcatcaacc cgctccaagg aatcgcgggc ccagtgtcac taggcgggaa	60
caccacagcg cgctgcgccc tggcaggaag atggctgtga gggacagggg agtggcgccc	120
tgcaatatTT gcattgcgct atgtgttctg ggaaatcacc ataaacgtga aatgtctttg	180
gatttgggaa tcttataagt tctgtatgag accactt	217

<210> SEQ ID NO 102
<211> LENGTH: 250
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' LTR

<400> SEQUENCE: 102

tggaagggct aattcactcc caacgaagat aagatctgct ttttgcctgt actgggtctc	60
tctgtttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta	120
agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgctctg ttgtgtgact	180
ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtagta	240
gttcatgtca	250

<210> SEQ ID NO 103
<211> LENGTH: 243
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 7SK promoter

<400> SEQUENCE: 103

ctgcagtatt tagcatgcc caccatctg caaggcattc tggatagtgt caaacagcc	60
ggaaatcaag tccgtttatc tcaaaactta gcattttggg aataaatgat atttgctatg	120
ctggttaaat tagatttttag ttaaatctcc tgctgaagct ctagtacgat aagcaacttg	180
acctaagtgt aaagttgaga tttccttcag gtttatatag cttgtgcgcc gcctggctac	240
ctc	243

<210> SEQ ID NO 104
<211> LENGTH: 132
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miR155 Tat

<400> SEQUENCE: 104

ctggaggctt gctgaaggct gtatgtgtgc cgcttcttcc tgccataggg ttttgccac	60
tgactgaccc tatggggaag aagcggacag gacacaaggc ctgttactag cactcacatg	120
gaacaaatgg cc	132

<210> SEQ ID NO 105
<211> LENGTH: 3992
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pRSV Rev

-continued

<400> SEQUENCE: 105

agcgcccaat acgcaaacgg cctctccccg cgcgttgccc gattcattaa tgcagctggc	60
acgacaggtt tcccgactgg aaagcgggca gtgagcgcaa cgcaattaat gtgagttagc	120
tcactcatta ggcaccccag gctttacact ttatgcttcc ggctcgtagt ttgtgtggaa	180
ttgtgagcgg ataacaattt cacacaggaa acagctatga ccatgattac gaattcgatg	240
tacggggcag atatacgcgt atctgagggg actaggggtg gtttaggcga aaagcggggc	300
ttcggttgta cgcggttagg agtccctca ggatatagta gtttcgcttt tgcataggga	360
gggggaaatg tagtcttatg caatacactt gtagtcttgc aacatggtaa cgatgagtta	420
gcaacatgcc ttacaaggag agaaaaagca ccgtgcacgc cgattggtgg aagtaagggtg	480
gtacgatcgt gccttattag gaaggcaaca gacaggtctg acatggattg gacgaaccac	540
tgaattccgc attgcagaga taattgtatt taagtgccta gctcgatata ataaacgcca	600
tttgaccatt caccacattg gtgtgcacct ccaagctcga gctcgtttag tgaaccgtca	660
gatcgccctg agacgccatc cacgctgttt tgacctccat agaagacacc gggaccgatc	720
cagcctcccc tcgaagctag tcgattaggc atctcctatg gcaggaagaa gcggagacag	780
cgacgaagac ctctcaagg cagtcagact catcaagttt ctctatcaaa gcaaccacc	840
tcccaatccc gaggggaccc gacaggcccg aaggaataga agaagaaggt ggagagagag	900
acagagacag atccattcga ttagtgaacg gatccttagc acttatctgg gacgatctgc	960
ggagcctgtg cctcttcagc taccaccgct tgagagactt actcttgatt gtaacgagga	1020
ttgtggaact tctgggacgc aggggggtggg aagccctcaa atattggtgg aatctctac	1080
aatattggag tcaggagcta aagaatagtg ctgttagctt gctcaatgcc acagctatag	1140
cagtagctga ggggacagat aggggttatag aagtagtaca agaagcttgg cactggccgt	1200
cgttttatcat gatctgagcc tgggagatct ctggctaact agggaaacca ctgcttaagc	1260
ctcaataaag cttgccttga gtgcttcaag tagtgtgtgc ccgtctgttg tgtgactctg	1320
gtaactagag atcacaagc aaccatagta cgcgccctgt agcggcgcat taagcgcggc	1380
gggtgtggtg gttacgcgca gcgtgaccgc tacacttgcc agcgcctag cgcgcgtcc	1440
tttcgcttcc ttccttctct ttctcgccac gttcgccggc tttcccgctc aagctctaaa	1500
tcggggggctc cctttagggt tccgatttag tgctttacgg cacctcgacc ccaaaaaact	1560
tgatttgggt gatgggtcac gtagtgggccc atcgccctga tagacggttt ttcgcccttt	1620
gacgttggag tccacgttct ttaatagtgg actcttgctc caaactggaa caaactcaa	1680
ccctatctcg ggctattctt ttgatttata agggattttg ccgatttcgg cctattgggt	1740
aaaaaatgag ctgatttaac aaaaatttaa cgcgaatttt aacaaaatat taacgtttac	1800
aattttatgg tgcactctca gtacaatctg ctctgatgcc gcatagttaa gccagccccg	1860
acaccgcgca acaccgcgtg acgcgcctg acgggcttgt ctgctcccg catccgctta	1920
cagacaagct gtgaccgtct ccgggagctg catgtgtcag aggttttcac cgtcacacc	1980
gaaacgcgcg agacgaaagg gcctcgtgat acgcctattt ttataggtta atgtcatgat	2040
aataatggtt tcttagacgt caggtggcac ttttcgggga aatgtgcgcg gaaccctat	2100
ttgtttatct ttctaaatac attcaaatat gtatccgctc atgagacaat aaccctgata	2160
aatgcttcaa taatattgaa aaaggaagag tatgagtatt caacatttcc gtgtcgccct	2220

-continued

tattcccttt tttgcggcat tttgccttcc tgtttttgct caccagaaa cgctggtgaa	2280
agtaaaagat gctgaagatc agtgggtgc acgagtgggt tacatcgaac tggatctcaa	2340
cagcggtaag atccttgaga gttttcgccc cgaagaacgt tttccaatga tgagcacttt	2400
taaagtctcg ctatgtggcg cgttattatc ccgtattgac gccgggcaag agcaactcgg	2460
tcgccgcata cactattctc agaatgactt ggttgagtac tcaccagtca cagaaaagca	2520
tcttacggat ggcatgacag taagagaatt atgcagtgcg gccataacca tgagtataaa	2580
cactgcggcc aacttacttc tgacaacgat cggaggacgg aaggagctaa ccgctttttt	2640
gcacaacatg ggggatcatg taactcgcct tgatcgttgg gaaccggagc tgaatgaagc	2700
cataccaaac gacgagcgtg acaccacgat gcctgtagca atggcaacaa cgttgcgcaa	2760
actattaact ggcgaaactac ttactctagc ttcccgga caattaatag actggatgga	2820
ggcggataaa gttgcaggac cacttctcgc ctccgcccct ccgctgggtt ggtttattgc	2880
tgataaatct ggagccgggtg agcgtgggtc tcgcggtatc attgcagcac tggggccaga	2940
tggttaagccc tcccgtagcg tagttatcta cagcagggg agtcaggcaa ctatggatga	3000
acgaaataga cagatcgtg agataggtgc ctactgatt aagcattggt aactgtcaga	3060
ccaagtttac tcatatatac tttagattga tttaaaactt catttttaaat ttaaaaggat	3120
ctagtggaag atcctttttg ataatctcat gaccaaactc ccttaacgtg agttttcgtt	3180
ccactgagcg tcagaccccg tagaaaagat caaaggatct tcttgagatc ctttttttct	3240
gcgcgtaatc tgctgcttgc aaacaaaaaa accaccgcta ccagcgggtg tttgtttgcc	3300
ggatcaagag ctaccaactc tttttccgaa ggtaactggc ttcagcagag cgcagatacc	3360
aaatactgtc cttctagtgt agccgtagtt aggccaccac ttcaagaact ctgtagcacc	3420
gcctacatac ctgcctctgc taatcctgtt accagtggct gctgccagtg gcgataagtc	3480
gtgtcttacc ggggttgact caagacgata gttaccggat aaggcgcagc ggtcgggctg	3540
aacgggggggt tcgtgcacac agcccagctt ggagcgaacg acctacaccg aactgagata	3600
cctacagcgt gagctatgag aaagcggcac gcttcccgaa gggagaaaagg cggacaggta	3660
tccgtaagc ggcagggtcg gaacaggaga gcgcacgagg gagcttcag ggggaaacgc	3720
ctggtatctt tatagtcctg tcgggtttcg ccacctctga cttgagcgtc gatttttgtg	3780
atgctcgtca gggggggcga gcctatggaa aaacgccagc aacgcggcct ttttacgggt	3840
cctggccttt tgctggcctt ttgctcacat gttctttcct gcgttatccc ctgattctgt	3900
ggataaccgt attaccgctt ttgagtgagc tgataccgct cggcgcagcc gaacgaccga	3960
gcgcagcgag tcagtgcgag aggaagcgga ag	3992

<210> SEQ ID NO 106

<211> LENGTH: 6363

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pCMV-VSV-G

<400> SEQUENCE: 106

gagcttggcc cattgcatac gttgtatcca tatcataata tgtacattta tattggctca	60
tgtccaacat taccgccatg ttgacattga ttattgacta gttattaata gtaatcaatt	120
acgggggtcat tagttcatag cccatatatg gagttccgcg ttacataact tacggtaaat	180

-continued

ggcccgccctg gctgaccgcc caacgacccc cgcccattga cgtcaataat gacgtatgtt	240
cccatagtaa cgccaatagg gactttccat tgacgtcaat ggggtggagta ttacggtaa	300
actgcccact tggcagtaca tcaagtgtat catatgccaa gtacgcccc tattgacgtc	360
aatgacggta aatggccgc ctggcattat gcccagtaca tgacctatg ggactttcct	420
acttggcagt acatctacgt attagtcac gctattacca tggatgatgcg gttttggcag	480
tacatcaatg ggcgtggata gcggtttgac tcacggggat ttccaagtct ccacccatt	540
gacgtcaatg ggagtttgtt ttggcaccaa aatcaacggg actttccaaa atgtcgtaac	600
aactccgccc cattgacgca aatggcggtt aggcgtgtac ggtgggaggt ctatataagc	660
agagctcggt tagtgaaccg tcagatcgcc tggagacgcc atccacgctg ttttgacctc	720
catagaagac accgggaccg atccagcctc cggtcgaccg atcctgagaa cttcagggtg	780
agtttgggga cccttgattg ttctttcttt ttcgctattg taaaattcat gttatatgga	840
gggggcaaaag ttttcagggt gttgtttaga atgggaagat gtcccttgta tcaccatgga	900
ccctcatgat aattttgttt ctttcacttt ctactctgtt gacaaccatt gtctcctctt	960
attttctttt cattttctgt aactttttcg ttaaacctta gcttgcatth gtaacgaatt	1020
tttaaattea cttttgttta ttgtcagat tgtaagtact ttctctaate actttttttt	1080
caaggcaate agggatatatt atattgtact tcagcacagt tttagagaac aattgttata	1140
attaatgat aaggtagaat atttctgcat ataaattctg gctggcgtgg aaatattctt	1200
attggtagaa acaactacac cctggtcacc atcctgcctt tctctttatg gttacaatga	1260
tatacactgt ttgagatgag gataaaatac tctgagtcca aaccgggccc ctctgctaac	1320
catgttcattg ccttcttctc ttctctacag ctctgggcca acgtgctggg tgtgtgctg	1380
tctcatcatt ttggcaaaaga attcctcgac ggatccgcca tgaagtgcct tttgtactta	1440
gcctttttat tcattggggg gaattgcaag ttcaccatag tttttccaca caacaaaaa	1500
ggaaactgga aaaatgttcc ttctaattac cattattgcc cgtcaagctc agattttaat	1560
tggcataatg acttaatagg cacagcctta caagtcaaaa tgcccaagag tcacaaggct	1620
attcaagcag acggttggat ggtgcatgct tccaaatggg tcaactactg tgatttcgcg	1680
tggataggac cgaagtatat aacacattcc atccgatcct tcaactcacc tgtagaacaa	1740
tgcaaggaaa gcattgaaca aacgaaacaa ggaacttggc tgaatccagg ctccctcct	1800
caaagtgtg gatatgcaac tgtgacggat gccgaagcag tgattgtcca ggtgactcct	1860
caccatgtgc tgggtgatga atacacagga gaatgggttg attcacagtt catcaacgga	1920
aatgcagca attacatatg cccactgtc cataactcta caacctggca ttctgactat	1980
aagggtcaaa ggcctatgtga ttctaaccct atttccatgg acatcacctt cttctcagag	2040
gacggagagc tatcatccct gggaaaggag ggcacagggt tcagaagtaa ctactttgct	2100
tatgaaactg gaggcaaggc ctgcaaaatg caatactgca agcattgggg agtcagactc	2160
ccatcagggt tctggttoga gatggctgat aaggatctct ttgctgcagc cagattccct	2220
gaatgcccg aagggtcaag tatctctgct ccatctcaga cctcagtgga tgtaagtcta	2280
attcaggacg ttgagaggat cttggattat tccctctgcc aagaacctg gagcaaaatc	2340
agagcgggtc ttccaatctc tccagtggtat ctacgtatc ttgctcctaa aaaccagga	2400
accggtcctg ctttcacat aatcaatggt accctaaaat actttgagac cagatacatc	2460

-continued

agagtcgata	ttgctgctcc	aatcctctca	agaatggctg	gaatgatcag	tggaactacc	2520
acagaaaggg	aactgtggga	tgactgggca	ccatatgaag	acgtggaaat	tggaaccaat	2580
ggagttctga	ggaccagtcc	aggatataag	tttcctttat	acatgattgg	acatggtatg	2640
ttggactccg	atcttcatct	tagctcaaag	gctcagggtg	tcgaacatcc	tcacattcaa	2700
gacgtgctt	cgcaacttcc	tgatgatgag	agtttatctt	ttggtgatac	tggtgctatc	2760
aaaaatccaa	tcgagcttgt	agaagggttg	ttcagtagtt	ggaaaagctc	tattgcctct	2820
ttttctctta	tcataggggt	aatcattgga	ctattcttgg	ttctccgagt	tggtatccat	2880
ctttgcatta	aattaaagca	caccaagaaa	agacagattt	atacagacat	agagatgaac	2940
cgacttgga	agtataagg	atccgtcgag	gaattcactc	ctcaggtgca	ggctgcctat	3000
cagaagggtg	tggtggtgt	ggccaatgcc	ctggctcaca	aataccactg	agatcttttt	3060
ccctctgcca	aaaattatgg	ggacatcatg	aagccccttg	agcatctgac	ttctggctaa	3120
taaaggaaat	ttattttcat	tgcaatagtg	tggtggaatt	ttttgtgtct	ctcactcgga	3180
aggacatatg	ggaggggcaa	tcatttataa	catcagaatg	agtatttggt	ttagagtttg	3240
gcaacatatg	cccatatgct	ggctgccatg	aacaaaggtt	ggctataaag	aggtcatcag	3300
tatatgaaac	agccccctgc	tgctcattcc	ttattccata	gaaaagcctt	gacttgaggt	3360
tagatttttt	ttatatattg	ttttgtgtta	tttttttctt	taacatccct	aaaaatttcc	3420
ttacatgttt	tactagccag	atttttcttc	ctctcctgac	tactccagct	catagctgtc	3480
cctcttctct	tatggagatc	cctcgacgga	tcggccgcaa	ttcgtaatca	tgctcatagct	3540
gtttcctgtg	tgaattgtgt	atccgctcac	aattccacac	aacatacgag	ccggaagcat	3600
aaagtgtaaa	gcctgggggtg	cctaattgagt	gagctaactc	acattaattg	cgttgcgctc	3660
actgcccgc	ttccagtcgg	gaaacctgtc	gtgccagctg	cattaatgaa	tcggccaacg	3720
cgcggggaga	ggcggtttgc	gtattgggcg	ctcttccgct	tcctcgctca	ctgactcgct	3780
gcgctcggtc	gttcgggtgc	ggcgagcggt	atcagctcac	tcaaaggcgg	taatacgggt	3840
atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	gcaaaaggcc	agcaaaaggc	3900
caggaaacct	aaaaaggcgg	cgttgctggc	gtttttccat	aggtccgccc	ccctgacga	3960
gcacacaaaa	aatcgacgct	caagtcagag	gtggcgaaac	ccgacaggac	tataaagata	4020
ccaggcggtt	ccccctggaa	gctccctcgt	gcgctctcct	gttccgaccc	tgccgcttac	4080
cggataacct	tccgcttttc	tcctctgggg	aagcgtggcg	ctttctcata	gctcacgctg	4140
taggtatctc	agttcggtgt	aggctgttcg	ctccaagctg	ggctgtgtgc	acgaaccccc	4200
cgttcagccc	gaccgctgcg	ccttatccgg	taactatcgt	cttgagtcca	acccggttaag	4260
acacgactta	tcgccactgg	cagcagccac	tggtaacagg	attagcagag	cgaggatatgt	4320
aggcggtgct	acagagttct	tgaagtgggt	gcctaactac	ggctacacta	gaagaacagt	4380
atttggtatc	tgcgctctgc	tgaagccagt	taccttcgga	aaaagagttg	gtagctcttg	4440
atccggcaaa	caaaccacgg	ctggtagcgg	tggttttttt	gtttgcaagc	agcagattac	4500
gcgcagaaaa	aaaggatctc	aagaagatcc	tttgatcttt	tctacggggg	ctgacgctca	4560
gtggaacgaa	aactcacggt	aagggatttt	ggctcatgaga	ttatcaaaaa	ggatcttcac	4620
ctagatcctt	ttaaatataa	aatgaagttt	taaatcaatc	taaagtatat	atgagtaaac	4680
ttggtctgac	agttaccaat	gcttaatcag	tgaggcacct	atctcagcga	tctgtctatt	4740

-continued

tcggttcaccc atagttgctt gactccccgt cgtgtagata actacgatac gggagggcctt	4800
accatctggc cccagtgtgt caatgatacc gcgagaccca cgctcaccgg ctccagattt	4860
atcagcaata aaccagccag ccggaagggc cgagcgagca agtgggtcctg caactttatc	4920
cgcctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccagttaa	4980
tagtttgccg aacgttgttg ccattgtctac aggcacgtgt gtgtcacgct cgctgtttgg	5040
tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat ccccatgtt	5100
gtgcaaaaaa gcggttagct ccttcgggtcc tccgatcgtt gtcagaagta agttggccgc	5160
agtgttatca ctcatggta tggcagcact gcataattct cttactgtca tgccatccgt	5220
aagatgcttt tctgtgactg gtgagtactc aaccaagtca ttctgagaat agtgtatgct	5280
gcgaccgagt tgctcttgcc cggcgctaat acgggataat accgcgccac atagcagaac	5340
tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc	5400
gctgttgaga tccagttcga tgtaacccac tcgtgcaccc aactgatctt cagcatcttt	5460
tactttcacc agcgtttctg ggtgagcaaa aacaggaagg caaaatgccg caaaaaagg	5520
aataaggcgc acacggaaat gttgaatact catactcttc ctttttcaat attattgaag	5580
catttatcag ggttatgtc tcattgagcg atacatat ttgaatgtattt agaaaaataa	5640
acaaataggg gttccgcgca catttccccg aaaagtgcc cctaaattgt aagcgtaaat	5700
attttgttaa aattcgcgtt aaatttttgt taaatcagct cattttttaa ccaataggcc	5760
gaaatcggca aaatccctta taaatcaaaa gaatagaccg agatagggtt gagtgttgtt	5820
ccagtttggg acaagagtcc actattaaag aacgtggact ccaacgtcaa agggcgaaaa	5880
accgtctatc agggcgatgg cccactacgt gaaccatcac cctaatcaag ttttttggg	5940
tcgaggtgcc gtaaaagcact aaatcggaac cctaaaggga gccccgatt tagagcttga	6000
cgggggaaagc cggcgaaagt ggcgagaaaag gaagggaaga aagcgaaagg agcgggcgct	6060
agggcgctgg caagtgtagc ggtcacgctg cgcgtaacca ccacaccgc cgcgcttaat	6120
gcgcgcgtac agggcgcgtc ccattcgcca ttcagggtgc gcaactgttg ggaagggcga	6180
tcggtgcggg cctcttcgct attacgccag ctggcgaaaag ggggatgtgc tgcaaggcga	6240
ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagttag	6300
cgcgcgtaat acgactcact ataggcgcaa ttggagctcc accgcggtgg cgcccgctct	6360
aga	6363

<210> SEQ ID NO 107

<211> LENGTH: 9496

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PSPAX2 delta Rev

<400> SEQUENCE: 107

gtcgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata	60
gccccatat ggagttccgc gttacataac ttacggtaaa tggcccgctt ggctgaccgc	120
ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag	180
ggactttcca ttgacgtcaa tgggtgggact atttacggta aactgcccac ttggcagtac	240
atcaagtgtg tcatatgcca agtacgcccc ctattgacgt caatgacggt aaatggcccc	300

-continued

cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg	360
tattagtcac cgctattacc atgggtcgag gtgagcccca cgttctgctt cactctcccc	420
atctcccccc cctccccacc cccaattttg tattttattta ttttttaatt attttgtgca	480
gcgatggggg cggggggggg gggggcgcg cccaggcggg gcggggcggg gcgagggggc	540
gggcggggcg aggcgggagag gtgcggcggc agccaatcag agcggcgcg tccgaaagt	600
tccttttatg gcgaggcggc ggcggcggcg gccctataaa aagcgaagcg cgcggcgggc	660
gggagtcgct gcgttgccct cgcctcgtc cccgctccg gccgcctcgc gcgcgcgcgc	720
ccggctctga ctgaccgcgt tactcccaca ggtgagcggg cgggacggcc cttctcctcc	780
gggctgtaat tagcgcttgg tttaatgacg gctcgtttct tttctgtggc tgcgtgaaag	840
ccttaaaggg ctccgggagg gccctttgtg cgggggggag cggctcgggg ggtgcgtgcg	900
tgtgtgtgtg cgtggggagc gccgcgtgcg gcccgcgctg cccggcggtt gtgagcgctg	960
cgggcgcggc gcggggcttt gtgcgctccg cgtgtgcgcg aggggagcgc ggccgggggc	1020
ggtgccccgc ggtgcggggg ggtgcgaggg ggaacaaagg ctgcgtgcgg ggtgtgtgcg	1080
tgggggggtg agcagggggg gtgggcgcgg cggtcgggct gtaaccccc cctgcacccc	1140
cctccccgag ttgctgagca cggcccggtt tcgggtgcgg ggctccgtgc ggggcgtggc	1200
gcggggctcg ccgtgcggg cgggggggtg cggcagggtg ggggtgcggg cggggcgggg	1260
ccgcctcggg ccggggaggg ctccggggag gggcgcgcg gccccggagc gccggcggtt	1320
gtcgaggcgc ggcgagcgc agccattgcc ttttatggtg atcgtgcgag agggcgagc	1380
gacttccctt gtcccaaatc tggcggagcc gaaatctggg aggcgcgcgc gcacccctc	1440
tagcgggcgc gggcgaagcg gtgcggcgcc ggcaggaagg aaatgggcgg ggagggcctt	1500
cgtgcgtcgc cgcgcgcgcg tcccccttc catctccagc ctccgggctg ccgcaggggg	1560
acggtgcctt tcggggggga cggggcaggg cgggggttcg cttctggcgt gtgaccggcg	1620
gctctagagc ctctgctaac catgttcatt ccttcttctt tttcctacag ctctgggca	1680
acgtgctggg tattgtgctg tctcatcatt ttggcaaaga attcgggcgc gccgcgttga	1740
cgcgcacggc aagaggcgag gggcggcgac tgggtagaga tgggtgcgag agcgtcagta	1800
ttaaagcggg gagaattaga tcgatgggaa aaaattcggg taaggccagg gggaaagaaa	1860
aaatataaat taaaacatat agtatgggca agcagggagc tagaacgatt cgcagttaat	1920
cctggcctgt tagaaacatc agaaggctgt agacaaatac tgggacagct acaaccatcc	1980
cttcagacag gatcagaaga acttagatca ttatataata cagtagcaac cctctattgt	2040
gtgcatcaaa ggatagagat aaaagacacc aagggaagctt tagacaagat agagggaag	2100
caaaacaaaa gtaagaaaaa agcacagcaa gcagcagctg acacaggaca cagcaatcag	2160
gtcagccaaa attaccctat agtcagaaac atccaggggc aaatggtaca tcaggccata	2220
tcacctagaa ctttaaatgc atgggtaaaa gtagtagaag agaaggcttt cagcccagaa	2280
gtgataccca tgttttcagc attatcagaa ggagccaccc cacaagattt aaacaccatg	2340
ctaaacacag tgggggggaca tcaagcagcc atgcaaatgt taaaagagac catcaatgag	2400
gaagctgcag aatgggatag agtgcatcca gtgcagcag ggctatttgc accaggccag	2460
atgagagaac caaggggaag tgacatagca ggaactacta gtacccttca ggaacaaata	2520
ggatggatga cacataatcc acctatccca gtaggagaaa tctataaaag atggataatc	2580

-continued

ctgggattaa	ataaaatagt	aagaatgtat	agccctacca	gcattctgga	cataagacaa	2640
ggaccaaagg	aaccctttag	agactatgta	gaccgattct	ataaaactct	aagagccgag	2700
caagcttcac	aagaggtaaa	aaattggatg	acagaaacct	tggtgggcc	aaatgcgaac	2760
ccagattgta	agactatttt	aaaagcattg	ggaccaggag	cgacactaga	agaaatgatg	2820
acagcatgtc	agggagtggg	gggacccggc	cataaagcaa	gagttttggc	tgaagcaatg	2880
agccaagtaa	caaatccagc	taccataatg	atacagaaag	gcaattttag	gaaccaaaga	2940
aagactgtta	agtgtttcaa	ttgtggcaaa	gaagggcaca	tagccaaaaa	ttgcagggcc	3000
cctaggaaaa	agggtctgtg	gaaatgtgga	aaggaaggac	accaaataaa	agattgtact	3060
gagagacagg	ctaatttttt	agggaagatc	tggtcttccc	acaaggggaag	gccagggaat	3120
tttcttcaga	gcagaccaga	gccaacagcc	ccaccagaag	agagcttcag	gtttggggaa	3180
gagacaacaa	ctccctctca	gaagcaggag	ccgatagaca	aggaactgta	tccttttagct	3240
tccctcagat	cactctttgg	cagcgacccc	tcgtcacaat	aaagataggg	gggcaattaa	3300
aggaagctct	attagatata	ggagcagatg	atacagtatt	agaagaaatg	aatttgccag	3360
gaagatggaa	acaaaaaatg	atagggggaa	ttggagggtt	tatcaaagta	ggacagtatg	3420
atcagatact	catagaaatc	tcggtgacata	aagctatagg	tacagtatta	gtaggacctt	3480
cacctgtcaa	cataattgga	agaaatctgt	tgactcagat	tggtgcact	ttaaattttc	3540
ccattagtcc	tattgagact	gtaccagtaa	aattaaagcc	aggaatggat	ggcccaaaag	3600
ttaaacaatg	gccattgaca	gaagaaaaaa	taaaagcatt	agtagaaatt	tgtacagaaa	3660
tggaagagg	aggaaaaatt	tcaaaaattg	ggcctgaaaa	tccatacaat	actccagtat	3720
ttgccataaa	gaaaaaagac	agtactaaat	ggagaaaatt	agtagatttc	agagaactta	3780
ataagagaac	tcaagatttc	tggaaggttc	aattaggaat	accacatcct	gcagggttaa	3840
aacagaaaaa	atcagtaaca	gtactggatg	tggtgcgatc	atatttttca	gttcccttag	3900
ataaagactt	caggaagtat	actgcattta	ccatacctag	tataaacaat	gagacaccag	3960
ggattagata	tcagtacaat	gtgcttcac	agggatggaa	aggatcacca	gcaatattcc	4020
agtgtagcat	gacaaaaatc	ttagagcctt	ttagaaaaca	aaatccagac	atagtcattc	4080
atcaatacat	ggatgatttg	tatgtaggat	ctgacttaga	aatagggcag	catagaacaa	4140
aaatagagga	actgagacaa	catctgttga	ggtggggatt	taccacacca	gacaaaaaac	4200
atcagaaaaga	acctccattc	ctttggatgg	gttatgaact	ccatcctgat	aaatggacag	4260
tacagcctat	agtgtgccca	gaaaaggaca	gctggactgt	caatgacata	cagaaattag	4320
tggaagaaat	gaattgggca	agtcagattt	atgcagggat	taaagtaagg	caattatgta	4380
aacttcttag	gggaacaaaa	gcactaacag	aagtagtacc	actaacagaa	gaagcagagc	4440
tagaactggc	agaaaacagg	gagattctaa	aagaaccggt	acatggagtg	tattatgacc	4500
catcaaaaga	cttaatatga	gaaatacaga	agcaggggca	aggccaatgg	acatatcaaa	4560
tttatcaaga	gccatttaaa	aatctgaaaa	caggaaaaata	tgcaagaatg	aagggtgccc	4620
acactaatga	tgtgaaacaa	ttaacagagg	cagtacaaaa	aatagccaca	gaaagcatag	4680
taatatgggg	aaagactcct	aaattttaa	taccataaca	aaaggaaaca	tggaagcat	4740
ggtggacaga	gtattggcaa	gccacctgga	ttcctgagtg	ggagtgtgtc	aatacccttc	4800
ccttagtgaa	gttatgttac	cagttagaga	aagaaccat	aataggagca	gaaactttct	4860

-continued

atgtagatgg	ggcagccaat	agggaaacta	aattaggaaa	agcaggatat	gtaactgaca	4920
gaggaagaca	aaaagtgtgc	cccctaacgg	acacaacaaa	tcagaagact	gagttacaag	4980
caattcatct	agctttgcag	gattcgggat	tagaagtaaa	catagtgcac	gactcacaat	5040
atgcattggg	aatcatcaca	gcacaaccag	ataagagtga	atcagagtta	gtcagtcaca	5100
taatagagca	gttaataaaa	aaggaaaaag	tctacctggc	atgggtacca	gcacacaaag	5160
gaattggagg	aaatgaacaa	gtagatgggt	tggtcagtcg	tggaatcagg	aaagtactat	5220
ttttagatgg	aatagataag	gcccagaag	aacatgagaa	atcacacagt	aattggagag	5280
caatggctag	tgattttaac	ctaccacctg	tagtagcaaa	agaaatagta	gccagctgtg	5340
ataaatgtca	gctaaaaggg	gaagccatgc	atggacaagt	agactgtagc	ccaggaatat	5400
ggcagctaga	ttgtacacat	ttagaaggaa	aagttatctt	ggtagcagtt	catgtagcca	5460
gtggatatat	agaagcagaa	gtaattccag	cagagacagg	gcaagaaaca	gcatacttcc	5520
tcttaaaatt	agcaggaaga	tgccagctaa	aaacagtaca	tacagacaat	ggcagcaatt	5580
tcaccagtac	tacagttaag	gccgcctgtt	ggtggggcgg	gatcaagcag	gaatttggca	5640
ttccctacaa	tcccaaaagt	caaggagtaa	tagaatctat	gaataaagaa	ttaaagaaaa	5700
ttataggaca	ggtaagagat	caggctgaac	atcttaagac	agcagtacaa	atggcagtat	5760
tcaccacaaa	ttttaaaga	aaagggggga	ttggggggta	cagtgcaggg	gaaagaatag	5820
tagacataat	agcaacagac	atacaaaacta	aagaattaca	aaaacaaatt	acaaaaattc	5880
aaaattttcg	ggtttattac	agggacagca	gagatccagt	ttggaaagga	ccagcaaagc	5940
tcctctggaa	agggtgaagg	gcagtagtaa	tacaagataa	tagtgacata	aaagtagtgc	6000
caagaagaaa	agcaagatc	atcagggatt	atggaaaaca	gatggcaggt	gatgattgtg	6060
tggaagtag	acaggatgag	gattaacaca	tggaattctg	caacaactgc	tgtttatcca	6120
tttcagaatt	ggaggagctt	tgttccttgg	gttcttggga	gcagcaggaa	gcactatggg	6180
cgcagcgtca	atgacgttga	cgttacaggc	cagacaatta	ttgtctggta	tagtgcagca	6240
gcagaacaat	ttgctgagg	ctattgaggc	gcaacagcat	ctgttgcaac	tcacagtctg	6300
gggcatcaag	cagctccagg	caagaatcct	ggctgtggaa	agatacctaa	aggatcaaca	6360
gctcctgggg	atttgggggt	gctctggaaa	actcatttgc	accactgctg	tgccttggaa	6420
tgctagtgtg	agtaataaat	ctctggaaca	gatttggaat	cacacgacct	ggatggagtg	6480
ggacagagaa	attaacaatt	acacaagctt	gctagcagat	ctttttccct	ctgccccaaa	6540
ttatggggac	atcatgaagc	cccttgagca	tctgacttct	ggctaataaa	ggaaatttat	6600
tttcattgca	atagtgtgtt	ggaatttttt	gtgtctctca	ctcggaagga	catatgggag	6660
ggcaaatcat	ttaaaacatc	agaatgagta	tttggttttag	agtttggcaa	catatgccat	6720
atgctggctg	ccatgaacaa	aggtggctat	aaagaggcca	tcagtatatg	aaacagcccc	6780
ctgctgtcca	ttccttatcc	catagaaaag	ccttgacttg	aggttagatt	ttttttatat	6840
tttggtttgt	gttatttttt	tctttaacat	ccctaaaatt	ttccttacat	gttttactag	6900
ccagattttt	cctcctctcc	tgactactcc	cagtcatagc	tgtccctctt	ctcttatgaa	6960
gatccctcga	cctgcagccc	aagcttggcg	taatcatggg	catagctgtt	tctgtgtgta	7020
aattgttata	cgtccacaat	tccacacaa	atacgagccg	gaagcataaa	gtgtaaagcc	7080
tgggtgtcct	aatgagtggg	ctaactcaca	ttaattgcgt	tgcgctcact	gcccgttttc	7140

-continued

cagtcgggaa	acctgtcgtg	ccagcggatc	cgcattctcaa	ttagtcagca	accatagtcc	7200
cgcctcctaac	tccgcccatac	ccgcccctaa	ctccgcccag	ttccgcccatt	tctccgcccc	7260
atggctgact	aatttttttt	atttatgcag	aggccgaggc	cgcctcggcc	tctgagctat	7320
tccagaagta	gtgaggaggc	ttttttggag	gcctaggctt	ttgcaaaaag	ctaacttgtt	7380
tattgcagct	tataatgggt	acaaataaag	caatagcatc	acaaatttca	caaataaagc	7440
atttttttca	ctgcattcta	gttgtgggtt	gtccaaactc	atcaatgtat	cttatcatgt	7500
ctggatccgc	tgcattaatg	aatcgcccaa	cgcgcgggga	gaggcggttt	gcgtattggg	7560
cgtcttccg	cttctcgtc	cactgactcg	ctgcgtcgg	tcgttcggct	gcggcgagcg	7620
gtatcagctc	actcaaaggc	ggtaatacgg	ttatccacag	aatcagggga	taacgcagga	7680
aagaacatgt	gagcaaaagg	ccagcaaaag	gccagggaacc	gtaaaaaggc	cgcgttgctg	7740
gcgtttttcc	ataggctccg	ccccctgac	gagcatcaca	aaaatcgacg	ctcaagtcatg	7800
agggtggcga	acccgacagg	actataaaga	taccaggcgt	ttccccctgg	aagctccctc	7860
gtgcgctctc	ctgttccgac	cctgcgcgtt	accggatacc	tgccgcctt	tctcccttcg	7920
ggaagcgtgg	cgttttctca	atgctcacgc	tgtaggatc	tcagttcggg	gtaggtcgtt	7980
cgtcccaagc	tgggtctgtg	gcacgaaccc	cccgttcagc	ccgaccgctg	cgccttatcc	8040
ggtaactatc	gtcttgagtc	caaccggta	agacacgact	tatcgccact	ggcagcagcc	8100
actggttaaca	ggattagcag	agcgaggat	gtaggcgggtg	ctacagagtt	cttgaagtgg	8160
tggcctaact	acggctacac	tagaaggaca	gtatttggtg	tctgcgctct	gctgaagcca	8220
gttaccttcg	gaaaaagagt	tggtagctct	tgatccggca	aacaaaccac	cgtggttagc	8280
ggtggttttt	ttgtttgcaa	gcagcagatt	acgcgcagaa	aaaaaggatc	tcaagaagat	8340
cctttgatct	tttctacggg	gtctgacgct	cagtggaaag	aaaactcacg	ttaagggatt	8400
ttggtcatga	gattatcaaa	aaggatcttc	acctagatcc	ttttaaatta	aaaatgaagt	8460
tttaaatcaa	tctaaagtat	atatgagtaa	acttggtctg	acagttacca	atgcttaatc	8520
agtgaggcac	ctatctcagc	gatctgtcta	ttctgttcat	ccatagttgc	ctgactcccc	8580
gtcgtgtaga	taactacgat	acgggagggc	ttaccatctg	gcccagtgcc	tgcaatgata	8640
ccgcgagacc	cacgctcacc	ggctccagat	ttatcagcaa	taaaccagcc	agccggaagg	8700
gccgagcgca	gaagtggtec	tgcaacttta	tccgcctcca	tccagtctat	taattgttgc	8760
cgggaagcta	gagtaagtag	ttcgccagtt	aatagtttgc	gcaacgttgt	tgccattgct	8820
acaggcatcg	tggtgtcacg	ctcgtcgttt	ggtatggctt	cattcagctc	cggttcccaa	8880
cgatcaaggc	gagttacatg	atcccccatg	ttgtgcaaaa	aagcggttag	ctccttcggt	8940
cctccgatcg	ttgtcagaag	taagttggcc	gcagtgttat	cactcatggt	tatggcagca	9000
ctgcataatt	ctcttactgt	catgccatcc	gtaagatgct	tttctgtgac	tggtgagtac	9060
tcaaccaagt	cattctgaga	atagtgtatg	cggcgaccga	gttgcctctg	cccggcgcca	9120
atacgggata	ataccgcgcc	acatagcaga	actttaaaag	tgctcatcat	tggaacacgt	9180
tcttcggggc	gaaaaactctc	aaggatctta	cgcgtgttga	gatccagttc	gatgtaaccc	9240
actcgtgcac	ccaactgac	ttcagcatct	tttaacttca	ccagcgtttc	tggttgagca	9300
aaaacaggaa	ggcaaaatgc	cgcaaaaaag	ggaataaggg	cgacacggaa	atgttgaata	9360
ctcactactct	tccttttttca	atattattga	agcattttatc	agggttattg	tctcatgagc	9420

-continued

ggatacatat ttgaatgtat ttgaaaaat aaacaaatag gggttccgcg cacatttccc 9480

cgaaaagtgc cacctg

9496

What is claimed is:

1. A method of treating cells infected with HIV, the method comprising:

- (a) contacting peripheral blood mononuclear cells (PBMC) isolated from a subject infected with HIV with a therapeutically effective amount of a stimulatory agent, wherein the contacting is carried out ex vivo;
- (b) transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element, wherein the at least one genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5 or at least one small RNA capable of targeting an HIV RNA sequence; and
- (c) culturing the transduced PBMC for at least 1 day.

2. The method of claim 1, further comprising positively selecting HIV-specific CD4+ T cells from the PBMC.

3. The method of claim 2, wherein the HIV-specific CD4+ T cells are positively selected using at least one physical method of selection.

4. The method of claim 1, further comprising infusing the transduced PBMC into a subject.

5. The method of claim 4, wherein the subject is a human.

6. The method of claim 1, wherein the stimulatory agent comprises a peptide.

7. The method of claim 6, wherein the peptide comprises a gag peptide.

8. The method of claim 1, wherein the stimulatory agent comprises a vaccine.

9. The method of claim 8, wherein the vaccine comprises a HIV vaccine.

10. The method of claim 9, wherein the HIV vaccine comprises a MVA/HIV62B vaccine or a variant thereof.

11. The method of claim 1, wherein the at least one genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5 and at least one small RNA capable of targeting an HIV RNA sequence.

12. The method of claim 1 or 11, wherein the HIV RNA sequence comprises a HIV Vif sequence, a HIV Tat sequence, or a variant thereof.

13. The method of claim 1 or 11, wherein the at least one genetic element comprises a microRNA or a shRNA.

14. The method of claim 13, wherein the at least one genetic element comprises a microRNA cluster.

15. The method of claim 13, wherein the at least one genetic element comprises a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 1.

16. The method of claim 13, wherein the at least one genetic element comprises SEQ ID NO: 1.

17. The method of claim 13, wherein the at least one genetic element comprises a microRNA having:

- a. at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 2; or
- b. at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 3.

18. The method of claim 13, wherein the at least one genetic element comprises SEQ ID NO: 2; or SEQ ID NO: 3.

19. The method of claim 14, wherein the microRNA cluster comprises a sequence having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 31.

20. The method of claim 14, wherein the microRNA cluster comprises SEQ ID NO: 31.

21. A method of treating HIV infection in a subject, the method comprising:

- (a) immunizing the subject with an effective amount of a first stimulatory agent;
- (b) removing leukocytes from the subject and purifying peripheral blood mononuclear cells (PBMC);
- (c) contacting the PBMC ex vivo with a therapeutically effective amount of a second stimulatory agent;
- (d) transducing the PBMC ex vivo with a viral delivery system encoding at least one genetic element; and
- (e) culturing the transduced PBMC for at least 1 day.

22. The method of claim 21, further comprising positively selecting HIV-specific CD4+ T cells from the PBMC.

23. The method of claim 22, wherein the HIV-specific CD4+ T cells are positively selected using at least one physical method of selection.

24. The method of claim 21, further comprising infusing the transduced PBMC into the subject.

25. The method of claim 21, wherein the first and second stimulatory agents are the same.

26. The method of claim 21, wherein at least one of the first and second stimulatory agents comprises a HIV vaccine.

27. The method of claim 26, wherein the HIV vaccine comprises a MVA/HIV62B vaccine or a variant thereof.

28. The method of claim 21, wherein the viral delivery system comprises a lentiviral particle.

29. The method of claim 21, wherein the at least one genetic element comprises a small RNA capable of inhibiting production of chemokine receptor CCR5 or at least one small RNA capable of targeting an HIV RNA sequence.

30. A method of treating HIV infection in a subject, the method comprising:

- (a) immunizing the subject with an effective amount of a first stimulatory agent;
- (b) removing leukocytes from the subject and purifying peripheral blood mononuclear cells (PBMC);
- (c) positively selecting HIV-specific CD4+ T cells from the PBMC;
- (d) contacting the HIV-specific CD4+ T cells ex vivo with a therapeutically effective amount of a second stimulatory agent;
- (d) transducing the HIV-specific CD4+ T cells ex vivo with a viral delivery system encoding at least one genetic element; and
- (e) culturing the transduced PBMC for at least 1 day.

* * * * *